SAT0157 THE SEX HORMONE BINDING GLOBULIN IN MALE OSTEOPOROSIS

ER Legrand, C Gibert, Y Gallois, E Mathieu, F Boux de casson, MF Basle, D Chappard, M Audran. *Service de Rhumatologie, Centre Hospitalier Universitaire, Angers, France*

10.1136/annrheumdis-2001.644

Background Several studies have shown that the increase in Sex Hormone Binding Globulin and the decrease in bioavailable estradiol and testosterone can explain age-induced bone loss in healthy men over 70 years.

Objectives To evaluate the role of the Sex Hormone Binding Globulin (SHBG) in the pathophysiology of osteoporosis in middle-aged men.

Methods 80 men with osteoporosis (T-score < -2.5) and 40 age-matched controls were recruited to evaluate the relationships between sex hormone levels, bone markers levels, bone mineral density and vertebral fractures. Fasting serum samples were assayed for total and free testosterone, total estradiol and SHBG. Bone remodelling was evaluated by measurement of urinary levels of the C-telopeptide of type I collagen (CTX) and free deoxyypyridinoline (D-Pyr), serum osteocalcin and bone specific alkaline phosphatase (BSAP).

Results

- There was no difference between controls and osteoporotic men according to age, BMI, total testosterone and estradiol. In contrast, serum SHBG level was significantly higher (+42.2%) whereas free androgen index was lower (-24.8%) in patients with osteoporosis.
- Stepwise linear regression analysis showed that SHBG was significantly correlated with D-Pyr ($r = 0.33$), CTX ($r = 0.25$) and BSAP ($r = 0.30$). (3) After adjustment for age and BMI, hip BMD was not associated with total testosterone or estradiol but only with serum SHBG ($r = -0.35$). (4) Among osteoporotic patients, spinal radiographs evidenced at least one vertebral crush fracture in 36 men and none in 44 patients. Serum SHBG concentration was significantly associated with the presence of vertebral fracture: odds ratio was 2.0 (95% CI: 1.2–3.5) for an increase of one standard deviation of SHBG.

Conclusion The present study showed that serum SHBG concentration is increased in most middle-aged men with primary or secondary osteoporosis and is correlated with bone remodelling markers, hip bone mineral density and vertebral fracture risk. These results might represent a plausible explanation of bone loss in young and middle-aged men with idiopathic osteoporosis and vertebral fractures.

SAT0158 ASSOCIATES WITH AND EXTENT OF REDUCED BONE MINERAL DENSITY (BMD) IN FEMALE RHEUMATOID ARTHRITIS (RA) PATIENTS: THE OSLO, TRURO, AMSTERDAM (OSTRA) COLLABORATIVE STUDY

¹G Haugeberg, ²M Lodder, ³W Lems, ¹T Uhlig, ²B Dijkman, ¹T Kvien, ³A Woolf. ¹Department of Rheumatology, Diakonhjemmet Hospital, Oslo, Norway; ²Department of Rheumatology, Vrije Universiteit, Amsterdam, Netherlands; ³Department of Rheumatology, Royal Cornwall Hospital, Truro, UK

10.1136/annrheumdis-2001.645

Background

Objectives To examine associates with and occurrence of reduced BMD in female RA patients from three European countries.

Methods 150 (50 from each country) consecutive female RA patients (ACR 1987 criteria, disease duration ≥ 5 yrs, age 50–70 yrs), visiting the out-patient rheumatology clinics were included. Study variables included clinical data, laboratory tests, hand X-ray (Larsen score) and BMD measurement. BMD (g/cm^2) was measured by dual energy X-ray absorptiometry (DXA) and adjusted for age, weight and height. A pooled European/US reference population, showing no substantial differences with published Norwegian, English, and Dutch reference populations, was used to calculate T-scores. Groups were compared by one way ANOVA and cross tabulation (Pearson Chi square test). To investigate associates for BMD reduction bi- and multi-variety statistical analyses were performed.

Results No significant differences between the countries were found for age, disease duration and rheumatoid factor, whereas weight and most disease related variables, including the use of prednisolone and treatment for osteoporosis, differed between the 3 groups. Percentage of patients having osteoporosis (T-score < -2.5 SD) at femoral neck was 21.3% in Oslo, 8.3% in Truro, and 18.4% in Amsterdam ($P = 0.19$) and at the spine 32.0%, 18.4% and 12.0% ($P = 0.04$), respectively. Mean (SD) BMD values are shown in the Table 1. Using multivariate linear regression age, weight, total hand X-ray Larsen score, previous history of clinical fracture and ever use of DMARDs were independent associates for femoral neck BMD. For spine L2–4 weight, mean CRP last 12 months, ever use of DMARDs and current use of prednisolone were independent associates.

Abstract SAT0158 Table 1

	Oslo	Truro	Amsterdam	P-value
Fem. neck	0.80 (0.06)	0.86 (0.10)	0.83 (0.07)	0.001
Spine L2–4	1.06 (0.06)	1.12 (0.12)	1.09 (0.09)	0.002

Conclusion Differences in femoral neck and spine BMD might reflect differences in life styles factors, disease activity and disease management between the countries.

SAT0159 NON-VERTEBRAL FRACTURES AND ITS ASSOCIATES IN FEMALE RHEUMATOID ARTHRITIS (RA) PATIENTS: THE OSLO, TRURO, AND AMSTERDAM (OSTRA) COLLABORATIVE STUDY

¹G Haugeberg, ²M Lodder, ³W Lems, ¹R Ørstavik, ²B Dijkman, ¹T Kvien, ³A Woolf. ¹Department of Rheumatology, Diakonhjemmet Hospital, Oslo, Norway; ²Department of Rheumatology, Vrije Universiteit, Amsterdam, Netherlands; ³Department of Rheumatology, Royal Cornwall Hospital, Truro, UK

10.1136/annrheumdis-2001.646

Background

Objectives To examine non-vertebral fractures (N-VF) in RA patients, of similar age, disease duration and rheumatoid factor status recruited from three European countries, and their association with bone mineral density (BMD), demographic and disease related associates.

Methods 150 (50 from each country) consecutive female RA patients (ACR 1987 criteria, disease duration > 5 yrs, age 50–70 yrs) visiting the out-patient Department of Rheumatology in the participating centres were included. Data on N-VF occurring after age 25 and after RA diagnosis, demographics and measures of disease activity were collected. Dual energy X-ray