Methods 16 patients with RA according to the 1987 ACR criteria, of whom 5 were in complete clinical remission, 6 patients with PsA, and 2 CO were studied. MRI was performed with a 0.2T dedicated device (Artoscan, Esaote, Genova, Italy). The sequence used was a Spin Echo (TR/TE = 100/16 ms, matrix = 160 × 128, FOV = 50 × 150, slice thickness = 5 mm). A bolus of 0.2 ml/kg of gadopentetate dimeglumine was injected intravenously and fast images of three slices of the wrist were repeated every 20 s thereafter. The enhancement ratio was calculated both as rate of early enhancement (REE) and as relative enhancement (RE). The signals from a small circular ROI in the region of maximal enhancement and from a larger segmented region including most of the enhanced synovial membrane were evaluated. The signal was normalised to the bone to reduced noise.

Results Both REE and RE were significantly higher in active RA vs. inactive RA ($p = 0.003$), PsA ($p = 0.002$), and CO ($p = 0.0002$). In addition, RE and REE were higher in inactive RA than in CO ($p = 0.008$). Results from the analyses using the small ROI and the segmented ROI showed a good correlation (mean $R = 0.75$). REE resulted slightly more sensitive than RE.

Conclusion Dynamic MRI is a useful method to evaluate synovitis by allowing quantitative estimation of the synovial enhancement. This promising technique is able to differentiate active from inactive RA, as well as RA from PsA and CO. The evaluation of a small area of the synovial membrane seems to be equally effective than the more time-consuming segmentation.

REFERENCES

- 1 Cimmino MA, et al. *Semin Arthritis Rheum.* 2000;**30**:180–95
- 2 Klarlund M, et al. *Scand J Rheumatol.* 2000;**29**:108–15

SAT0207 DETERMINATION OF SYNOVIAL INFLAMMATION IN THE WRIST BY DOPPLER ULTRASONOGRAPHY AND MRI IN PATIENTS WITH ARTHRITIS

¹L Terslev, ²A Savnik, ¹E Qvistgaard, ³S Torp-Pedersen, ¹B Danneskiold-Samsøe, ⁴HS Thomsen, ¹H Bliddal. ¹Parker Institute; ²Department of Radiology, Frederiksberg Hospital; ³Department of Radiology, University Hospital at Gentofte; ⁴Department of Radiology, University Hospital at Herlev, Copenhagen, Denmark

10.1136/annrheumdis-2001.704

Background Synovial membrane volumes obtained by Magnetic Resonance Imaging (MRI) have been suggested as a marker of joint disease activity.¹ Graded power Doppler images has been able to detect treatment response in joint synovium.² If ultrasonographic (US) parameters correspond with findings in MRI an alternative method in assessing inflammatory activity would be available.

Objectives To compare the evaluation of signs of synovial activity in patients with arthritis as determined by MRI and DUS respectively.

Methods 11 patients (5 women and 6 men, median age 52,6 years, range (32–69)) with arthritis were included. All patients were examined by DUS with a 15 MHz linear transducer and by 1.5 T MRI before and after contrast injection. The area and thickness of the synovial membrane were outlined after contrast injection on MRI. US flow pattern of the synovium in the wrist was evaluated by quantitative spectral DUS ($n = 10$), recording resistance index (RI). The relative RI (rRI) = meanRI/extra synovial RI, was calculated to adjust for changes in peripheral resistance unrelated to the joint. The synovial vascularisation was determined by colour Doppler ($n = 10$). Using a colour recognition function, all marked pixels were changed to a reference

colour and the total amount of pixels finally depicted using a colour histogram.³

Each patient was evaluated by ESR, CRP and joint assessment.

Results The colour Doppler pixel (CDP) measurements correlated significantly with both max area and thickness of the synovial membrane on post contrast MR images ($R_s = 0.88$ and 0.84 , $P < 0.002$). Also the rRI values correlated significantly with both max area and thickness of the synovial membrane on post contrast MR images ($R_s = 0.89$ and 0.93 , $p < 0.001$). Kappa statistics were made on MRI and DUS: Kappa was 0,8 between the max area and thickness of the synovial membrane on post contrast MR images and the CDP measurements. Kappa was 1,0 between the max area and thickness of the synovial membrane on post contrast MR images and the rRI values. No correlation was found between ESR, CRP and joint assessment and the imaging modalities.

Conclusion The estimation of synovial inflammatory activity on DUS was comparable with post contrast MRI. DUS appears to be an alternative method to determine inflammatory activity in patients with rheumatic diseases.

REFERENCES

- 1 Ostergaard M, et al. Magnetic resonance imaging determined synovial membrane volume as a marker of disease activity and a predictor of progressive joint destruction in the wrists of patients with rheumatoid arthritis. *Arthritis Rheum.* 1999;**42**(5):918–29
- 2 Newman JS, et al. Power Doppler sonography of synovitis: assessment of therapeutic response preliminary observations. *Radiology* 1996;**198**:582–4
- 3 Qvistgaard E, et al. Quantitative ultrasonography in rheumatoid arthritis evaluation of inflammation by Doppler technique. Submitted

SAT0208 THE PREDICTIVE ROLE OF BONE SCINTIGRAPHY IN OSTEOARTHRITIS, RELATIONSHIP WITH AGE AND SEVERITY OF THE DISEASE

¹S Bal, ²H Erturk, ³M Aydin, ³O Ugur, ¹R Celiker. ¹Physical Medicine and Rehabilitation; ²Physical Medicine and Rehabilitation, State Hospital, Nevsehir, Turkey; ³Nuclear Medicine, Hacettepe University Medical School, Ankara

10.1136/annrheumdis-2001.705

Background Osteoarthritis (OA) is the most common cause of musculoskeletal diseases in the developed countries. The diagnosis of OA is usually dependent on clinical and radiological evaluations which have some limitations in measuring the outcome.

Objectives The aims of this study were to evaluate the predictive role of bone scintigraphy in OA, and to study the relationship with age and severity of the disease.

Methods Ninety-nine female subjects were included in this study. Conventional x-rays of the hand, knee, hip and lumbar regions were obtained from each subjects. According to Kellgren and Lawrence Grading System the radiological score of the joint which had the highest grade was accepted as the Maximum Radiological Score (MRS). A bone scan was performed for each subjects 4 h after the injection of 740 MBq of technetium ^{99m}-MDP. Spot images of hand, knee, lumbar vertebrae and pelvis were acquired using a gamma camera (Toshiba GCA 601). Besides the increased or normal uptake of activity at the affected joints, the ratios of lesion/non-lesion were also calculated. In the areas where the increased uptake of activity were present, the maximum lesion/non-lesion ratio was accepted as Maximum Scintigraphic Score (MSS). According to the American College of Rheumatology- clinical and radiological ? criteria for diagnosis of OA 46 patients were included in OA group (Group 1) and

53 patients were included in control group (Group 2). The Lequesne Index was used to determine the severity of the disease.

Results All patients in group 1 and 31 patients in group 2 had knee pain and the mean duration of pain was 51.6+77.5 months in group 1 and 23.4+46.5 months in group 2. The mean Body Mass Index (BMI) was 31.7+5.0 in group 1 and 36.7+4.0 in group 2. According to the results of scintigraphic evaluation, healthy controls were separated into two groups. In group 2A patients who had an increase in uptake at least one joint and in group 2B patients who had no uptake. The mean MSS was 4.2 +1.4 in group 1 and 3.3+1.6 in group 2A. Also the mean MRS of the patients in group 1 were 2.9+0.9 and 0.6+0.5, 0.2+0.4 in group 2A and 2B respectively. The mean score of the Lequesne Index was 8,9+3,6 in group 1 patients. There was a positive correlation between age and MSS ($p < 0,05$) and BMI and MRS ($p < 0,05$). Also there was a strong correlation between Lequesne OA Severity Index Score the age of the patients in group 1 but no correlation with the MSS. According to the clinical and/or radiological criteria sensitivity of bone scintigraphy was 98.2%, specificity was 57.1%, positive predictive value was 76% and negative predictive value was 96%.

Conclusion Our results suggest that 99 m technetium scintigraphy is a better quantifying method in assessing the changes of OA especially in early cases.

SAT0209 MRI EVALUATION OF CHONDRAL LESIONS IN KNEE OSTEOARTHRITIS: ARTHROSCOPIC CORRELATION

¹C Acebes, ²A Robledo, ³J Granizo, ¹G Herrero-Beaumont. ¹Rheumatology; ²Radiology; ³Epidemiology, Fundación Jiménez Díaz, Madrid, Spain

10.1136/annrheumdis-2001.706

Background Arthroscopic exam evaluation (chondroscopy) of the articular cartilage is considered a relevant instrument of OA outcome for research purposes. Nevertheless availability of other accurate but non-invasive methods for chondral evaluation in OA, such as MRI, should be desirable.

Objectives To evaluate the capacity of MRI to score chondropathy in knee OA compared with chondroscopy.

Methods 24 consecutive patients diagnosed of knee OA were referred for knee hyaline cartilage MR imaging of medial chondile and tibial plateau with a fat-suppressed spoiled gradient-echo sequences (SPGR) before chondroscopy. MRI and chondroscopic cartilage lesions evaluated by two methods were registered in an scale diagram and total score chondropathy was calculated using a software programme (Autocad). Correlation was determined by Sperman's test for both evaluation ($p < 0,05$).

Results High and significant intraobserver correlation was observed for evaluation of medial chondile and tibial plateau by chondroscopy $r = 0,806$ and $r = 0,721$ and by MRI $r = 0,805$ and $r = 0,680$ respectively. Correlation coefficient between chondroscopic and MRI score for chondile and tibia was $r = 0,781$ and $r = 0,531$ respectively.

Conclusion Fat-suppressed SPGR imaging seems to be a relevant tool for evaluation of cartilage lesions in OA of the knee showing a high correlation between both procedures. Both chondroscopy and MRI have a lower correlation for the detection of chondral lesions in tibial plateau than in femoral chondile.

Supp. by a grant of Health Research Funding (98/626).

SAT0210 BRAIN MAGNETIC RESONANCE IMAGING (MRI) AND PROTON MAGNETIC RESONANCE SPECTROSCOPY (1H-MRS) IN SYSTEMIC LUPUS ERYTHEMATOSUS (SLE)

¹M Govoni, ²P Colamussi, ¹N Rizzo, ¹D Santilli, ³R Ricci, ¹F Trotta. ¹Sez. Di Reumatologia; ²Ist. Di Medicina Nucleare, Università Di Ferrara, Ferrara; ³Sevizio Di Neuroradiologia, Ospedale Bellaria, Bologna, Italy

10.1136/annrheumdis-2001.707

Background

Objectives We evaluated the usefulness of 1H-MRS in patients with SLE, with or without neurological involvement, in whom single photon emission computed tomography (SPECT) showed cerebral perfusion abnormalities in brain areas with no detectable changes at MRI.

Methods 9 female SLE patients, with a mean age of 28.7 years (range 14–46) and a mean duration of disease of 2.5 years, underwent cerebral blood flow (CBF) study with 99mTc-SPECT and brain MRI with 1H-MRS analysis. 4 patients complained of headache, 1 patient was being treated for partial epilepsy and 4 patients were asymptomatic and with a negative neurological examination. In brain areas shown to be hypoperfused by SPECT, peak levels of signals from brain metabolites N-acetylaspartic acid (NAA), choline (CHO), and creatine (CRE) were measured on a total of 20 brain voxels (volume = 8 cm³) selected on transverse MR images. Values were expressed as NAA/CHO, CHO/CRE and NAA/CRE ratios. Ratios measured in 9 hypoperfused brain areas were compared with those obtained in 11 normoperfused areas. MRI showed a normal appearance in all of the above brain areas. Statistical analysis was performed by means of classical t-test.

Results NAA/CHO ratio was decreased in 78% of hypoperfused and in 18% of normoperfused brain areas (1.70 ± 0.15 vs 2.55 ± 0.16 ; $p = 0.0017$). CHO/CRE ratio was increased in 89% of hypoperfused and in 9% of normoperfused brain areas (1.34 ± 0.08 vs 0.80 ± 0.05 ; $p < 0.0001$). No significant differences were detected in NAA/CRE ratios. The comparison of spectroscopic ratios measured both in hypoperfused and normoperfused areas in patients with neurological symptoms with those obtained in asymptomatic patients did not show any significant differences.

Conclusion In SLE patients, independently from the presence of overt clinical neurological involvement, 1H-MRS is a valuable tool to detect neuronal biochemical changes in brain regions with reduced CBF at SPECT. Although the precise meaning of these abnormalities remains to be determined, their localization in areas with reduced CBF suggests that they could represent an early sign of neuronal injury in SLE patients even in the presence of a normal brain MRI appearance. If proven so, 1H-MRS could be considered a sensitive and accurate measure of neuronal dysfunction in SLE and this would demonstrate that the detection of reduced CBF by SPECT in areas with a normal MRI appearance is not to be regarded as a false positive result.

SAT0211 INTEROBSERVER AGREEMENT OF ULTRASOUND EXAMINATION IN ASSESSMENT OF THE SMALL JOINTS OF THE HANDS AND FEET IN RHEUMATOID ARTHRITIS

¹M Szkudlarek, ²M Court-Payen, ¹S Jacobsen, ³H Thomsen, ⁴M Ostergaard. ¹Department of Rheumatology; ²Department of Ultrasound; ³Department of Radiology, Herlev Hospital, Copenhagen, Denmark; ⁴The Danish Research Center of Magnetic Resonance, Hvidovre Hospital, Hvidovre

10.1136/annrheumdis-2001.708