therapy (1 gr/kg in three divided doses) was administered each month for six months.

Marked improvement in dermal eruptions was observed following the second administration of IVIG therapy. Now in the eighth month of follow up she has been complaining of dyspnea on exertion for twenty days. High resolution computerised tomography (HRCT) of lung revealed paranchimal changes suggesting that bronchiolitis obliterans organising pneumonia (BOOP).

Could this be the result of methotrexate therapy or a result of amyopathic dermatomyositis? Lung involvement in amyopathic dermatomyositis is rare. Pulmonary drug toxicity is increasingly being diagnosed as a cause of acute and chronic lung disease. BOOP, which is commonly caused by cyclophosphamide and bleomycin as well as methotrexate and gold salts, appears on HRCT scans as poorly defined nodular areas of consolidation, centrilobular nodules, and bronchial dilatation. Knowledge of the drugs most frequently involved can facilitate diagnosis and institution of appropriate treatment. The results of lung biopsy will be discussed in poster session. If lung biopsy will reveal the pathology methotrexate will be discontinued.

Objectives

Methods

Results

Conclusion

AB0219 QUANTITATIVE HEMOSTATIC INDICES IN SECONDARY AND MYELOPROLIFERATIVE THROMBOCYTOSIS

¹J Parada-Turska, ²B Sokolowska, ²A Walter-Croneck, ²A Dmoszynska, ¹L Szczepanski, ²J Brzozowska, ²M Sledzinska, ²M Kokoszka. ¹Department of Rheumatology; ²Department of Hematology, Medical University, Lublin, Poland

10.1136/annrheumdis-2001.740

Background An elevated platelet count may lead to an increased incidence of systemic thromboembolism or unexpectedly to thrombohemorrhagic complications.

Both, incidences of systemic thromboembolism and unexpected thrombohemorrhagic complications (bleeding) were described in patients with an elevated platelet count.

In the present study the quantitative hemostatic indices were measured in 46 patients with thrombocytosis (platelet count > 400.000 per microliter), including 23 patients with secondary thrombocytosis (17 patients with rheumatoid arthritis, 2 with lung cancer, 1 with systemic lupus erythematosus, 1 with granuloma Wegeneri) and 23 patients with myeloproliferative thrombocytosis (12 with chronic myeloid leukaemia, 8 with polycythemia vera, 3 with idiopathic thrombocytosis).

In both studied groups, von Willebrand factor activity and epinephrine-induced platelet aggregation were decreased, and activated partial thromboplastin time was prolonged, thus, suggesting increased risk of bleeding complications.

On the other hand, the concentrations of fibrinogen and d-dimers were enhanced and bleeding time was shortened. These findings argue for an enhanced risk of thrombosis.

Since no differences between two studied groups were found, it seems that the observed abnormalities were directly related to the elevated platelet count and were not associated with primary disease.

The low incidence of hemostatic complications in patients with secondary and myeloproliferative thrombosis may suggest that the coexistence of an additional component is critical for the clinical manifestation of such disturbances.

Objectives

Methods

Results

Conclusion

AB0220 OCHRONOTIC SPONDYLOARTHROPATHY WITH MULTISYSTEMIC INVOLVEMENT: A CASE REPORT

K Nas, A Gür, R Çevik, AJ Saraç, S Söker (Çakmak), S Em, M Karakoç, S Kiliç. *Physical Medicine and Rehabilitation, Dicle University, Diyarbakir, Turkey*

10.1136/annrheumdis-2001.741

Background Ochronosis is a musculoskeletal manifestation of alcaptonuria, a rare hereditary metabolic disorder characterised by the absence of the enzyme homogentisic acid oxidase and associated with various systemic abnormalities related to the deposition of homogentisic acid pigment.

Objectives We aimed to review a case with ochronosis, a disorder of rare and multisystemic involvement, in the light of literature.

Methods A housewife, 53 years of age, was admitted to our clinic with complaints of low back and left shoulder pains. Due to low-back pain complaints, she was operated on two times with diagnosis of lumbar discal hernia.

Results The radiographic appearance in the lumbar spine of the patients was almost pathognomonic of ochronotic spondyloarthropathy. Typical water-like calcifications were seen in the intervertebral discs, with narrowing of disc spaces and osteoporotic rarefaction of the vertebral bodies. Shoulder joint, narrowing of the joint space, a marked subchondral sclerosis, and small osteophytes were observed; similarly, knee, sacroiliac joints and symphysis pubis show at narrowing of the joint spaces and subchondral bone sclerosis.

In the eyes of the patient, ochronotic pigmentation was seen by light microscopy. The blue-black pigmentation was bilateral and asymmetric in the cornea, conjunctiva and sclera. In addition, black-grey pigmentation was present in the ear cartilage and hand nails. In the laboratory test, homogentisic acid was found increase (2740 mmol HJA/mmol creatinin; N: 10 mmol HJA/mmol) creatinin in urine.

Conclusion

AB0221 FAMILIAL MEDITERRANEAN FEVER: ANALYSIS OF 45 TUNISIAN PATIENTS

SS Turki, CH Ben Taarit, R Goucha, E Abderrahim, F Ben Moussa, H Ben Maiz. *Department of Nephrology and Internal Medicine, Charles Nicolle Hospital, Tunis, Tunisia*

10.1136/annrheumdis-2001.742

Background Familial mediterranean fever (FMF) is an inherited multisystem disease of unknown aetiology, characterised by recurrent, painful, self-limited episodes.

Objectives The objectives of this study is to describe clinical profile, course and complications of 45 cases of FMF observed over a period of 25 years.

Methods Retrospective review of 45 cases with FMF.

Results Forty patients (88%) started their illness below the age of 10 years. Peritonitis occurred in 84,4%, arthritis in 35,5%, pleuritis in 17,7% and erysipelas-like lesions in 31,1%. 21 patients developed renal amyloidosis and 17 patients were subjected to unnecessary operative surgery.

The arthritis was monoarticular in 85% and polyarticular in 15% and was seronegative in all patients (Rheumatoid factor and antinuclear antibodies). The synovial attack showed a wide variation in the clinical presentation, course and duration of arthritis causing diagnostic difficulties. Renal amyloidosis occurred in 21 patients after a mean duration of FMF 10,5 years (2 to 28 years). One patient had hematuria, kidney biopsy showed mesangial IgA glomerulonephritis. Among 32 patients treated with colchicine, 80% achieved a therapeutic response.

Conclusion FMF is an inherited disease; colchicine remains the only efficient treatment to prevent both acute manifestations and amyloidosis.

AB0222 EVALUATION OF THE VISUAL ANALOGICAL SCALES (VAS) OF PAIN AND THEIR CORRELATION WITH THE OTHERS ACR ACTIVITY CRITERIA OF RHEUMATOID ARTHRITIS (RA)

C Diaz, M Cortes, C Geli, M Moreno, H Corominas, C Diaz-Torne, A Rodriguez, JM Llobet, G Vazquez. *Rheumatology, Hospital de La Santa Creu I Sant Pau, Barcelona, Spain*

10.1136/annrheumdis-2001.743

Background

Objectives Evaluate the VAS for global, activity, rest and nocturnal pain in relation with other activity parameters including Ritchie index.

Methods 228 outpatients with ACR-87 criteria for RA diagnostic were studied. We measure the following clinical parameters: n° of joints with pain and inflammation, VAS for global, activity, rest and nocturnal pain, VAS for global impression of the patient and the doctor, HAQ, morning stiffness, Ritchie index and the analytical data: ESR and CRP. The correlations among the different pain-VAS and those with other activity criteria of RA were evaluated by Pearson's correlation with the statistical packet SPSS. They are positive if $r > .5$.

Results 82% were women. Age was 60 ± 16 . Years of evolution were 13 ± 10 . 31%, 28%, 29% and 12% correspond to the functional stages I, II, III and IV respectively. 48% were seropositive. VAS-G has the highest correlations with VAS-A, R and N. there are also significant correlation among VAS-Pain and other activity parameters of RA: HAQ, Ritchie index and the global impression of the patient and the doctor. The impression of the patient correlations are higher than the doctor's.

Conclusion VAS-Pain, and specially VAS-G has a high correlation among themselves, HAQ, patient and doctor's impression and with Ritchie Index. There are no correlation among n° of articulations with pain, articulations with inflammation, CRP or ESR. The lack of the last correlation maybe related to the low inflammatory activity of the sample.

AB0223 EVALUATION OF CLASSICAL NSAIDS AND COX-2 SELECTIVE INHIBITORS ON PURIFIED OVINE ENZYMES AND HUMAN WHOLE BLOOD

¹X De Leval, ¹JM Dogne, ¹J Delarge, ²JY Reginster, ²Y enroin. ¹Department of Medicinal Chemistry; ²Bone and Cartilage Metabolism Research Unit, Institute of Pathology, CHU Start-Tilman, Liège, Belgium

10.1136/annrheumdis-2001.744

Background

Objectives In this study, we evaluated the cyclooxygenases (COX) inhibitory profile of NSAIDs in a cellular assay using human whole blood.

Methods Human whole blood assay was performed using fresh blood from normal volunteers. COX-1 assay: Fresh blood was collected into tubes containing no anticoagulants, vortexed and incubated at 37°C under constant agitation for 1 h to allow the blood clot. The serum was then obtained by centrifugation and was assayed for thromboxane B₂ (TXB₂) using a enzyme immunoassay. COX-2 assay: Fresh blood was collected in heparinized tubes and activated by treatment with lipopolysaccharide at the final concentration of 100 µg/ml. After 24 h of incubation with the drugs, plasma was collected and assayed for prostaglandin E₂ (PGE₂). PGE₂ production was quantified by a specific radioimmunoassay. Drugs were tested at the concentrations ranged among 0.01 to 100 µmol/l.

Results

Abstract AB0223 Table 1

	COX-1	IC50 (µM) COX-2	COX-1/COX-2
Acetofenac	> 100	0.77	> 130
NS-398	8.3	0.34	24
Rofecoxib	7.8	0.43	18
Nimesulide	22	1.3	17
Celecoxib	12	0.77	16
Diclofenac	0.6	0.04	15
Tenoxicam	7.2	5.6	1.3
Indomethacin	0.07	0.17	0.4

Conclusion In the whole blood test and based on the IC50 values, the majority of the NSAID tested preferentially inhibits COX-2. Acetofenac shows the best selectivity whereas indomethacin preferentially inhibited COX-1.

OP0122 TUMOUR NECROSIS FACTOR-RECEPTOR ASSOCIATED PERIODIC SYNDROME IN A FINNISH FAMILY

¹T Pettersson, ²H Nevala, ²L Karenko, ²S Stjernberg, ²M Raatikainen, ²H Suomalainen, ³J Rauta, ⁴MF McDermott, ³P Peterson, ²A Ranki. ¹Department of Medicine; ²Department of Dermatology, Helsinki University Central Hospital, Helsinki; ³Institute of Medical Technology, University of Tampere, Tampere, Finland; ⁴Medical Unit, St. Bartholomew's and the Royal London Hospital School of Medicine and Dentistry, London, UK

10.1136/annrheumdis-2001.745

Background TNF receptor-associated periodic syndrome (TRAPS) is a recently defined dominantly inherited autoinflammatory syndrome, which is caused by mutations in the extracellular domains of the 55 kDa TNF receptor, TNFRSF1A. TRAPS is characterised by periodic fevers, sterile peritonitis, pleurisy, arthritis, erysipeloid erythema of the skin, and conjunctivitis. Affected individuals have shown a shedding defect of TNFRSF1A upon stimulation and, consequently, low levels of soluble TNFRSF1A in serum.

Objectives We studied a three-generation Finnish family, where recurrent attacks of fever and abdominal pain appeared to segregate as an autosomal dominant trait.

Methods The TNFRSF1A gene was sequenced in both affected and unaffected family members. Flow cytometric analysis and