Methods The total population of the city is 1,5 million population in the city of Yekaterinburg. The medical records of four city hospitals where patients with hip fractures were admitted as well as from outpatient emergency units were analysed for 1992–1997. The city administration provided statistical data on relevant population number and age distribution. Hip fracture has been defined as a fracture of the femoral neck or intertrochanteric region fracture (ICD IX code 820.0 and 820.2). There were no exclusion criteria other than the hip fracture occurring as a result of major trauma.

Results In the health care areas studied there were 2 777 hip fractures: 599 males and 2 178 females. The mean age of males was $69,0 \pm 11,2$ years of age and females – $76,5 \pm 9,3$ respectively ($p < 0,0001$). The incidence of hip fracture for the examined period made up 132,3/100 000 population in those over 50 years (76,9/100 000 for males and 164,8/100 000 for females). The ratio of male to females cases was 1:2. The incidence of hip fracture rose exponentially with age in both genders. The occurrence of hip fracture was maximal in subjects over 80 years old and made up 744,1/100 000 (476,9/100 000 males and 818,2/100 000 females). Male/female ratio was 1:1,7. The incidence of hip fracture in our population has not seasonal variation.

Conclusion More recent data showed significant differences in the incidence of hip fractures between Russian regions. The lowest rate of hip fracture was found in the western part of Russia and the highest in the Ural (Yekaterinburg). There was a large difference in incidence between regions, which suggests important environmental factors in the causation of hip fracture.

OP0108 AGGRESSIVE LYMPHOMAS (DIFFUSE LARGE B CELL LYMPHOMAS) ARE OVERREPRESENTED IN RA PATIENTS

¹E Baecklund, ²C Sundström, ³A Ekbom, ³N Feltelius, ³L Klareskog. ¹Department of Rheumatology; ²Department of Pathology, University Hospital, Uppsala; ³Department of Medicine, Karolinska Hospital, Stockholm, Sweden

10.1136/annrheumdis-2001.780

Background Patients with rheumatoid arthritis (RA) have an increased risk to develop lymphomas. In a previous case-control study we found a strong association between high disease activity in RA and the risk of lymphoma.

To get a better understanding of the biological background of this connexion we examined lymphoma subtypes in RA patients. **Objectives** To study the distribution of lymphoma subtypes in RA patients compared to the general population.

Methods In a population-based cohort of 11,683 patients with RA in the Uppsala Health Care region in Sweden, 42 cases of lymphoma were identified through record linkages with the Swedish cancer registry 1965 through 1984.

The medical records and paraffin embedded lymphoma tissues were collected.

The lymphomas were reclassified according to the recently described REAL classification. The distribution was compared to lymphoma subtypes in the general population according to the Non-Hodgkin's Lymphoma (NHL) Classification Project.¹

Results One patient was excluded as the RA diagnosis was wrong. Tissues from 36 of the remaining patients were found and reviewed. NHL was found in 33 patients, Hodgkin's disease in 2 and 1 patient did not have lymphoma.

Compared with the expected there was a more than 2-fold increased number of diffuse large B-cell lymphoma (22 patients, 69% of NHL's compared with 31% expected).

Few of the patients had been treated with immunosuppressive drugs.

Conclusion The distribution of lymphoma subtypes in RA patients differs from the expected.

This supports the idea of a specific underlying mechanism behind lymphoma development in RA patients. This mechanism may also be of importance for RA development and activity. Genetics and other characteristics of those RA patients who develop lymphomas deserve further studies.

REFERENCE

- 1 The Non-Hodgkin's Lymphoma Classification Project: A Clinical Evaluation of the International Lymphoma Study Group. *Blood* 1997;**89**(11):3909–18

OP0109 HIGHER INCIDENCE OF THROMBOEMBOLIC EVENTS AMONG PATIENTS WITH RHEUMATOID ARTHRITIS VS. OSTEOARTHRITIS, AND VS. NO ARTHRITIS, IN THE GENERAL PRACTICE RESEARCH DATABASE (GPRD)

DJ Watson, T Rhodes. *Epidemiology, Merck Research Laboratories, West Point, USA*

10.1136/annrheumdis-2001.781

Background Persons with Rheumatoid arthritis (RA) have been noted to have increased all-cause and cardiovascular (CV) disease mortality relative to those without RA. Data on mortality in patients with RA vs those with osteoarthritis (OA), and data on nonfatal + fatal CV incidence in patients with RA relative to those with OA and those with no arthritis are lacking.

Objectives To compare the incidence of all cause mortality, myocardial infarction (MI), sudden death (SD), cerebrovascular events (stroke; excluding transient ischaemic attack), CV death (fatal MI, sudden death, fatal stroke), and all thromboembolic events in RA patients compared with patients with OA but not RA, and compared with those with no arthritis, in the GPRD.

Methods The GPRD is a general practice research database that currently contains over 35 million patient years of data from the UK and represents 6% of general practice patients in England and Wales. This was an observational cohort study of patients >40 years old from GPRD practices in which >80% of patients had recorded visits. Patients with history of MI or stroke prior to beginning of follow-up were excluded. Person-time was calculated for each patient by gender, age (5 year intervals) and diagnosis (RA; OA but not RA, and no arthritis). The rate/1000 person-years was computed for each gender, age, and diagnosis stratum. For each patient, only the first incident endpoint was counted and attributed to the stratum in which it occurred. Gender and age-adjusted incidence rates and ratios of rates of each endpoint during person-time with RA vs that with OA but not RA, and vs that with no arthritis were computed using Poisson regression.

Results 594 practices contributed 6,533,474 patients to the analysis. The incidence of RA was 0.68/1000 patient years in men and 1.33/1000 patient years in women. Age-adjusted rates of all endpoints were significantly increased for both men and women with RA compared to those with OA but not RA (except for sudden death, which was significantly increased in women only). Age- and gender-adjusted rate ratios (95% CI) were: all-cause mortality 1.72 (1.67, 1.78); MI 1.32 (1.24, 1.41); SD 1.33 (1.09, 1.63); stroke 1.26 (1.18, 1.34); CV death 1.41 (1.24, 1.61); and all thromboembolic events 1.31 (1.25, 1.39). Similar results were seen when patients with RA were compared with those with no arthritis: age- and gender-adjusted rate ratios (95% CI) with RA vs no arthritis were: all-cause mortality 1.60