REFERENCES

- 1 Havelin LI, Espehaug B, Vollset SE, Engesaeter LB, Langeland N. The Norwegian Arthroplasty Register. A survey of 17,444 total hip replacements. *Acta Orthop Scand*. 1993;**64**:245–51
- 2 National Health Screening Service, The cardiovascular disease study in Norwegian counties: results from second screening. Oslo: National Health Screening Service, 1988
- 3 Waaler HT. Height, weight and mortality. *Acta Med Scand Suppl*. 1984;**679**:1–56

THU0240 EPIDEMIOLOGY OF OSTEOARTHRITIS IN NORTHERN ETHNIC GROUPS OF RUSSIA

¹S Erdesz, ²R Protopopova, ²V Krivoschapkin. ¹Epidemiology, Institute of Rheumatology, Moscow; ²Institute of Health, Yakutsk, Russia

10.1136/annrheumdis-2001.784

Background Till present there are practically no information on the prevalence and risk factors of osteoarthritis in populations residing in severe climatic zones of Russia and subject to the influence of a number of unfavourable environmental factors.

Objectives To study the epidemiology of osteoarthritis among native population of rural areas of Siberian North – in Yakutia.

Methods Cross-sectional study of residents of four Yakutian settlements (North/North-Eastern part of Siberia) was performed. At the moment of the study 1282 persons aged 18 and more resided lived there. Among them 1216 persons (94.8%) were examined. 1002 persons were Yakutians (Saha) and 214 – Evenks.

Results Osteoarthritis was diagnosed in 218 settlement residents. Correspondingly the prevalence of OA in the population under study was 18.00.8%. Among women the OA prevalence was 22.51.4% and among men 12.61.4% ($p < 0.05$). There was no reliable difference in prevalence between Yakuts and Evenks (correspondingly 17.3% and 21.0%, $p < 0.05$). We found linear correlation ($r = 0.96$) between frequency of OA and age.

Conclusion Thus, epidemiological study of osteoarthritis among rural population of Russian North demonstrated that the prevalence of the disease is 18.0% and women prevail and there is direct age relation between age and frequency of the disease.

THU0241 BEHCET'S DISEASE: DOES LACK OF KNOWLEDGE RESULT IN UNDERDIAGNOSIS?

¹Y Yazici, ¹D Erkan, ²G Neshet, ³E Kural, ³N Seyahi, ⁴A Ince. ¹Rheumatology, Hospital for Special Surgery, New York, USA; ²Rheumatology, Shaare-Zedek Medical Center, Jerusalem, Israel; ³Internal Medicine, Cerrahpasa Medical Faculty, Istanbul, Turkey; ⁴Rheumatology, St Louis University Health Sciences Center, St Louis, USA

10.1136/annrheumdis-2001.785

Background Prevalence of Behcet's disease has been estimated at 8–38/10,000 in Turkey and 1–2/30,000 in the USA. Pathergy test positivity is a major criterion for making the diagnosis of Behcet's disease.

Objectives To determine the knowledge among internal medicine (IM) residents from 2 different Mediterranean regions and 2 different regions in the USA regarding general rheumatology (GR), Behcet's disease, pathergy test and PPD test for tuberculosis.

Methods A 26-item questionnaire was developed including both closed and open-ended items, testing knowledge of GR, Behcet's disease, pathergy test, and PPD test. It was given to IM residents from 4 university centres (St. Louis University Health Sciences Centre (A) and Weill Medical College of Cornell University (B) in the USA, Shaare-Zedek Medical Centre (C) in Israel, and

Cerrahpasa Medical Faculty (D) in Turkey). In a single session, residents were asked 10 questions each about GR and Behcet's disease and 3 questions each about pathergy test and PPD. Rates of correct responses between IM residents from the USA and IM residents from Israel/Turkey were compared using Fischer's Exact test.

Results 69 residents participated in this study (A: 26, B: 13, C: 10, D: 20). The Table 1 shows the proportion of correct answers given for each category of questions. There was a statistically significant difference between the rate of correct responses among IM residents in the USA compared to those in Israel/Turkey with respect to the categories of GR, Behcet's disease, and pathergy test. This difference was more pronounced in the questions regarding Behcet's disease and pathergy test.

Abstract THU0241 Table 1

	General rheumatology	PPD	Behcet's disease	Pathergy test
United States	33%	61%	35%	21%
Israel/Turkey	48%	70%	65%	66%
p value	<0.0002	0.05	<0.0001	<0.0001

Conclusion Knowledge of Behcet's disease, and especially pathergy test among IM residents tested is significantly lower in the USA than among those in Israel/Turkey. The difference in the prevalence of Behcet's disease in these two regions of the world may partly explain this disparity. Contrarily, lack of GR and Behcet's disease knowledge may contribute to decreased recognition and thus underdiagnosis of Behcet's disease resulting in the reported low prevalence in the US. Only by improving the education of internists with respect to Behcet's disease will the true prevalence of this condition be realised.

THU0242 CAN SOCIAL SUPPORT AND SOCIAL NETWORK PREDICT THE EVOLUTION OF QUALITY OF LIFE OVER TIME IN EARLY RHEUMATOID ARTHRITIS? THE EURIDISS COHORT STUDY

¹V Demange, ¹F Guillemin, ¹M Baumann, ²TP Suurmeijer, ³T Moum, ⁴D Doeglas, ¹S Briancon, ²WJ Van den Heuvel. ¹School of Public Health, Faculty of Medicine, University of Nancy, Vandoeuvre-Les-Nancy, France; ²Interuniversity Centre for Social Science Theory and Methodology; ³Department of Behavioural Sciences in Medicine, University of Oslo, Oslo, Norway; ⁴Department of Clinical Psychology, University of Groningen, Groningen, The Netherlands

10.1136/annrheumdis-2001.786

Background There is clear epidemiological evidence for a relationship between social support and health. Though, the mechanism remains unclear whether a poor social support precedes or follows a poor health status. Rheumatoid arthritis (RA) is a chronic inflammatory disease affecting particularly quality of life.

Objectives First, to investigate cross-sectional relationships between quality of life and social support (SS) and support network (SN) in patients with early RA. Then to look at how baseline SS or SN influence quality of life over time; and last, to explore how variations over time in SS or SN affect quality of life over time.

Methods Subjects were members of the EURIDISS population-based cohort from France, the Netherlands and Norway, and had suffered from RA for less than 5 years. Yearly assessments

were made of SS, SN, psychological well-being (GHQ), and functional disability (HAQ). Variance and covariance analyses with repeated measures were calculated and adjusted for potential confounding variables.

Results A total of 542 subjects had yearly assessment for three years. On average, those receiving a greater amount of specific SS (daily emotional support or social companionship) ($p = 0.05$ to $p = 0.0001$) or surrounded by a stronger specific SN (those aspects providing emotional support) ($p < 0.001$) experienced a better quality of life. No statistically significant relationship was observed between baseline level of SS or SN and changes in quality of life over time. Similarly, changes in SS or SN did not appear to be associated with improvements in quality of life over time.

Conclusion Better specific SS and specific SN are cross-sectionally associated with better quality of life in early RA patients. Baseline or changes in SS or SN do not seem to predict changes in quality of life over time for these patients. It may be useful to measure SS in early RA patients in order to identify individuals who receive little specific SS. Further work is needed to devise simple ways to measure SS, to develop interventions that provide additional support, and to investigate whether provision of structured SS by relevant professionals would be more useful than increasing non-specific intervention.

THU0243 FINDINGS OF AN NSAID-INDUCED ULCER STUDY IN JAPAN – AMONG THE JAPAN RHEUMATISM FRIENDSHIP ASSOCIATION MEMBERS -

T Saito. *Institute of Rheumatology, Tokyo Women's Medical University, Shinjuku-Ku, Japan*

10.1136/annrheumdis-2001.787

Background Non-steroidal Anti-inflammatory Drugs (NSAIDs) play a vital role in reducing morning stiffness and/or joint pain of patients receiving Rheumatoid Arthritis (RA) drug therapy. NSAID-induced gastric ulcers, however, have been identified as a major side effect, highlighting the need for countermeasures. Asymptomatic gastric ulcers, in particular, have been receiving much attention as a blind spot in RA drug therapy.

In this context, a questionnaire survey was conducted involving 3,000 cases of the listed 22,101 cases of the Japan Rheumatism Friendship Association (incorporated non-profit organisation), and subsequent analysis focused on gastrointestinal disorders.

Objectives The objective was to aid future countermeasures by studying and analysing facts regarding drug-induced side effects (gastrointestinal disorders in particular), clinical practices and issues taking place between physicians and RA patients.

Methods Mail survey of 3,000 patients randomly sampled from the listed 22,101 members of the Japan Rheumatism Friendship Association (non-profit organisation).

Results A total of 1,764 responses were returned (response ratio 58.8%).

As known, many RA patients visit multiple medical institutions for treatment of their complications including hypertension, cardiopulmonary disorders, infections, etc., thus receiving prescriptions for NSAIDs from non-RA specialists as well. Gastric ulcer incidence was significantly higher among these patients who tend to double-dose NSAIDs.

A total of 1,146 RA patients were discovered to have experienced gastrointestinal disorders caused by NSAIDs, which translates to 65% of the overall respondents. 325 (18%) experienced

ulcers (including gastric and duodenal ulcer). Among the 280 patients who experienced gastric ulcers, 97 (35%) had been hospitalised—65 (67%) of whom for 3 weeks or longer.

The number of gastric ulcer patients was proportionate to the period of RA morbidity. On the other hand, despite RA patients' concerns about NSAID-induced side effects, only 652 (37%) received explanations from their physicians. Only 717 (41%) of the overall respondents were recommended to receive endoscopic examinations. Many aspects of this study have shed new light on the facts regarding RA treatment in Japan, revealing many issues that require closer attention.

Conclusion RA is a general and progressive disease causing lesions not only in joints but also in multiple organs, which means many different drugs other than those for RA tend to be added to patients' regimen. Results of this study revealed many issues to be resolved in RA treatment in Japan: double-dosing of NSAIDs, the fact that only 652 (37%) received explanation from physicians about possible side effects and that only 717 (41%) were recommended to receive endoscopic check-ups despite the high incidence of gastrointestinal disorders (1,146 patients or 65%) and gastric and duodenal ulcers (325 patients or 18%). Results of this study indicate urgent need to improve RA treatment in Japan through educating not only physicians and patients but also the general public.

THU0244 CROSS-SECTIONAL STUDY OF OSTEOARTHRITIS IN SOME NATIVE POPULATIONS OF RUSSIA

LI Alexeeva, S Erdesz, NM Mylov, AV Smirnov, LI Benevolenskaya. *Epidemiology and Genetics, Institute of Rheumatology, Moscow, Russia*

10.1136/annrheumdis-2001.788

Background

Objectives To study the prevalence of knee and hand osteoarthritis (OA) in Ugro-Finnish populations of Russia.

Methods We examined by cross-sectional epidemiological methods three populations from Russian Ugro-Finnish language family: Erzya (E), Moksha (Mo), Marij (Ma). Using standard questionnaire we examined 1312 rural residents aged 15 yrs and over (representativity >80%). Knee and hand OA was diagnosed by using ACR criteria. Films were readed using Kellgren-Lawrence scales.

Results 339 from 1312 had knee joint pain. The prevalence of knee OA was 11,2% in E, 12,5% in Mo, 8,1% in Ma in individuals aged over 15 and 14,1%, 23,8%, 14,4% accordingly in individuals aged over 50.

212 from 1312 had hand joint pain. Hand OA was diagnosed in 10,0% E, in 3,7% Mo and in 3,8% Ma. The prevalence of hand OA increased with age: 19,8%, 12,2%, 13,5% accordingly in individuals aged over 60.

Knee and hand OA were more higher among women than men.

Conclusion These data demonstrate that knee and hand OA is widespread in Ugro-Finnish populations especially in individuals aged over 50. The ACR criteria for hand and knee OA may be used in population studies.