days of withdrawing TNF signals. Dose-response studies indicated that TNF-treated T cells required TCR engagement with higher peptide concentrations for longer periods of time for commitment to IL-2 production. Subsequent experiments revealed that chronic TNF exposure led to a reversible loss of TCR ζ chain expression, in part through a reduction in gene transcription. Downregulation of TCR ζ impaired TCR/CD3 assembly and expression at the cell surface, and uncoupled membrane proximal tyrosine phosphorylation events, including phosphorylation of the TCR ζ chain itself, CD3 ϵ , ZAP-70 protein tyrosine kinase and LAT. Intracellular Ca²⁺ mobilisation was also suppressed in TNF-treated T cells. Signals transduced through a single chain Fv antibody/FcR γ chain chimeric receptor that recognises native type II collagen, and which utilises TCR ζ for IL-2 gene transcription, were also attenuated. We propose that TNF may contribute to T cell hyporesponsiveness in chronic inflammatory and infectious diseases, such as rheumatoid arthritis, by mechanisms which include downregulation of TCR ζ expression. We speculate that by uncoupling proximal TCR signals TNF could also interrupt mechanisms of peripheral tolerance which are dependent upon intact TCR signal transduction pathways.

Objectives

Methods

Results

Conclusion

OP0032 INVASION OF SYNOVIOCYTES IS INHIBITED BY GENE TRANSFER OF TNF-BP OR IL10 IN AN *IN VITRO* INVASION MODEL

¹EJ Pieterman, ¹PH Goossens, ²WH vd Laan, ¹AL Huidekoper, ³MJ Rabelink, ³RC Hoeben, ¹FC Breedveld, ¹TW Huizinga. ¹Rheumatology; ²Gaubius Laboratory, TNO Prevention and Health, Leiden, The Netherlands; ³Department of Molecular Cell Biology, Leiden University Medical Centre

10.1136/annrheumdis-2001.817

Background In RA fibroblast like synoviocytes (FLS) degrade and invade into adjacent cartilage. An *in vivo* model is the SCID-mouse/Human cartilage/synoviocyte model (S. Gay). Previous studies indicate that the invasive behaviour of fibroblast like synoviocytes can be tested *in vitro* in a matrigel transwell system. Matrigel, mainly composed of laminin and collagen IV, serves as a model for cartilage.

Objectives The aim of this study was to compare the invasive behaviour of FLS from RA and osteoarthritis (OA) patients and to investigate the effect of adenoviral (Ad) transfer of genes encoding IL-10 and TNF-binding protein (p55) (TNF-BP) on the invasive behaviour of FLS from RA patients.

Methods FLS from 43 RA and 28 OA patients obtained from synovial tissue harvested at joint replacement surgery, were seeded at confluency in serum free medium on top of a matrigel coated transwell filter with 8 μ m pores. Medium with 10% Fetal Calf Serum and 10% Human Serum was used in the lower compartment. Three days post incubation cells were fixated, stained and the invaded synoviocytes on the lower side of the filter were counted. To test the effect of IL10 and TNF-BP, RA synoviocytes were infected overnight with 5, 10, 50 and 100 plaque forming units (pfu)/cel of Ad. IL10, Ad. TNF-BP or Ad. luciferase (negative control) and then tested in the invasion model.

Results Significantly more RA synoviocytes invaded through the matrigel (median = 4035 cells) as compared to OA synoviocytes (median = 1900 cells; $p < 0.001$). IL10 and TNF-BP gene

transfer both resulted in a dose dependent inhibition of invasion with a maximal inhibition of 93.7% (± 10.7) and 86.6% (± 14.5) respectively, while luciferase gene transfer showed a maximal inhibition of 17.6% (± 1.7).

Conclusion In conclusion, the invasive behaviour of FLS can be studied in the matrigel transwell system. This assay discriminates between the invasive behaviour of RA and OA FLS. The invasive behaviour of RA synoviocytes can be strongly inhibited by IL10 indicating that IL10 is able to downregulate the proteins involved in invasive growth. The inhibitive effect of TNF-BP indicates that continuous production of TNF is involved in invasive behaviour of synoviocytes.

OP0033 SYSTEMIC ADENOVIRUS MEDIATED GENE THERAPY USING A CHIMERIC SOLUBLE IL-1 RECEPTOR TYPE II ? IGG PROTEIN REDUCES ACUTE INFLAMMATION IN ANTIGEN-INDUCED ARTHRITIS

J Dudler, R Salvi, N Busso, V Chobaz-Pédat, A So. Service de Rhumatologie, CHUV, Lausanne, Switzerland

10.1136/annrheumdis-2001.818

Background The anti-inflammatory effects of IL-1 inhibition are well established but most studies have focused on the endogenous IL-1 receptor antagonist (IL-1Ra) protein. However, IL1ra has a very short plasma half-life and hundred-fold excess over IL-1 is required to obtain a 50% inhibition of IL-1 responses. Another naturally occurring inhibitor, soluble IL-1 receptor type II (sIL-1R2) may offer a higher IL-1 neutralising activity and several studies have emphasised its potential function as a regulatory "decoy" for IL1b.

Objectives To construct soluble IL-1 receptor type II chimeric protein with an IgG heavy chain and to evaluate its therapeutic potential in a murine arthritis model using a gene therapy approach.

Methods The murine IL-1R2 extracellular domain was cloned using a RT-PCR protocol from total cytoplasmic RNA, sequenced and a chimeric expressing gene constructed by the addition of a XbaI fragment encoding the constant and hinge regions of a murine IgG1 heavy chain. The IL-1R2IgG coding insert was subcloned into an E1-deleted adenoviral shuttle expression plasmid under the control of a CMV promoter and the resulting plasmid used to construct a standard first generation E1-deleted recombinant adenovirus (Ad-sIL1R2 Ig) using standard homologous recombination techniques in 293 cells. Stock viral preparations were produced in 293 cells and purified by ultracentrifugation in two CsCl² gradients. Viral titer (pfu/ml) was determined by plaque-assay. An EGFP (enhanced green fluorescent protein) expressing recombinant adenovirus (Ad-EGFP) was prepared using a similar protocol. Expression was confirmed by Western blotting using transient expression on HeLa cells under serum free-conditions.

The therapeutic effects of this approach was tested in the acute murine antigen-induced arthritis model (AIA).

Results Western blotting under reducing and nonreducing conditions confirmed expression and dimerization of the chimeric protein. More importantly, adenovirus-mediated systemic expression of sIL1R2 Ig suppressed acute inflammation *in vivo* in murine AIA. Mice were injected into the tail vein with 109 pfu Ad-sIL1R2Ig or Ad-EGFP three days before induction of arthritis by intraarticular injection of 100 μ g mBSA into the right knee. The left knee was injected with PBS as a control. Joint inflammation

determined by ⁹⁹technetium pertechnetate uptake⁶⁹ was statistically significantly decreased at day 3 and 7 post-injection in animal treated with Ad-sIL1R2 Ig as compared to Ad-EGFP (10 animals per group, 2 experiments) but proteoglycan depletion of cartilage did not appear to be diminished.

Conclusion Our study demonstrated the value of a soluble IL1R2Ig chimeric protein has a IL1 inhibitor *in vivo*. It also demonstrated adenovirus-mediated gene expression allow for sufficient level of expression. But our preliminary data also raise new questions of the respective values and roles of IL1 or TNF α inhibition regarding acute inflammation and cartilage degradation.

OP0034 INFLAMMATION-INDUCIBLE INTRA-ARTICULAR PRODUCTION OF HUMAN IL-1 RECEPTOR ANTAGONIST RESULTS IN MORE EFFICIENT INHIBITION OF COLLAGEN-INDUCED ARTHRITIS THAN DOES CONSTITUTIVE LOCAL EXPRESSION OF THE SAME TRANSGENE

¹FA Van de Loo, ¹AC Bakker, ¹LA Joosten, ¹OJ Arntz, ²AW Varley, ²RS Munford, ¹WB Van den Berg. ¹Rheumatology Research Laboratory, ²Molecular Host Defence Laboratory, UMC St-Radboud, Nijmegen, The Netherlands

10.1136/annrheumdis-2001.819

Background Achieving biologically effective yet safe levels of recombinant anti-inflammatory proteins by gene therapy may require three regulatory features: 1) a basal level that is sufficiently low to avoid or minimise chronic immunosuppression 2) transcriptional regulation over a wide dynamic range, and 3) induced expression levels that are sufficiently high to achieve the desired biological effects. The C3-Tat/HIV promoter construct seems to have these properties.¹ In this two-component expression system the complement factor 3 (C3) promoter regulates production of the HIV transactivator of transcription, and the Tat protein then stimulates expression of the desired transgene, which is regulated by the HIV-LTR promoter. Both C3 and HIV LTR promoters are inducible by proinflammatory cytokines.

Objectives To achieve disease-inducible expression of recombinant anti-inflammatory proteins in order to allow autoregulation of drug dose by natural homeostatic mechanisms.

Methods We compared a disease-inducible, two component expression system (C3-Tat/HIV) with the constitutive immediate early cytomegalovirus (ieCMV) promoter in the polyarticular Collagen-Induced Arthritis (CIA) model in mice. DBA/I mice were immunised with bovine type II collagen and boosted on day 22. On day 22, mice without any clinical signs of arthritis were selected and adenoviral vectors (Ad. CMV-Luc, Ad. CMV-IL-1Ra, or Ad. C3-Tat/HIV-IL-1Ra) that contained luciferase or the human IL-1Ra gene under control of one of the two promoters were used to transfect the synovial lining of both knees. The injected knee joints and ipsilateral paws were then scored for signs of arthritis and at the end histology was taken.

Results Inducible promoter-driven IL-1Ra expression resulted in significantly improved inhibition of CIA than did CMV-driven IL-1Ra production. Moreover, overexpression of IL-1Ra in the knee joints also prevented CIA in the ipsilateral paws.

Conclusion Our data (1) demonstrate the feasibility of an inducible expression system for producing a recombinant transgene for treatment of arthritis and (2) show that this system is more effective than strong, constitutive transgene expression for preventing collagen-induced arthritis in mice.

REFERENCE

- 1 Varley AW, Geisler SM, Gaynor RB, Munford RS. A two-component expression system that responds to inflammatory stimuli *in vivo*. *Nat Biotechnol.* 1997;**15**(10):1002-6

OP0035 THE PRODUCTION OF CXCR3 AGONISTIC CHEMOKINES BY SYNOVIAL FIBROBLASTS FROM PATIENTS WITH RHEUMATOID ARTHRITIS

¹A Ueno, ¹M Yamamura, ¹M Iwahashi, ¹T Aita, ¹A Okamoto, ²K Nishida, ²H Inoue, ¹H Makino. ¹Department of Medicine III, ²Department of Orthopedic Surgery, Okayama University Medical School, Okayama, Japan

10.1136/annrheumdis-2001.820

Background IFN-g-producing Th1 or Th0 cells predominate in the synovium of rheumatoid arthritis (RA). It is now known that the accumulation of Th1 cells is mediated by Th1 cell expression of the chemokine receptors CCR5 and CXCR3 and the local expression of their agonistic chemokines. Accordingly, T cell infiltrates in RA synovium have been shown to mostly express CCR5 and CXCR3.

Objectives Since synovial fibroblasts are known as an active participant in joint destruction and chronic inflammation, we investigated the expression of CXCR3 agonists such as IP-10, Mig, and I-TAC in synovial fibroblasts, as well as synovial tissues from RA patients. In addition, the importance of these fibroblast-derived chemokines in the chemotactic response of RA blood CD4⁺ T cells was determined.

Methods Synovial fibroblast cell lines, prepared from RA synovial tissues, were incubated with or without IFN-g, IL-1, TNF- α , IL-17, or various combinations. Levels of protein and mRNA for IP-10, Mig, and I-TAC were measured by ELISA and RT-PCR, respectively. The distribution of chemokine-producing cells in RA synovial tissue was determined by immunohistochemistry. Fibroblasts were cultured overnight on chamber slides and their chemokine synthesis was detected by intracellular staining and a confocal laser scanning microscopy. The chemotactic response of RA blood CD4⁺ T cells to fibroblast culture supernatants with or without antibodies against IP-10, Mig, and I-TAC was determined by an *in vitro* chemotaxis microchamber technique.

Results Concentrations of CXCR3 agonistic chemokines in RA synovial fluid were much higher than those in RA peripheral blood or osteoarthritis (OA) synovial fluid. Immunohistochemical analysis of RA synovial tissue showed that these chemokines were expressed mostly on large cells and vascular endothelial cells in the sublining layer, as well as in the lining layer. The staining was prominent around the lymphocyte aggregate. Synovial tissues from RA more strongly expressed mRNA for all three CXCR3 agonists than OA tissues, and spontaneously released large amounts of their proteins. Synovial fibroblast cell lines isolated from RA, when stimulated with IFN-g, were able to produce these chemokines at the mRNA and protein levels, and this chemokine production was significantly augmented by IL-1, TNF- α , and IL-17. Culture supernatants of activated fibroblasts contained the chemotactic activity for blood CD4⁺ T cells from RA, and this activity was partially but significantly inhibited by antibodies against these CXCR3 agonists.

Conclusion These results indicate that synovial fibroblasts may contribute to the continued Th1 response in RA in part by secreting CXCR3 agonists. In addition, inflammatory cytokines derived from Th1 cells and macrophages, IFN-g, IL-17, IL-1, and TNF- α , are important inducers of these chemokines in the joint.