Background The mechanism by which methotrexate (MTX), a well known antifolate inhibits inflammation is still unknown. The inflammation in rheumatoid arthritis is mediated by pro-inflammatory cytokines such as TNF α and IFN γ .

Objectives To assess whether cytokine production in blood of RA patients is inhibited after taking MTX.

Methods Blood was obtained from 25 RA patients at T = 0 h and T = 2 h after taking MTX (10 mg or) for the first time. Heparinized Whole Blood Cultures (WBC; dilution 1/4) using anti-CD3 and anti-CD28 as stimulus were performed, with and without addition of folic acid (40 g/ml) to the culture. Because MTX has been reported to induce IL-10 production all experiments were also carried out with and without excess neutralising mAb to IL-10. After incubation for 72 h supernatants were measured by ELISA.

Results IFN γ production by WBC of 25 RA patients before (T = 0 h) and after (T = 2 h) MTX.

Abstract THU0044 Table 1

	IFN γ *ng/ ml	IFN γ , FA added	IFN γ , anti IL 10 added	IFN γ , FA and anti IL 10
T = 0 h	53(12)	39(10) ²	94(17)	78(16)
T = 2 h	19(5) ¹	48(12) ³	27(6) ⁴	93(17)

*Mean (SEM). ¹2h after intake of MTX, IFN γ production is inhibited (p = 0.0001); ²Folic acid by itself inhibits IFN γ production (p < 0.0001); ³Folic acid completely reverses the inhibition by MTX; ⁴In the presence of anti-IL-10 similar results were obtained.

Conclusion MTX inhibits *ex vivo* IFN γ production in RA patients by a folate dependent mechanism. MTX does not act via induction of IL-10. Similar results were found for TNF α and GM-CSF. IL-8 production is not affected by MTX.

THU0045 LOCAL PRODUCTION AND MODULATION OF COMPLEMENT PROTEINS IN RHEUMATOID ARTHRITIS SYNOVIUM

¹U Müller-Ladner, ¹I Tarner, ²S Barnum, ²J Echols, ¹E Neumann, ¹M Fleck, ¹M Jüdex, ¹F Kullmann, ²J Mountz, ¹J Schölmerich, ³S Gay. ¹Department Internal Medicine I, University Hospital, Regensburg, Germany; ²Departments Microbiology and Rheumatology, UA, Birmingham, AL, USA; ³Ctr. Exp. Rheum., USZ, Zürich, Switzerland

10.1136/annrheumdis-2001.842

Background

Objectives Rheumatoid arthritis (RA) is a severe chronic disease characterised by inflammation of synovial tissue, which causes pain and dysfunction of joints and leads to their destruction. In RA, complement has been implicated in inflammation and joint destruction based on studies showing reduced levels of native complement components and increased levels of complement metabolites in plasma, synovial fluid and synovial tissue of RA patients. However, there is limited information on local production and activation of key factors of the complement cascade in RA synovium and their potential modulation by novel anti-cytokine therapies.

Methods Using *in situ* hybridization, immunohistochemistry and Western blot techniques, the presence of complement proteins C3, factor B and Sc5b-9 and of the complement receptors C3aR and C5aR were evaluated in RA and osteoarthritic (OA) synovium. In addition, C3 and factor B levels in synovial fluid were

determined by ELISA. Functional assessment was performed by examination of the effects of adenovirus-based sTNF-R p55 gene transfer into RA synovial fibroblasts in the SCID mouse model.

Results Complement proteins C3 and factor B were synthesised mainly around terminal vasculature in all (6/6) RA synovial specimens whereas complement receptors C3aR and C5aR were expressed in adjacent inflammatory infiltrates. In contrast, only limited local production of these factors could be observed in OA. Different activation of the complement cascade was illustrated by more intensive degradation of factor B in RA as compared to OA synovial fluid (mean: 28 vs. 48 μ g/ml). Overexpression of sTNFRp55 revealed upregulation of C5 but not C3 receptor expression in the transduced RA synovial fibroblasts indicating differential effects of anti-cytokine therapies on complement factors in RA synovium.

Conclusion We conclude that local complement production and activation may play an important role in RA synovium. Moreover, modulation or inhibition of local complement production, e.g. by biologic agents and anti-C5 antibodies, which are currently investigated in clinical trials, could be an attractive therapeutic approach for RA.

THU0046 SERUM SOLUBLE SELECTINS LEVELS IN PATIENTS WITH RHEUMATOID ARTHRITIS AND SYSTEMIC SCLEROSIS

A Ates, G Kinikli, M Turgay, M Duman. *Clinical Immunology and Rheumatology, Ankara University Medical School, Ankara, Turkey*

10.1136/annrheumdis-2001.843

Background Soluble forms of adhesion molecules may be found in the circulation, and increased levels of selectins have been reported in some connective tissue diseases. It was suggested that the measurement of levels of circulating soluble selectins is useful as an indicator of inflammatory disease.

Objectives To determine serum levels of soluble (s) E-, L-, and P-selectin in patients with rheumatoid arthritis (RA) and systemic sclerosis (SSc), and to determine whether levels of these molecules correlate with measures of disease activity and pulmonary involvement.

Methods Serum levels of sE-, L-, and P-selectin were determined by sandwich ELISA in 34 patients with RA, 30 patients with SSc and 15 healthy subjects. The patients with RA were classified according to disease activity, functional status, Larsen score reflecting anatomical joint damage, and the presence of pulmonary involvement. The presence of pulmonary fibrosis and DLCO were determined in patients with SSc.

Results The mean soluble selectin levels are shown in Table 1. sE- and sL-selectin levels were significantly higher in patients with active RA compared to inactive patients (respectively p < 0.05 and p < 0.05). Although sP-selectin level was higher in active RA patients than those in inactive patients, the difference did not reach statistically significant (p = 0.06). Significant correlations were found between sL-selectin and CRP (p < 0.05), and white blood cell count (p < 0.01) and between sP-selectin and CRP (p < 0.01), and erythrocyte sedimentation rate (p < 0.01). There were no correlations between selectin levels and Larsen score. sE-selectin level was increased in RA patients with advanced functional impairment (56.0 $\pm$ 39.8 in stage III-IV vs. 32.7 $\pm$ 13.6 ng/ml in stage I-II, p < 0.05). The presence of pulmonary involvement in patients with RA was not correlated with soluble selectin levels. sE-selectin levels was significantly higher

in SSc patients with pulmonary fibrosis compared to other SSc patients ($p < 0.05$). There was a significant correlation between sE-selectin level and DLCO ($r = -0.536$, $p < 0.01$).

Abstract THU0046 Table 1

	sE-selectin (ng/ml)	sL-selectin (ng/ml)	sP-selectin (ng/ml)
RA	43.7 ± 30.9 ^a	378 ± 168 ^{b, c}	383 ± 198 ^c
SSc	44.9 ± 22.3 ^d	552 ± 224	246 ± 163
Healthy subjects	24.9 ± 12.9	672 ± 140	292 ± 199

^a $p < 0.05$ vs. healthy subjects; ^b $p < 0.001$ vs. healthy subjects; ^c $p < 0.01$ vs. SSc; ^d $p < 0.01$ vs. healthy subjects.

Conclusion These results suggest that the measurement of levels of serum soluble selectins may provide a useful additional marker for disease activity in RA patients and for pulmonary involvement, especially sE-selectin, in SSc patients.

THU0047 MATRIXINS, CYTOKINES AND ADHESIVE IMMUNOGLOBULINS WITHIN OSTEOARTHRITIC JOINT

J ??tovickova, H Hulejova, V Pesakova, M Adam.

10.1136/annrheumdis-2001.844

Research Centrum, Institute of Rheumatology, Prague, Czech Republic

Background The most abundant joint disease is osteoarthritis. Recently it was classified as cartilagenous, synovial or osseous but the unique forms may be combined. The main feature of the disease is imbalance between synthetic and degradative processes within joint tissues. Executive part in degradative processes play matrix metalloproteinases, so called matrixins. *In situ*, they are induced by many stimuli, namely by cytokines and by interaction of different adhesive molecules with their ligands. Having been secreted into the extracellular space, they are activated by serine proteinases, or by other matrixins and inhibited by tissue inhibitors of matrix metalloproteinases, TIMPs.

Objectives In the sera of OA patients we found matrixins MMP-9, MMP-2 and MMP-3 and TIMP-1 together with cytokines IL-8 and TNF α and adhesive molecules ICAM-1 and VCAM-1. To distinguish between the forms of osteoarthritis, we search for these molecules in all joint tissues that is in cartilage, synovium, granulous tissue within subchondral bone, synovial fluid and blood serum. The joint tissues and fluids we obtained from surgery during total hip or knee replacement.

Methods The joint tissues were derived from 22 patients. The patients were 17 female and 6 male, 67,5 ± 9,6 years old. The metalloproteinase spectrum in the body fluids and extracts of joint tissues were evaluated by zymography. The concentration of metalloproteinases and TIMP-1 were assessed by ELISA (Bio-trak, Amersham Pharmacia Biotech), the concentration of adhesive immunoglobulins by Bender ELISA (MedSystems Diagnostics GmbH), and the concentration of IL-8 and TNF α were determined on Immulite (DPC, USA).

Results We found that the most abundant matrixine within cartilage and synovium is either gelatinase A (MMP-2) or stromelysin-1 (MMP-3). On the other hand, within granulous tissue, blood sera and synovial fluid, the prevailing matrixine is gelatinase B or stromelysin-1. In cartilage, synovial fluid and blood serum TIMP-1 is abundant over sum of matrixins, whereas in synovium and granulous tissue the sum of matrixins is abundant over TIMP-1. There is about 2 to 10 times greater concentration of IL-8 in the cartilage than in other joint compartments.

We found significant correlation between levels of sICAM-1 and sVCAM-1; between TNF α and MMP-3, TIMP-1, sICAM-1, sVCAM-1, and IL-8; and between IL-8 and sICAM-1 in all solid joint tissues. In the synovial fluid and blood serum the correlation was less pronounced.

Conclusion There is a great degradative potential within all compartments of osteoarthritic joint due to presence of matrixins, namely that of stromelysin-1. Within cartilage and synovial fluid the proteolytic potential is equilibrated by TIMP-1 whereas within synovium and granulous tissue is not. Synovium and granulous tissue are highly probably the propellers of degradative process in osteoarthritis.

REFERENCE

- 1 Yoshihara Y, *et al.* Matrix metalloproteinases and tissue inhibitors of metalloproteinases in synovial fluids from patients with rheumatoid arthritis or osteoarthritis. *Ann Rheum Dis.* 2000;59:455–61

THU0048 ADDITION OF THE RGD MOTIF IN THE FIBRE KNOB OF AN ADENOVIRAL VECTOR IMPROVED THE TRANSFECTION EFFICIENCY AND WHEN USED FOR OVEREXPRESSION OF IL-1RA RESULTED IN A MORE EFFECTIVE INHIBITION OF ARTHRITIS

¹FA Van de Loo, ¹AC Bakker, ¹LA Joosten, ²V Krasnykh, ²DT Curiel, ¹WB Van den Berg. ¹Rheumatology Research Laboratory, ²Gene Therapy Center, UMC St-Radboud, Nijmegen, The Netherlands

10.1136/annrheumdis-2001.845

Background Adenoviral (Ad) vectors expressing anti-inflammatory cytokines are used for therapy in experimental arthritis. Cell recognition and entry is dependent on the initial recognition of the coxsackievirus and adenovirus receptor (CAR) on cells. Recently introducing an Arg-Gly-Asp (RGD) motif in the HI loop of the fibre knob modified the tropism of the virus.¹ The RGD motif is capable of mediating a CAR-independent cell entry via interactions with the α v β 3 and α v β 5 integrins.

Objectives In this study we explored the transfection efficiency of the RGD modified viruses and investigated the effectiveness of intra-articular IL-1Ra overexpression in the collagen-induced arthritis (CIA) model mediated by the RGD modified vector compared to wildtype viruses.

Methods 10e7 pfu of the AdLuc, AdLucRGD, AdmIL-1Ra, or AdmIL-1RaRGD were intra-articularly injected in all experiments. The CIA model was used to compare the wild type and the RGD viruses on their ability to downregulate arthritis. DBA/1 mice were immunised with collagen type II, and boosted 22 days later. Four days after the booster the adenoviral vectors were injected into both knee joints of the mice. Clinical scores of the knee joints were on a scale of 0–3.

Results Twenty-four hours after intra-articular injection of 10e7 pfu virus into normal knee joints, luciferase production in AdLucRGD injected joints was around 22 times higher than in AdLuc injected joints. On day 26 of the CIA model, 10e7 pfu of AdLuc, AdLucRGD, AdmIL-1Ra, or AdmIL-1RaRGD were injected into the knee joints. Onset of the CIA was significantly inhibited in the AdmIL-1RaRGD group (clinical score in the knee joints at day 38 was 0.82 ± 0.24) compared to the other groups (clinical scores were 1.57 ± 0.24 for the AdmIL-1Ra group and 1.86 ± 0.21 for the AdLucRGD group).

Conclusion Transduction of the synovial lining cells in normal knee joints was up to 30 times more efficient when the RGD mediated pathway was used. CIA was significantly better