Objectives To evaluate safety, clinical improvement, and radiographic outcomes in patients receiving ENBREL or MTX for up to 2 years.

Methods Patients enrolled in this 2-year study continued to receive the therapy to which they had been randomised in an open-label fashion after the study was unblinded (at a mean of 17.3 months). The 2-year endpoints were ACR responses (using last on-drug observation carried forward analyses) and change in Sharp score. Readers were blinded to treatment and chronological order of the x-rays. All 632 patients were included in the analyses.

Results 154 of 207 patients (74%) who received ENBREL 25 mg completed 2 years of treatment, compared to 129 of 217 patients (59%) who received MTX. Twice as many patients (12%) discontinued MTX for adverse events as did those taking ENBREL. The ACR 20 response at 2 years was 72% in those randomised to ENBREL and 59% in those randomised to MTX ($P = 0.005$). Radiographic evaluations demonstrated that ENBREL 25 mg was superior to MTX in arresting radiographic progression over the 2-year period. Mean (median) changes in total Sharp score were 1.3 (0) and 3.2 (0.5) units in the ENBREL 25 mg and MTX groups, respectively ($P = 0.001$) and changes in erosion score were 0.7 (0) and 1.9 (0) units, respectively ($P = 0.001$). ENBREL 25 mg prevented radiographic progression in 63% of patients, compared to 51% of MTX patients who did not progress ($P = 0.017$). ENBREL 25 mg was also superior to the 10 mg dose in inhibiting radiographic progression. Significantly more ENBREL 25 mg patients than MTX patients had a clinically meaningful (at least 0.5 units) improvement from baseline in HAQ disability score. ENBREL continued to be well tolerated with over 2 years of exposure.

Conclusion This 2-year study demonstrated that ENBREL was superior to MTX in reducing clinical signs and symptoms of RA and inhibiting radiographic progression.

REFERENCE

- 1 Bathon JM, Martin RW, Fleischmann RM, et al. *NEJM* 2000;**343**:1586–93

SLE – Etiology and pathogenesis/Animal models

AB0069 ANTI BETA 2 GLYCOPROTEIN ANTIBODIES AND THROMBOSIS IN SLE PATIENTS

MA Mansour, M Badr, H El Bebawi, A El Ghobarey. *Rheumatology, Meadi Armed Force Hospital, Cairo, Egypt*

10.1136/annrheumdis-2001.1223

Background

Objectives Beta 2 glycoprotein was identified as a cofactor for the binding of antiphospholipid antibodies. Anticardiolipin antibodies were found to correlate with thrombosis in SLE patients. We measured the subclasses of anti B2 GP1 in LSE patients with or without history of thrombosis.

Methods Fourty four SIE patients with mean ($\pm$ SD) age of 26.7 ($\pm$ 9.9) years, nine of whom had a history of vascular thrombosis along with 30 healthy age and sex matched normal control subjects were studied.

Results There was a correlation between the presence of IgG anti B2 GP I and the incidence of thrombosis ($P < 0.05$). There was, also, a positive correlation between IgG anti B2 GPt and

IgG aCL antibodies and a negative correlation between IgG anti B2 GP1 and thrombocytopenia. However, serum levels of IgG and 1 gM anti B2GPI were increased in patients who had positive LAC activity. This thesis confirm the pathologic role of IgG anti B 2 GP1 (and not 1gM) in the incidence of thrombosis in SLE patients. Also, its relationship to the presence of anti ds DNA antibodies, anti CLA and lupus anticoagulant in SIE is discussed.

Conclusion

AB0070 THE EFFECT OF PROLACTIN ON DISEASE ACTIVITY IN SLE

LC Te Boome, PM Welsing, AA Eijbouts, FH Van den Hoogen. *Rheumatology, University Medical Center, Nijmegen, Netherlands*

10.1136/annrheumdis-2001.1224

Background Prolactin, a hormone with immunomodulatory effects, possible plays a role in the disease activity of Systemic Lupus Erythematosus (SLE). Results of studies are contradictory regarding the correlation between hyperprolactinemia and disease activity.

Objectives To establish whether determination of prolactin is helpful in assessing disease activity in SLE by reviewing relevant literature.

Methods Studies were identified through Medline search. Also references from these articles were reviewed.

Results Ten articles that sought to establish a relationship between prolactin and SLE activity were identified.

Number of patients in the studies varied from 29 to 192, aged 16–73 years, and the percentage of elevated serum prolactin levels in SLE patients was between 2.2–40%. The SLEDAI was used as disease activity index in 7/10 studies. The presence of conditions known to be associated with hyperprolactinemia were either reported or excluded. Steroids treatment was accepted in all studies, DMARDs were used in 8/10 studies. In 5/10 studies bloodsamples were collected midmorning and in 4/10 patients had to fast. Single measurement of serum prolactin concentration and disease activity was done in 6 studies. The other 4 studies had 2–4 measurements per patient. Pearson's correlation coefficient was the used statistical analysis in 7/10 studies. A relation between prolactin and disease activity was reported in one study. The number of included patients in the studies was small and different statistical methods were used.

Conclusion The correlation between serum prolactin levels and disease activity is still uncertain. A cross-sectional as well as longitudinal study with using the correct statistics, power calculation and proper recording of secondary causes of elevated prolactin and medication need to be done.

FRI0083 ELEVATED IL-16 LEVEL IS A NON-GENETIC CHARACTERISTIC OF PATIENTS WITH SEVERE SYSTEMIC LUPUS ERYTHEMATOSUS

¹LR Lard, ²BO Roep, ¹TW Huizinga. *¹Rheumatology, ²Immunohaematology and Bloodbank, Leiden University Medical Centre, Leiden, Netherlands*

10.1136/annrheumdis-2001.1225

Background IL-16, originilly named lymphocyte chemoattractant factor, is a cytokine which is mainly produced by CD8⁺ T cells. Several reports have described that increased levels of IL-