Methods Using flowcytometric whole blood and platelet rich plasma double staining assay systems we analysed CD40 L expression. PE labelled CD40L (cloneTRAP1, Pharmingen) was employed, FITC-CD42b as detection system for platelets. 40 patients with diagnosis of SLE were included into this study, among them 10 with secondary APS and compared to 15 normal healthy donors (NHD). Results were analysed using Cellquest software (Becton Dickinson, San Jose, USA) differentiating single from aggregated platelets under nonstimulatory and stimulatory conditions (thrombin 30 μ M, ADP 10 μ M).

Results We found that CD 40 L expression was higher on aggregated compared to single platelets, where fluorescence means were close to baseline. We compared the whole blood method with platelet rich plasma and found lower background activity in whole blood test. Using established stimulators of platelets we could increase CD40L expression on platelets, especially on aggregated platelets. Analysing CD40L expression in SLE patients we found no significant difference between SLE and NHD, neither on single or on aggregated platelets nor upon stimulation with ADP or thrombin. High CD40 L expression was observed in APS with highly significant differences compared to NHD and SLE analysing single platelets ((NHD < APS ($p < 0,0001$) SLE < APS ($p < 0,0001$)) as well as aggregated platelets. Upon stimulation with thrombin or ADP results were comparable indicating significantly higher CD40 L expression on platelets.

Conclusion We first describe high CD40 L expression on single and aggregated platelets in APS establishing CD40L as a novel platelet activation marker in APS. Interplay between CD40L on platelets and endothelial cells may contribute to procoagulatory conditions in APS. On the other hand we could not show altered CD40L expression in patients with SLE without clinical APS.

OP0075 CHROMATIN CLEARANCE FROM NECROTIC CELLS BY A CONCERTED ACTION OF DNASE 1 AND SERINE PROTEASES RESEMBLES APOPTOTIC DNA-DEGRADATION

M Napirei, H Karsunky, R Paddenberg, B Zevnik, S Wulf, T Mörröy, HG Mannherz. *Ruhr-Universität Bochum, Abteilung Für Anatomie Und Embryologie, Bochum, Germany*

10.1136/annrheumdis-2001.1239

Background Systemic lupus erythematosus (SLE) is an autoimmune disease characterised by the presence of antinuclear antibodies (ANA) which tend to accumulate as immunodeposits in different tissues and cause inflammations. One main feature of SLE-pathogenesis seems to be a reduced elimination of cellular debris after cell death.^{1,2} The recently generated mice deficient for the endonuclease Dnase 1 has underscored the important influence of defects of chromatin clearance for the onset of SLE.³

Objectives

Methods Cell nuclei of MCF-7 cells were isolated and either incubated with serum from Dnase 1 wildtype and Dnase 1 deficient mice or with recombinant human DNASE 1 alone and in combination with different serine proteases. Subsequently the DNA was isolated and analysed by agarose gel-electrophoresis.

DNA-degradation of MCF-7 cells incubated with different amounts of native or heat inactivated bovine serum in DMEM was analysed under apoptotic (2 μ M staurosporin) and necrotic conditions (1 mM H₂O₂) or after complement lysis due to the

application of 6.7% (v/v) rabbit complement. Subsequently the DNA was isolated and analysed.

Results Recombinant human Dnase 1 could induce an apoptotic-like DNA-fragmentation pattern in isolated MCF-7 cell nuclei by a concerted action with different serine proteases like plasmin, trypsin, a-chymotrypsin, thrombin and kallikrein. Dnase 1 alone only induces an unspecific, time-retarded DNA-degradation. Serum from Dnase 1 wildtype mice also induced DNA-laddering in cell nuclei whereas serum from knockout mice did not due to the lack of Dnase 1. The laddering capacity of wildtype serum could be inhibited by serine protease inhibitors. In contrast to apoptotic cells native bovine serum and not heat-inactivated serum induces internucleosomal DNA-laddering in MCF-7 cells lysed by rabbit complement or by the cytotoxic action of H₂O₂.

Conclusion This study gives insight into a new function of Dnase 1 which could be linked with its protective role towards the induction of an anti-DNA immunoreaction and subsequently towards SLE-pathogenesis.^{2,3} Dnase 1 in connexion with serine proteases is able to effectively degrade the genomic DNA of nuclei and of cells lysed by complement or the cytotoxic action of H₂O₂. In contrast apoptotic cells were not influenced by serum Dnase 1 and serine proteases. Since apoptotic cells undergo a cell-autonomous intracellular DNA-degradation process one can conclude that Dnase 1 in connexion with serine proteases fulfills this process for necrotic cells.

REFERENCES

- 1 Vanholder R, De Keyser F, Kips J, Praet M, Naeyaert JM. The pathophysiology of lupus erythematosus. *Eur J Dermatol.* 1998;1:4-7
- 2 Walport MJ. Lupus, Dnase and defective disposal of cellular debris. *Nat Genet.* 2000;25:135-6
- 3 Napirei M, Karsunky H, Zevnik B, Stephan H, Mannherz HG, Mörröy T. Features of systemic lupus erythematosus in Dnase 1-deficient mice. *Nat Genet.* 2000;25:177-81

OP0076 KIDNEYS OF AUTOIMMUNE NZB/W MICE CONTAIN LONG-LIVED PC

AH Radbruch, G Cassese, S Lindenau, B De Boer, S Arce, A Hauser, G Riemekasten, C Berek, F Hiepe, RA Manz. *Cell Biology, Deutsches Rheuma-Forschungszentrum Berlin (DRFZ), Berlin, Germany*

10.1136/annrheumdis-2001.1240

Background NZB/W mice develop a disease similar to human systemic lupus erythematosus (SLE), including autoantibody production, hypergammaglobulinaemia and inflammation of the kidneys. Large numbers of lymphocytes infiltrate the kidneys of these mice but the role of this organ for the production of antibodies is not clear.

Objectives Here, we compare the role of bone marrow, spleen and inflamed kidneys of NZB/W mice for the activation of B cells and for the persistence of antibody secreting cells (ASC).

Methods NZB/W mice were immunised i.p.i with Ovalbumin and then boosted 1 month later. The number of antigen-specific PC present in spleen, bone marrow and kidneys was determined by Elispot. The frequency of different cell types was determined by flow-cytometry. The distribution of PC in the different organs was investigated by microscopy.

Results ASC are present in the kidneys of mice with full blown disease, as many as in the spleen and bone marrow, and 50 times more than in the kidneys of normal mice. In the kidneys, ASC are located mainly in the outer medulla, close to B- and T cell infiltrates. The specificity of the ASC in the inflamed kidneys is not restricted to self-antigens. After immunisation of NZB/W