Methods In November 1999 all physical therapists working within the catchment area of two rheumatological centres (Leiden, Enschede) were invited to participate. Education programmes were organised in parallel in both regions and consisted of a 5-day postgraduate training programme, followed by 2-monthly seminars. The programme focussed on examination and treatment of patients with rheumatic diseases and on communication with patients with a chronic disease. Communication among health professionals was enhanced by teaching practices within the rheumatological centres and by developing a draft guideline on communication among health professionals. Outcome measurements included questionnaires on the organisation, didactics and contents of the training programme (6-point scales, range 1; very bad to 6; very good), knowledge, interdisciplinary communication and patient satisfaction.

Results 63 of the 64 physical therapists (98.4%) finished the education programme and were formally registered. Evaluation of the training programme measured directly after each part of it showed positive scores with respect to contents, didactics and organisation (all medians 5, ranges 2–6). Knowledge scores increased significantly directly following the training programme ($p < 0.001$; Mann Whitney U). Measurements regarding the effectiveness on quality of care (interdisciplinary communication and patient satisfaction) will be completed in 2001.

Conclusions The preliminary results indicate that setting up a system of networks and continuing education for physical therapists with respect to the treatment of patients with rheumatic diseases is feasible. The complete set of results will be available by the end of 2001. Further research will focus on the implementation of the system on a larger scale. Funded by The Health Research and Development Council of The Netherlands; ZON.

SP0170 IMPLEMENTING EDUCATIONAL TRAINING PROGRAMS FOR HEALTH CARE PERSONNEL IN ARTHRITIS

LX Hansson. *Research and Development, The Swedish Rheumatism Association, Stockholm, Sweden*

10.1136/annrheumdis-2001.1267

Co-operation for education of health care personal in arthritis.

In Sweden we have 24 County Councils that are responsible for health care.

One of the most important tasks the Swedish Rheumatism Association, the SRA, is to give health care professionals more knowledge so they can discover early arthritis. If they do we today have a good arsenal of therapies to give people with arthritis a hopeful future. The communities themselves don't have resources to give their personnel further education in medical specialties. In Sweden as in other countries the medical companies offer the doctors a lot of specialised education. This is criticised by the public and there are forces in movement to change the situation.

For a long time the Swedish Arthritis/Rheumatism Association has given courses to health care personnel but this has only been the case in some parts of Sweden. Therefore our organisation, with support from 6 medical companies, now has started an educational project. This last autumn we negotiated with the medical companies to support us and in January we employed one person to contact all the County Councils to get a co-operation project with them to plan for educational days i.e. training for the medical profession.

We have a board, a steering group for this which includes representatives for the organisation of the GPs, for the organisation SRF (the rheumatologists), for Sverefo which represents the professionals from the arthritis team, for our own hospital Spenshult (owned by the SRA) representing the Patient Partner Project in Sweden which is employed by us and some other additional important partners.

The contents of the education is decided upon in cooperation between representatives from the SRA and the county councils mainly on the basis of some educational programs in arthrosis, rheumatoid arthritis, fibromyalgia och osteoporosis which the board picks out. Our patient owned hospital Spenshult in Halmstad on the west coast of Sweden has the main responsibility for the implementation of the educational days and the medical quality. We also try to stress patient influence and the point of view of the patients.

The companies can have exhibitions in the vicinity and treat the participants to food and an attractive free time program. The curriculum usually doesn't contain themes from rheumatology but instead fields like leadership, conflict management and personal development. The educational days vary from a group of 20 participating doctors working with dialogue pedagogy to lectures with 150 participants. We stress a qualitatively good documentation.

The year 2001 is a formative year for the project. Until now we have carried out three educational programs this spring and plan seven for the fall. In 2002, which together with the Swedish Pharmacies has been declared the year of Arthritis/Rheumatism, we plan to carry through approximately forty such educational programs in all county councils in Sweden.

Another project which I want to mention here also covers the education of health care professionals is to build up "e-learning" for the rheuma team.

At our next conference we hope to be able to present some more experiences from both these projects.

Health services, economics and outcome research

HP0034 RESPONSIVENESS OF SELECTED DISEASE-SPECIFIC PATIENT-ASSESSED INSTRUMENTS IN ANKYLOSING SPONDYLITIS (AS)

¹KL Haywood, ²AM Garratt, ³K Dziedzic, ⁴PT Dawes. ¹Department of Health Sciences and Clinical Evaluation, University of York, York; ²Unit of Health Care Epidemiology, Institute of Health Sciences, Oxford; ³Primary Care Sciences Research Centre, Keele University, Newcastle-Under-Lyme; ⁴Staffordshire Rheumatology Centre, North Staffordshire Acute Hospitals Trust, Stoke-on-Trent, UK

10.1136/annrheumdis-2001.1268

Introduction Five instruments, including an individualised measure of AS-related quality of life, the Patient-Generated Index (PGI-AS), have been assessed for measurement properties of reliability, validity and responsiveness in a large group of AS patients. Responsiveness refers to the ability to detect clinically important change and is an essential requirement for evaluative instruments.

Objective To establish comparative evidence in support of instrument longitudinal validity and responsiveness.

Methodology A questionnaire containing all instruments was mailed to over 450 patients randomly selected from six AS databases. Patients completed the questionnaire at baseline (response

rate 76.9%; males 74.2%; mean age 46.0 yrs (SD 12.6), range 18–75 yrs; mean symptom duration 19.8 yrs) and 6 months. Self-reported AS and general health transition at 6 months were the criteria used to assess validity and responsiveness. The relationship between changes in instrument scores and responses to transition items was assessed for a linear trend.

Results Significant correlations between mean change in instrument scores and both forms of transition question supports the longitudinal validity of all instruments ($p < 0.001$): the largest levels of change were found for the PGI-AS and the Bath AS Disease Activity Index (BASDAI), both demonstrating a stronger relationship with general transition. The modified standardised response mean (MSRM) was used to assess responsiveness (Table 1). The Body Chart, a measure of AS specific pain, had a moderate relationship with AS transition only (better -0.44 , worse 0.51), which may support the importance of specific pain evaluation.

Conclusion Although certain domains may not be expected to change over 6-months, this period reflects normal practice in AS evaluation in many rheumatology centres. The responsiveness of the PGI-AS and BASDAI may make them suitable for routine monitoring where management may result in small but important changes in health related quality of life.

Abstract HP0034 Table 1

	AS transition – MSRM Better (n = 56)	AS transition – MSRM Worse (n = 73)	General transition – MSRM Better (n = 51)	General transition – MSRM Worse (n = 62)
AS Quality of Life (ASQoL)	–0.29	0.35	–0.55	0.26
BASDAI	–0.60	0.22	–0.67	0.43
Revised Leeds	–0.32	0.28	–0.37	0.34
Disability (RLDQ)				
PGI-AS (informed/ open)	0.42 (n = 20)	–0.48 (n = 21)	0.74 (n = 25)	–0.75 (n = 18)

HP0035 ASSOCIATION BETWEEN TESS FUNCTIONAL QUESTIONNAIRE AND TIMED TESTS OF FUNCTION

¹KL Barker, ²SE Lamb, ³AH Simpson. ¹Physiotherapy Research Unit, Nuffield Orthopaedic Centre NHS Trust, Oxford; ²School of Health and Social Sciences, Coventry University, Coventry; ³Department of Orthopaedic Surgery, University of Edinburgh, Edinburgh, UK

10.1136/annrheumdis-2001.1269

Background It is important to evaluate the functional result of any orthopaedic intervention. This is commonly achieved by completing tests which evaluate function in a controlled environment under a specific set of conditions, allowing objective measures to be gathered. Alternatively patient-reported assessments may be used, which reveal the limitations that the patients experience when carrying out everyday activities in their own environment. This study investigated the association between these two methods.

Method 40 adult patients undergoing lower limb reconstruction surgery completed the Toronto Extremity Salvage Score (TESS) pre-operatively and at 6, 12 and 24 months post surgery. TESS evaluates a single domain, physical disability, based on patients' reports of their function. It is applicable to a heterogeneous

population being sensitive to change across a range of levels of disability.¹ They also performed 3 timed tests of stair climbing, sit-to-stand and walking.

Results Within subjects changes were analysed using the Wilcoxon matched-pairs test and associations between variables by a Spearman's Rank Correlation Coefficient.

Subjects, on average, improved on all of the functional measures, with significant improvements from the baseline score at 12 and 24 months ($P < 0.001$). Changes at 6 months were not significant. There was a poor correlation between TESS and the timed function tests. To explore this further the association between specific questions in TESS and the objective performance tests were tested. This showed significant correlations between walking speed and the question about walking indoors (0.53 , $P < 0.003$), between stair climbing and the question about going upstairs (0.64 , $P < 0.001$) and downstairs (0.42 , $P < 0.025$) and sit-to-stand and the question about getting out of a chair (0.41 , $P < 0.03$).

Conclusions The association between the self-reported and observed outcome measures was not as strong as may have been expected when both purport to measure the same dimension of physical function. One explanation for this may be that the observed outcomes were all timed activities and patients' satisfaction with their functional abilities may not be related to time. Whilst timed tests may be reliable tests of function, they may not be as valid or useful a measure as patient-reported measures.

REFERENCE

- 1 Davis AM, Bell RS, Badley EM, *et al.* Evaluating functional outcome in patients with lower extremity sarcoma. *Clin Orthop Relat Res.* 1999;**358**:90–100

SP0169

NEW APPROACH TO THE CARE OF RHEUMATIC PATIENTS WITHIN THE PROJECT "PATIENT PARTNERS"

M Dobesova. Out-Patients Department for Rheumatic Patients, Hospital in Bruntal, Bruntal, Czech Republic

10.1136/annrheumdis-2001.1270

New approaches to the care of rheumatic patients within the Project "Patients Partner". An important component of the care of patients with a chronic rheumatic disease is a complex care and an individual approach. Rheumatism is generally thought of as a disease of elderly persons, although certain types of rheumatism attack also younger age groups. There are various topics that I feel deserve to be given attention to:

- Communication and information – the patient should learn communication skills and should receive a maximum of information about his disease and its treatment.
- Complex care and the importance of teamwork. Of importance is a good collaboration among the various specialists. The role of the nurse in providing complex care is indispensable.
- Establishment of national and international organisations of rheumatic patients – a good opportunity for meeting other patients, exchanging experience, giving advice about how to cope with pain. A good example is the Revma League in Prague and its branch at Bruntal.
- Social problems – equal rights for healthy and disabled persons (Public transport, abolishment of legal discrimination etc.).