Objectives Aim of current study was to estimate predictive value of cytology of induced sputum for progressive pulmonary involvement in patients with connective tissue diseases.

Methods Forty-five patients with established diagnosis of connective tissue disease (SLE, PSS, PM/DM), were examined for pulmonary involvement (X-ray, spirometry, diffusion lung capacity, high resolution CT scan, event.bronchoscopy with BAL). Induced sputum was obtained after inhalation of 5 ml of 3 percent solution of sodium chloride by ultrasonographic nebuliser. The induced sputum was cultivated, cytological and immunophenotypisation examinations were made. Pulmonary function tests and induced sputum examinations were repeatedly done after 9–12 months.

Results We have detected progression of pulmonary involvement in 16 patients. Cytological features of leucocyte predominance were found in 12 of them. In 29 non-progressive patients was leucocyte predominance found in only six cases. Respiratory infection was excluded.

Conclusion Method of induced sputum seems, after thoroughgoing appreciation, to be the interesting possibility of simple, non-invasive method for screening of pulmonary involvement progression in patients, suffering by connective tissue diseases.

REFERENCES

- 1 Olivieri D, DiPolito R, Chetta A. Induced sputum: diagnostic value in interstitial lung disease. *Curr Opin Pulm Med*. 2000;**6**(5):411–14
- 2 Bradna P, Parakova Z, Kral B, Hrnčíř Z, et al. Method of induced sputum in screening of pulmonary involvement in connective tissue diseases. *Rheumatologia* 2000;**14**(2):91

FRI0136 IMMUNE ADSORPTION IN HIGHLY ACTIVE SLE INCREASES THE RISK OF BACTERIAL, BUT NOT OF SEVERE VIRAL INFECTIONS

¹GH Stummvoll, ¹M Aringer, ²M Jansen, ²A Goldammer, ²K Derfler, ¹JS Smolen, ¹WB Graninger. ¹Rheumatology/Internal Medicine III; ²Nephrology/Internal Medicine III, University of Vienna, Vienna, Austria

10.1136/annrheumdis-2001.171

Background Active SLE patients undergoing plasmapheresis and cyclophosphamid therapy are at high risk of severe bacterial and viral infections. Immune adsorption (IAS) is a more novel approach to autoantibody removal with an unclear risk of infections in SLE.

Objectives We therefore decided to investigate the risk of infections in patients undergoing longterm IAS.

Methods We analysed the records of ten consecutive active SLE patients treated with long term IAS (mean 38 months) for the occurrence of severe infectious disease. We compared the IAS patients to historical control groups¹ of 9 patients under plasmapheresis and cyclophosphamid (PP) and 12 patients treated with cyclophosphamid (CX) alone for similar time periods.

Results One IAS patient died from gram-negative septicemia. Three localised bacterial infections (two gram-positive, one gram-negative) were treated with antibiotics. All infections occurred in the subgroup of five highly active patients (SIS >15, mean 17.7 ± 0.9) who all underwent additional CX therapy, showing characteristics similar to the CX and PP control groups (SIS 18.7 ± 2.4 and 20.3 ± 3.7, respectively). Bacterial infections were more frequent than in the CX group (4/5 vs 1/12, p < 0.01), but similar to the PP group. No severe viral infections were detected.

Conclusion IAS did not promote severe viral infections, but was associated with an increased rate of potentially life-threatening bacterial infections in SLE patients with very active disease.

REFERENCE

- 1 Aringer M, Smolen JS, Graninger WB. Severe infections in plasmapheresis-treated systemic lupus erythematosus. *Arthritis Rheum*. 1998;**41**(3):414–20

FRI0137 PROLACTIN IN SYSTEMIC LUPUS ERYTHEMATOSUS

¹L Moszkorova, ¹C Dostal, ²J Marek, ²Z Lacinova, ¹L Musilova. ¹1st Department, Institute of Rheumatology, Prague, Czech Republic; ²3rd Med. Department, 1st Med. Faculty Charles University, Prague, Czech Republic

10.1136/annrheumdis-2001.172

Background Prolactin (PRL) is an anterior pituitary hormone which is known to stimulate humoral and cell mediated immune responses, and may have a role in the pathogenesis of systemic lupus erythematosus (SLE).

Objectives To determine whether elevated serum prolactin associates the SLEDAI activity and/or specific organ involvement in SLE when recent reports have been till now discrepant.

Methods Basal serum PRL levels were determined in 55 consecutive SLE patients (pts) (49 females, 6 males) and in 41 healthy controls (31 females, 10 males) by RIA method. The estimated normal values of serum PRL were 200 mUI/l for males and 450 mUI/l for females.

Results The idiopathic hyperprolactinemia (hyper-PRL) was found in 7 SLE females (14,29%; 828,682 ± 391,410 mUI/l) and in 3 males (50,00%; 305,469 ± 65,751 mUI/l); and in one male in healthy controls (10,00%; 348,352 mUI/l) only. In remaining 42 SLE females and in 3 males PRL serum level was normal (365,783 ± 282,796 mUI/l; 142,854 ± 84,512 mUI/l). After 6 months repeatedly determined serum PRL in 2 females and 2 males with initial hyper-PRL, has shown normalisation in 1 female and 1 male. As to the proposed association between hyper-PRL and disease activity and specific organ involvement and presence of anti-ds-DNA there are some data on the Table 1.

Abstract FRI0137 Table 1

	Hyper-PRL	SLEDAI activity >4	Lupus nephritis (n = 15)	CNS (n = 11)	anti-ds-DNA (n = 28)
Females	7 (14%)	5 (71%)	3 (20%) *ns	0 *ns	3
Males	3 (50%)	2	3	1	2

Conclusion There was significantly higher frequency of hyper-PRL in 55 SLE pts compared to 41 healthy controls (p < 0,001). The value of serum PRL has normalised in 2 pts with hyper-PRL from 4 pts with initial hyper-PRL without any specific treatment after 6 months by repeated estimation. We document a new idiopathic hyper-PRL in 1 patient with normal initial serum PRL. The association between hyper-PRL and activity and/or specific organ involvement and presence of anti-ds-DNA was not statistically significant due to low number of observations.

REFERENCES

- 1 Walker SE, et al. Effect of prolactin in stimulating disease activity in SLE. *Proc NY Acad Sci* 1998;**840**:762–72

- 2 Neidhart M. Prolactin in autoimmune diseases. *Proc Soc Exp Bio Med.* 1998;**217**:408–79

FRI0138 ANTICARDIOLIPIN ANTIBODY ISOTYPE PROFILE AND LUPUS NEPHRITIS SEVERITY BASED ON RENAL BIOPSY PATHOLOGY

JC Tseng, CC Chiu, CM Hsu, RJ Hu, LY Lu, HH Cheng. *Allergy, Immunology and Rheumatology, Veterans General Hospital, Kaohsiung, Taiwan, R.O.C*

10.1136/annrheumdis-2001.173

Background Lupus nephritis remains the most prominent organ system manifestation of systemic lupus erythematosus (SLE) and is associated with high morbidity and mortality, particularly in the diffuse proliferative form of the World Health Organisation classification. Evidences have shown that antiphospholipid antibodies may be of pathogenetic significance in lupus nephritis, since glomerular capillary thrombi are an indicator of subsequent renal dysfunction.

Objectives To determine anticardiolipin antibody isotype profile in SLE patients with and without lupus nephritis, and the association with lupus nephritis severity based on renal biopsy classification.

Methods 200 Chinese SLE patients, followed between Jan. 1991 to May 1996 at Kaohsiung Veterans General Hospital Rheumatology Clinic were included in this study. All patients fulfilled ARA criteria for SLE. Anticardiolipin antibody (aCL) was measured by the ELISA according to international standardised method in 100 SLE patients with lupus nephritis verified by renal biopsy (Group A), 100 SLE patients without lupus nephritis (Group B), and 30 patients with renal biopsy-proved nephropathy not due to SLE (Group C). Renal biopsy morphology was assessed using WHO criteria for the classification of lupus nephritis. aCL isotype profile was analysed in each group and in the subgroups of lupus nephritis patients according to WHO classification.

Results

Conclusion In this study, the frequency of aCL in our SLE patients was low as compared to previous reports. It may be due to racial difference. From our preliminary data, though not statistically significant, no difference in the frequency of aCL was identified amongst patients with and without lupus nephritis. No correlation between aCL isotype profile and the severity of lupus nephritis could be demonstrated.

FRI0139 CLINICAL IMPORTANCE OF ASYMPTOMATIC BACTERIURIA IN WOMEN SUFFER FROM SYSTEMIC LUPUS ERYTHEMATOSUS – 5 YEARS OBSERVATION

¹M Koseda-Dragan, ¹M Hebanowski, ²I Szychlińska, ²J Galinski. ¹Connective Tissue Diseases Unit Department of Family Medicine; ²Department of Microbiology, Medical University of Gdansk, Gdansk, Poland

10.1136/annrheumdis-2001.174

Background Infections are frequent cause of death in lupus patients because of disease depending mechanisms and immunosuppressive treatment. Kidneys are often involved in course of lupus. We found asymptomatic bacteriuria in 16,5% women with SLE in first year of our study but longer 5 years observation could help us to understand its clinical importance.

Objectives

- Estimation of incidence of asymptomatic bacteriuria (a.b.) in population of women with SLE in 5 years observation.
- Examination of asymptomatic bacteriuria maintenance in following years of observation.
- Evaluation of incidence of symptomatic pyelonephritis in SLE women with a.b.
- Evaluation of need of antibacterial treatment in SLE patients with a.b.

Methods 85 women aged 24–77 (mean 45,2) years with mean SLE duration 10,2 years (2–33) were examined for 5 years. All of them were fulfilling ARA criteria for diagnosis of SLE. We did bacteriological analysis of SLE women urina 5 times in the 5 following years. Patients were observed clinically and parameters of biochemical and immunological activity were done. Among group with a.b. were counted women who had significant bacteriuria higher than 10^4 in ml urina. We were analysing type of bacteria and antibiotic resistancy.

Results In the following 5 years of observation asymptomatic bacteriuria was found with incidence from 12,8% to 22,5% (mean 17,6%). From 60% to 80% patients with a.b. (mean 70%) eliminated bacteria from urina in period one year without antibacterial treatment. In other patients mean 30% in second urina cultures which was done 1 year after first with a.b., significant bacteriuria was persistent. 50% of women with persistent bacteriuria after one year had in urina the same type of bacteria but in 50% patients type of bacteria was changed. In 5 years only 3 incidence of symptomatic pyelonephritis in SLE women with a. b. was observed. In all 3 cases nephrolithiasis was coexisted with SLE.

Conclusion

- Incidence of asymptomatic bacteriuria in women with SLE was 17,6%.
- 70% patients with a.b. eliminated bacteriuria without antibacterial treatment in one year.

Abstract FRI0138 Table 1

	Age, mean	Sex (F/M)	aCL+ (%)	aCL IgG+ (%)	aCL IgG+, mean (GPL/dl)	aCL IgA+ (%)	aCL IgA+, mean (APL/dl)	aCL IgM+ (%)	aCL IgM+, mean (MPL/dl)
Group A (SLE with nephritis)	29.61	89/11	17	15	32.8	1	31.0	11	16.7
Group B (SLE without nephritis),	34.77	94/6	14	11	28.0	5	37.4	10	30.0
Group C (non-SLE nephropathy),	48.3	10/20	3	2	22.0	1	51.0	1	65.0

Anticardiolipin antibody isotype profile.