Behcet's disease is an uncommon multisystemic inflammatory disorder characterised by recurrent oral aphthous ulcers, genital ulcers, uveitis, and skin lesions. Involvement of the large vessels is less frequent. Vascular involvement in Behcet's disease is divided into venous and arterial thrombosis and arterial aneurysmal formation. Arterial involvement of Behcet's disease most frequently affects the abdominal aorta, followed by the femoral and the pulmonary arteries.

Results In this case, he fulfils the described criteria for Behcet's disease by the International Study Group for Behcet's disease in 1990. Our diagnosis of Behcet's disease was based mainly on the recurrent oral ulceration, recurrent genital ulceration, eye lesion, skin lesion, and positive HLA-B51 and pathergy test.

Conclusion To our knowledge, the present case is the first patient described with Behcet disease who developed celiac trunk aneurysms. We should be aware of this complication in patients with Behcet's disease who develop abdominal pain.

REFERENCES

- 1 Junewick JJ, Grant TH, Weiss CA, *et al.* Celiac artery aneurysm: color Doppler evaluation. *J Ultrasound Med.* 1993;**12**:355
- 2 Sakane T, Takeno M, Suzuki N, Inaba G. Behcet's disease. *N Engl J Med.* 1999;**341**(17):1284-91
- 3 Vohra R, Carr HM, Welch M, *et al.* Management of celiac artery aneurysms. *Br J Surg.* 1991;**78**:1373
- 4 Lukes P, Wihed A, Tidebrant G, *et al.* Angiography of visceral aneurysms. *Eur Radiol.* 1994;**4**:75
- 5 International Study Group for Behcet's Disease. Criteria for diagnosis of Behcet's disease. *Lancet* 1990;**335**:1078-80

AB0083 TAKAYASU ARTERITIS WITH BILATERAL RENAL ARTERY STENOSIS IN A 17-YEAR-OLD GIRL

¹K Aksu, ¹G Keser, ¹V Inal, ²A Memis, ¹F Oksel, ³N Hizli, ¹Y Kabasakal, ¹G Gümüşdis, ¹E Doganavsargil. ¹Rheumatology; ²Radiodiagnosics; ³Rheumatology, Atatürk State Hospital, Izmir, Turkey

10.1136/annrheumdis-2001.190

Background Takayasu arteritis, commonly seen in young women, is a large vessel vasculitis, primarily affecting aorta and its major branches. Hypertension frequently accompanies Takayasu arteritis and in approximately 75% of the cases it is due to renal artery involvement, causing stenosis. Hereby, we report a 17-year-old girl with Takayasu arteritis, having bilateral renal artery stenosis, with unilateral atrophic and afunctional kidney.

This patient, who was diagnosed as Takayasu arteritis 3 years ago, with typical physical findings such as absent upper extremity peripheral pulses and classical angiographical findings involving arcus aorta and its major arteries, has been referred to our rheumatology department because of uncontrolled hypertension. Renal Doppler ultrasound studies and angiography revealed bilateral renal artery stenosis with atrophic right kidney. Successful angioplasty, followed by stent replacement corrected the stenosis in the left renal artery. However, the stenosis in the right renal artery involved a longer segment and angioplasty or stent replacement was not possible. From technical point of view, there was no way for surgical bypass for the stenosis in the right renal artery. Furthermore, no function was observed in the right kidney during renal scintigraphy. Nephrectomy of the right kidney was planned, but it was not accepted by the patient. Since the left renal artery stenosis was corrected, renovascular hypertension maintained by the right renal artery stenosis was treated successfully by captopril and the patient was closely monitored with respect to renal function tests.

In conclusion, renal artery stenosis should always be kept in mind in Takayasu arteritis with hypertension.

Objectives

Methods

Results

Conclusion

AB0084 BEHÇET'S DISEASE AND BUDD-CHIARI SYNDROME: TWO CASES WITH DIVERSE PROGRESSES

¹K Aksu, ¹G Keser, ¹V Inal, ¹F Oksel, ¹H Kocanaogullari, ²Y Tokat, ³Y Batur, ¹G Gümüşdis, ¹E Doganavsargil. ¹Rheumatology; ²General Surgery; ³Gastroenterology, Ege University, Faculty of Medicine, Izmir, Turkey

10.1136/annrheumdis-2001.191

Background Behçet's disease (BD) is a systemic vasculitis of unknown aetiology that can lead to thrombosis of both veins and arteries of various sizes. Thrombosis of hepatic veins, known as Budd-Chiari syndrome (BCS) may be seen in BD, with variable clinical courses as described in 2 patients:

Case 1: Thirty-four year-old male patient had oral and genital ulcers and erythema nodosum for 2 years. Within two weeks after presentation, thrombosis of hepatic veins and vena cava inferior was found, while evaluating the patient for abdominal pain and liver function test abnormalities. The patient well responded to the treatment consisting high dose prednisolone, pulse cyclophosphamide and anticoagulant drugs with recanalisation of the veins. At the end of two years, the patient has still been in clinical remission.

Case 2: A twenty year-old male patient had oral and genital ulcerations for 4 years and had an episode of uveitis 4 years ago. He was referred to our clinic because of abdominal distension and oedema of lower extremities lasting for 4 months. The laboratory analyses disclosed ascites, grade III varicosity of oesophagus and thrombosis of hepatic veins and vena cava inferior. Mesoatrial shunt operation was performed when the initial treatment with pulse cyclophosphamide and high dose prednisolone failed. Obliteration of the shunt was observed by doppler ultrasonography, performed one week after the operation. He did not respond to the supportive therapy after the shunt operation and was lost with gastrointestinal bleeding.

Objectives

Methods

Results

Conclusion These two male Behçet's patients with BCS are examples of diverse prognoses, probably depending upon how early the diagnosis is made, and how efficient the treatment is. BCS in BD, may have a better prognosis if the diagnosis is made earlier and the treatment started sooner in the course of the disease.

AB0085 THE EFFECTIVENESS OF MELOXICAM IN TREATING ERYTHEMA NODOSUM SYNDROME

El Chmidt, KV Zhukova. *Rheumatology, The 1-St City Hospital, Moscow, Russia*

10.1136/annrheumdis-2001.192

Background

Objectives We used meloxicam at a dose of 15 mg a day for treating erythema nodosum (EN) of various aetiology in 47 patients (44 females and 3 males aged 15-68). Not depending

on the EN aetiology, the main symptoms were fever (in 38 patients), joint pain (in 41 patients), foot and leg oedema (in 44 patients). The duration of EN syndrome before beginning meloxicam was from 2 weeks to 4 months. During this period all the patients underwent therapy by various NSAIDs (mainly diclofenac, ketoprofen, indometacin), 4 patients by steroids, antibiotics. In all the patients the previous therapy was either non-effective or little effective. Using meloxicam as a monotherapy on the 2-nd, 3-d day of taking the drug resulted in normalisation of temperature, disappearance of joint pain and oedema, quick decreasing of EN even in those patients who had no effect from steroids. In 4 patients the attempt of cancelling meloxicam lead to relapse of EN, fever and joint pain, which demanded resuming of meloxicam treatment. 4 patients had an allergy in the form of skin eruption and itch. That's why meloxicam in those patients was cancelled in spite of a good effect. Meloxicam turned out to be ineffective in 2 patients. In 45 patients (95.7%) with pronounced EN, fever and oedema meloxicam showed great effectiveness.

Methods

Results

Conclusion

AB0086 OCCURRENCE OF GIANT CELL ARTERITIS UNDER METHOTREXATE THERAPY FOR RHEUMATOID ARTHRITIS

J Gerster, I Carey, AM Chamot. *Department of Rheumatology, University Hospital (CHUV), Lausanne, Switzerland*

10.1136/annrhumdis-2001.193

Background There is controversy about methotrexate (MTX) as a therapy of giant cell arteritis (GCA).

Objectives The objective is to present a case who developed GCA during therapy with MTX in association with corticosteroids for rheumatoid arthritis.

Methods A 66-year-old-woman was diagnosed in 1998 as sero-negative rheumatoid arthritis involving knee, shoulder and wrist joints. ESR was 46 mm/1 h, synovial fluid contained 29 G/L leukocytes but no crystals were observed. The patient was treated with MTX at a weekly dose of 7,5 mg, dosage increased to 10 mg/week in February 2000. In addition, she received prednisone 5 mg per day. In November 2000 she started to feel some weakness and to suffer from frontal headaches. The ESR was 120 mm/1 h, CRP 121 mg/l and Hb 80 g/L. The biopsied right temporal artery, which was tender and indurated, demonstrated GCA. The internal elastic lamina was fragmented; there were infiltrates of histiocytes, lymphocytes and giant cells and the lumen was in some places occluded.

Results

Conclusion In this case, occurrence of GCA was not prevented by MTX therapy. This suggests that MTX at an average dose is probably insufficient to prevent or treat this disease.

AB0087 TAKAYASU'S ARTERITIS: REVIEW AND FOUR CLINICAL CASES

G Sequeira, F Saraiva, A Marques, J Romeu, T Costa, V Queiroz. *Medicine IV C, Hospital de Santa Maria, Lisbon, Portugal*

10.1136/annrhumdis-2001.194

Background Takayasu's arteritis is a rare disease, mainly affecting young women. The authors describe four cases of Takayasu's disease from the rheumatology outpatient clinic of Hospital de Santa Maria in Portugal. A revision of the disease is made concerning its clinical manifestations, laboratory, diagnosis, treatment and prognosis.

Objectives

Methods

Results All patients were female, three were white and one black. Beginning of symptoms was between 7 and 56 years, satisfying the American College of Rheumatology criteria for the classification of Takayasu's arteritis. The spectrum of presentation, disease activity and pace of disease progression were very variable. The delay between the onset of first symptoms and diagnosis was less than three years in three patients and twenty in the other. The first symptoms of disease were in two patients systemic reactions such as malaise, fever and anorexia, absent pulses in the lower limbs in another and claudication of the upper extremities in the last. Vascular ischaemic symptoms, hallmark of the disease, were present in the four patients and all of them had angiograms that involved visualisation of at least primary tributaries that showed lesions of the aortic arch and its branches. All patients needed corticosteroid treatment in the acute inflammatory phase of the disease. Cytotoxic agents were used in three of them and in two cases vascular surgery was done because of fear of occlusion of the involved vessels.

Conclusion Females and caucasian race predominated. The diagnosis of Takayasu's arteritis should be considered in the presence of symptoms of vascular ischemia such as pulse defects or signs of bruits and in the angiographic study. The prominence of inflammatory features antedating obvious symptoms often delayed diagnosis for substantial periods of time. Fever, myalgia/arthralgia with no identified cause, elevated erythrocyte sedimentation rate, claudication, diminished or absent pulses, bruits, vascular pain and the typical angiographic features are criteria for disease activity. The mainstay of treatment is oral corticosteroids. They are very effective in controlling clinical manifestations and arresting disease progression and sometimes in achieving return of previously absent pulses. Cytotoxic therapy has been helpful in patients failing to respond to corticosteroids.

AB0088 PULSE CYCLOPHOSPHAMIDE THERAPY IN A PATIENT OF GIANT CELL ARTERITIS WITH DIPLOPIA DUE TO UNILATERAL ABDUCENS NERVE PALS

JC Tseng, HH Cheng, RJ Hu, LY Lu. *Allergy, Immunology and Rheumatology, Veterans General Hospital, Kaohsiung, Taiwan, R.O.C*

10.1136/annrhumdis-2001.195

Background Patients with giant cell arteritis (GCA) occasionally complain of double vision. The neurologic manifestations of the GCA are protean. All parts of the visual pathway from the cornea to the cortex have been described to be involved.

Objectives We described a patient of GCA with diplopia due to left abducens nerve palsy which was refractory to corticosteroid therapy initially and recovered with monthly pulse cyclophosphamide therapy five months later. To the best of our knowledge this is the first case described in the medical literature to be treated with monthly pulse intravenous cyclophosphamide and to achieve a successful outcome.

Methods

Results