Fibromyalgia (FM) is a syndrome of unknown aetiology characterised by chronic wide spread pain, increased tenderness to palpation and additional symptoms such as disturbed sleep, stiffness, fatigue and psychological distress. Familial aggregation of FM is common. Depressive symptoms and psychological distress is a frequent finding in FM patients and their relatives.¹ These findings suggest that genetic and familial factors may play a role in the etiopathogenesis of this syndrome. In a recent linkage analysis of 40 FM multicase families it was shown that the HLA region might be a candidate for genetic transmission of vulnerability for the disorder.²

The pathophysiological mechanisms underlying fibromyalgia have been linked to disturbances in serotonin (5-HT) metabolism and transmission. Among the different parameters of the serotonin pathway the genes coding for the 5-HT receptors and transporters are interesting candidates to investigate a genetically driven predisposition to FM. Recent studies have analysed silent and functional polymorphic variants of these genes in patients with fibromyalgia.^{3,4,5}

- The human 5-HT transporter (5-HTT) is encoded by a single gene on chromosome 17q12. A functional polymorphism in the transcriptional region has been described ("long allele" L or "short allele" S). In a study of 62 FM patients a higher frequency of the S/S genotype of the 5-HTT was found compared to healthy controls. The S/S subgroup revealed higher means in depression and psychological distress.
- The 5-HT_{2A} receptor gene is located on chromosome 13q14–21. A silent T102C polymorphism of the 5-HT_{2A} receptor gene has recently been detected. In a sample of 168 FM patients a statistically significant different distribution of genotypes was found with a decrease in T/T and an increase in both T/C and C/C genotypes compared to controls. A correlation of genotypes with pain variables revealed higher scores in the tenderpoints score in patients with the T/T genotype. Correlations with psychopathological symptoms were not significant.
- The 5-HT_{2C} receptor gene is located on the X chromosome. In a sample of 65 FM patients the presence of the 5-HT_{2C} (C32S) mutant was significantly correlated with measures of psychological distress. In the same sample the tryptophan hydroxylase (TPH) gene was analysed. The TPH gene is located on the short arm of chromosome 11 and a polymorphism in intron 7 has recently been detected. The TPH is the rate limiting enzyme that determines the concentrations of serotonin *in vivo*. No significant different frequencies of the alleles were found in this group of FM patients compared to control subjects.

The implication of these findings for the etiopathogenesis of fibromyalgia is far from clear and have to be replicated in larger and ethnical different samples. But these results might point to a disturbance on 5-HT-receptor and transporter level at least in a subgroup of FM patients and emphasises the importance of the serotonergic system in the pathophysiology of FM.

REFERENCES

- 1 Offenbaecher M, Glatzeder K, Ackenheil M. Self reported depression, familial history of depression, fibromyalgia (FM) and psychological distress in patients with FM. *Z Rheumatol*. 1998;**57**:94–6
- 2 Yunus MB, Khan MA, Rawlings KK, Green JR, Olson JM, Shah S. Genetic linkage analysis of multicase families with fibromyalgia syndrome. *J Rheumatol* 1999;**26**:408–12
- 3 Offenbaecher M, Bondy B, de Jonge S, Glatzeder K, Krueger M, Schoeps P, Ackenheil M. Possible association of fibromyalgia with a polymorphism in the serotonin transporter gene regulatory region. *Arthritis Rheum* 1999;**42**:2482–8
- 4 Bondy B, Spaeth M, Offenbaecher M, Glatzeder K, Stratz T, Schwarz M, de Jonge S, Krüger M, Engel R, Faerber L, Pongratz D, Ackenheil M. The T102C polymorphism of the 5-HT_{2A}-receptor gene in fibromyalgia. *Neurobiol Dis*. 1999;**6**:433–9

- 5 Offenbaecher M, Bondy B, de Jonge S, Glatzeder K, Behrens S, Schoeps P, Ackenheil M. Evidence for involvement of 5-HT_{2A} and 5-HT_{2C} receptor genes in fibromyalgia. Poster presented at the 9th world congress on pain, 1999

SP0064 DRUG THERAPY IN FIBROMYALGIA. A REVIEW

E Thomas, F Blotman. *Rheumatology, Lapeyronie Hospital, Montpellier, France*

10.1136/annrheumdis-2001.20

Since the cause and pathophysiology of fibromyalgia (FM) remain unknown, treatment is often limited; fewer than 50% of patients have significant and sustained clinical improvement. No pharmacologic agent are indicated specifically for FM treatment, although many have been tried to date. Controlled trials with non-steroidal anti-inflammatory drugs have been shown them to be of little help, despite their use in a majority of patients. Steroidal anti-inflammatory drugs are not efficacious and often deleterious. Among analgesics, preliminary data suggest that tramadol, affecting norepinephrine at the receptor site, may be useful for FM pain. No clinical data are available to suggest efficacy of opioids in FM. Tricyclic antidepressants and selective serotonin reuptake inhibitors may diminish the sleep disturbance, pain and depression associated with FM. they remain the cornerstone of drug therapy in FM. Blockade of the serotonin receptor subtype 5-HT₃ with Tropisetron or Ondansetron has been reported to significantly decrease the number of tender points, despite poor digestive tolerance. The findings of low levels of insulin-like growth factor 1 in FM led to give daily growth hormone (GH) injections to 22 women (vs placebo to 23), enrolled in a controlled study. A 9 months, the tender point score was significantly improved with GH. other agents like 5-hydroxytryptophan, gamma-hydroxybutyrate and S-adenosyl-L-Methionine also shown promising results. FM symptoms may be alleviated in a short term manner with various pharmacologic treatments but a minority of patients still experience remission at a few years follow-up.

SP0065 FIBROMYALGIA: PAIN MANAGEMENT STRATEGIES

¹C Cedraschi, ²J Desmeules, ¹TL Vischer. ¹Rheumatology Division, Geneva, Switzerland; ²Clinical Pharmacology and Toxicology Division, Geneva University Hospital, Geneva, Switzerland

10.1136/annrheumdis-2001.21

At present, there is no specific treatment for FM, partly because of the lack of understanding of its aetiology and pathogenesis. Self-management based programs have been proposed to help the patients cope with their symptoms and improve their quality of life. These programs mainly use information, physical conditioning, and cognitive-behavioural techniques.

Information in chronic pain patients has been emphasised as an important component as a well-informed patient is an essential member of a successful chronic illness management team. Acknowledging FM patients' pain and suffering is also a key point in these programs.

FM patients have been shown to have functional impairment which include their ability to perform daily activities. They have decreased muscle strength and endurance, and low levels of physical fitness. Exercise programs are various and may include range-of-motion exercises, cycling, walking, swimming, specific muscular reinforcement, etc. Pool exercises have been used in