



OPEN ACCESS

EXTENDED REPORT

Comprehensive disease control (CDC): what does achieving CDC mean for patients with rheumatoid arthritis?

Paul Emery,¹ Arthur Kavanaugh,² Yanjun Bao,³ Arijit Ganguli,³ Parvez Mulani³

Handling editor Tore K Kvien

¹Leeds Institute of Rheumatic & Musculoskeletal Medicine, University of Leeds, Chapel Allerton Hospital and NIHR Leeds Musculoskeletal Biomedical Research Unit, Leeds Teaching Hospitals NHS Trust, Leeds, UK

²University of California San Diego, La Jolla, California, USA
³AbbVie Inc, North Chicago, Illinois, USA

Correspondence to

Professor Paul Emery, Leeds Institute of Rheumatic & Musculoskeletal Medicine, University of Leeds, Chapel Allerton Hospital and NIHR Leeds Musculoskeletal Biomedical Research Unit, Leeds Teaching Hospitals NHS Trust, Chapeltown Road, Leeds LS7 4SA, UK; manuscripts@jkmed.com

Received 27 January 2014
Revised 10 June 2014
Accepted 26 July 2014
Published Online First
19 August 2014



Open Access
Scan to access more
free content



CrossMark

To cite: Emery P, Kavanaugh A, Bao Y, et al. *Ann Rheum Dis* 2015;**74**:2165–2174.

ABSTRACT

Background This study assessed the impact of simultaneous achievement of clinical, functional and structural efficacy, herein referred to as comprehensive disease control (CDC), on short-term and long-term work-related outcomes, health-related quality of life (HRQoL), pain and fatigue.

Methods Data were pooled from three randomised trials of adalimumab plus methotrexate for treatment of early-stage or late-stage rheumatoid arthritis (RA). CDC was defined as 28-joint Disease Activity Score using C reactive protein <2.6, Health Assessment Questionnaire <0.5 and change from baseline in modified Total Sharp Score ≤0.5. Changes in scores at weeks 26 and 52 for work-related outcomes, Short Form 36 (SF-36) physical (PCS) and mental component scores (MCS), a Visual Analogue Scale measuring pain (VAS-Pain) and Functional Assessment of Chronic Illness Therapy-Fatigue (FACIT-F) were compared between patient groups defined by achievement of CDC at week 26 using linear regression with adjustment for baseline scores.

Results Patients with RA who achieved CDC at week 26 (n=200) had significantly greater improvements in VAS-Pain (46.9 vs 26.9; p<0.0001), FACIT-F (13.3 vs 7.5; p<0.0001), SF-36 PCS (19.7 vs 8.9; p<0.0001) and SF-36 MCS (8.1 vs 5.0; p=0.0004) than those who did not (n=1267). Results were consistent at week 52 and among methotrexate-naïve patients with early RA, methotrexate-experienced patients with late-stage RA and patients with inadequate response to methotrexate.

Conclusions Patients with RA who achieved CDC at week 26 had improved short-term and long-term HRQoL, pain, fatigue and work-related outcomes compared with patients who do not. These results demonstrate that the joint achievement of all CDC components provides meaningful benefits to patients.

Trial registration numbers DE019: NCT00195702, PREMIER: NCT00195702, OPTIMA: NCT00195702.

INTRODUCTION

Rheumatoid arthritis (RA) is characterised by progressive inflammation of the joints and leads to irreversible joint damage that reduces function and increases disability.^{1 2} Current treatment options have shown to reduce inflammation, slow or prevent joint damage and improve patient-related outcomes. As per the treat-to-target recommendations, the primary goal in the treatment of RA should be achieving the control of symptoms, prevention of structural damage, normalisation of function and social participation.³ In the European

League Against Rheumatism (EULAR) 2013 guidelines, achievement of clinical, functional and structural efficacy is also the basis for treatment comparison. However, the impact of such achievement on patient outcomes has not been adequately and explicitly assessed.

While the achievement of all clinical, functional and structural efficacy is the ultimate goal for disease management, in real world practice, remission has been recommended as a key treatment target for adjusting therapy to optimise outcomes accordingly.³ Over the years, remission has been defined in several ways, but even with the recently more stringent American College of Rheumatology (ACR)-EULAR criteria, nearly 23% of the patients who achieved ACR-EULAR remission did not achieve the desired radiographic inhibition.⁴ It is, therefore, not guaranteed that remission target achievement alone would ensure functional and structural goal achievement. While remission could be the target for adjusting therapy, the goal of every treatment should be inhibition of structural damage and normalise function, in addition to symptom control.

In defining the three elements of the treatment goal, there has been a greater evolution in the definition of clinical remission relative to what is defined for normalised function based on Health Assessment Questionnaire disability index (HAQ-DI) and for structural non-progression based on modified Sharp score. Remission criteria derived from Disease Activity Score (DAS) and 28-joint Disease Activity Score (DAS28) have been the dominant treatment target in clinical practice in this evolution. Recently developed ACR-EULAR remission is so stringent that it is difficult to be achieved by most patients especially those with long-standing disease who actually constitute the majority of patients in clinical care. Hence, the EULAR 2013 guidelines recommended less stringent measures as a good alternative target for many patients who cannot attain stringent forms of remission.⁵ Given this, in the current study, we define clinical remission based on DAS28. A cut-off point of DAS28 using C reactive protein (DAS28(CRP)) <2.6 is an acceptable, as well as stringent enough, quantitative treatment target for the signs and symptoms of RA inflammation.⁶

Patient-reported outcomes (PROs) provide important assessments of functioning and well-being from the patient's perspective and may provide additional tangible impact for a patient,

and thus be more sensitive to the effects of treatment than physician-assessed measures.^{7–10} In some cases, work limitations are important considerations in the treatment of RA given that the disease affects patients in their productive years.^{11–12} RA has also been noted to critically impact the health-related quality of life (HRQoL) of patients through the clinical manifestations of the disease and by socioeconomic, personal and environmental factors.^{7–13} Pain and fatigue are other important features of RA which are especially relevant from the patient perspective.^{8–14–16}

In this study, we sought to quantify the impact of simultaneous achievement of clinical, functional and structural efficacy on work-related outcomes, HRQoL, pain and fatigue by pooling data from three randomised clinical trials of adalimumab for the treatment of RA among patients with both early-stage and late-stage disease.

PATIENTS AND METHODS

Patients

Data were drawn from three separate randomised controlled trials in RA: DE019 (NCT00195702), which enrolled patients with experienced RA (disease duration of at least 3 years with moderately to severely active RA despite a minimum of 3 months of treatment with methotrexate (MTX)); PREMIER (NCT00195663), which enrolled MTX-naïve patients with early RA (disease duration <3 years) and OPTIMA (NCT00420927), which enrolled patients with early RA and included patients who failed to achieve stable low disease activity (LDA) (DAS28(CRP) <3.2 at weeks 22 and 26) with MTX. The design and primary results of the DE019, PREMIER and OPTIMA trials have been described elsewhere.^{17–20}

A pooled intent-to-treat population included all patients from these trials who were randomised and received at least one dose of study drug. The present analysis includes all patients from DE019 and PREMIER. Due to the rerandomisation in the OPTIMA trial, only patients who failed to achieve LDA (DAS28 (CRP) <3.2) with MTX during Period 1 of OPTIMA were included in the analysis. For these patients, the beginning of Period 2 was considered the baseline period. Patients with missing DAS28(CRP), HAQ-DI or change in modified total Sharp score (Δ mTSS) at baseline or week 26 were excluded from the present study.

The simultaneous achievement of stringent control of the signs and symptoms of inflammation, normal physical function and the absence of radiographic progression, herein referred as comprehensive disease control (CDC), was evaluated at week 26. Stringent control of the signs and symptoms of inflammation was assessed by DAS28(CRP) and defined as DAS28(CRP) <2.6.^{2–6–21} Normal physical function was assessed using the Disability Index of the HAQ-DI²² and defined as HAQ-DI <0.5.^{4–23} Radiographic progression was assessed using radiographs of the hands/wrists and feet scored using the mTSS method.^{24–26} Absence of radiographic progression was defined as a ≤ 0.5 unit change from baseline in mTSS.²⁷ Patients who achieved all three components were considered to have achieved CDC ('CDC achievers'), while patients who achieved either none or any one or two (partial achievers) of the three components were considered to have not achieved CDC ('CDC non-achievers').

Outcomes

Work-related outcomes were measured in DE032, the economic companion to PREMIER, and in OPTIMA. Work outcomes were analysed in each trial separately because of different definitions across the trials. Work outcomes in PREMIER/DE032

were measured using the Patient Health Economic Questionnaire (P-HEQ) and included the number of work days missed and the level at which work performance has been affected for employed patients with RA and homemakers separately. The level at which work performance has been affected was measured using a Visual Analogue Scale (VAS) scale ranging from 0 to 100.²⁸ Work outcomes in OPTIMA were measured using the Work Productivity and Activity Impairment (WPAI) questionnaire²⁹ and the RA-Work Instability Scale (RA-WIS).³⁰ WPAI consists of six questions regarding the ability to work and perform regular activities during the past 7 days. WPAI evaluates absenteeism, presenteeism and work impairment among employed patients and activity impairment among all patients. RA-WIS is a 23-item questionnaire which quantifies 'the state in which the consequences of a mismatch between an individual's functional abilities and the demands of his or her job can threaten continuing employment if they are not resolved'.³⁰

HRQoL was measured using the Short Form 36 (SF-36) health survey. The SF-36 is a 36-item generic HRQoL measure to assess the patient's view of his or her health consisting of two summary scales: physical (PCS) and mental component scores (MCS). For each component, a normalised summary score was calculated such that the average in the general population is 50 (with an SD of 10) using eight subdomains: Physical Functioning, Role-Physical, Bodily Pain and General Health for PCS, and Vitality, Social Functioning, Role-Emotional and Mental Health for MCS. Scores range from 0 to 100, with greater scores reflecting better health status.^{31–34} The minimal clinically important difference (MCID) for the SF-36 PCS and MCS is a 2.5-point increase from baseline.³⁵ SF-36 information was not collected in the OPTIMA trial.

Pain was measured using a VAS ranging from no pain to worst pain imaginable, with a higher score indicating worse pain. The MCID for VAS-Pain was a 10-mm decrease from baseline.³⁶ Fatigue was measured using the Functional Assessment of Chronic Illness Therapy-Fatigue (FACIT-F), a symptom-specific scale validated for patients with RA. The 13-item subscale is self-administered using a 5-point Likert rating scale and ranges between 0 and 52 with higher values indicating less fatigue.³⁷ The MCID for FACIT-Fatigue is a 4-point increase from baseline.³²

Statistical analysis

Baseline characteristics were described for patients who achieved and did not achieve CDC at week 26. The change from baseline to week 26 for each outcome variable was compared between CDC achievers and non-achievers using linear regression adjusting for the baseline value of each outcome. Work outcomes measured using the P-HEQ were analysed in PREMIER only, while work outcomes measured using WPAI and RA-WIS were analysed in OPTIMA only. Similar analyses were performed to compare change from baseline to week 52. VAS-Pain and FACIT-F were analysed by pooling data from all three randomised clinical trials and in each trial separately. SF-36 PCS and SF-36 MCS were analysed by pooling DE019 and PREMIER because these HRQoL measures were not collected in OPTIMA. Separate analyses were conducted within the early RA (PREMIER), the MTX-failure (OPTIMA) and the experienced RA (DE019) populations. To gauge the incremental benefit of the simultaneous achievement of the three CDC components, the differences in SF-36 PCS, MCS, FACIT-F and VAS-Pain were compared between patients achieving all three components compared with achieving each component individually using a full interaction model adjusting for the baseline value of each

Table 1 Patient characteristics of week 26 CDC achievers and non-achievers

	CDC achievers N=200	Non-achievers N=1267
Demographics		
Age, mean±SD	48.5±13.3	53.9±13.0
Female, n (%)	123 (61.5)	973 (76.8)
Trial, n (%)		
OPTIMA	58 (29.0)	255 (20.1)
PREMIER	98 (49.0)	563 (44.4)
DE019	44 (22.0)	449 (35.4)
Study outcomes, mean±SD		
Continuous CDC components		
DAS28	5.2±1.3	5.8±1.1
HAQ-DI	0.9±0.6	1.5±0.6
mTSS	20.9±31.6	37.1±44.7
Patient-reported outcomes		
SF-36 physical component score*, †	34.2±8.9	29.7±8.0
SF-36 mental component score*, †	48.2±12.4	45.9±12.0
VAS-Pain*	47.8±25.9	56.3±23.5
FACIT-Fatigue‡	35.3±10.8	29.0±11.2
Lab values		
Haemoglobin (g/dL)	13.1±1.5	12.8±1.5
Creatine (mg/dL)	0.8±0.2	0.7±0.2
Work-related outcomes in OPTIMA§		
Absenteeism*	9.6±23.1	17.3±30.9
Presenteeism*	25.9±25.2	45.6±27.2
Activity impairment*	26.9±24.2	47.8±23.9
Work impairment*	29.4±26.5	49.9±30.9
Work instability scale	6.7±6.6	11.7±7.2
Work-related outcomes in PREMIER/DE032¶		
Number of work days missed among employed patients in the last 4 weeks	3.2±6.1	5.2±8.6
Level work performance has been affected among employed patients in the last week*	45.6±32.6	53.5±32.0
Number of work days missed among homemakers in the last 4 weeks	4.7±8.3	7.2±10.1
Level work performance has been affected among homemakers in the last week*	43.6±26.8	62.4±26.4
Additional measures		
Clinical characteristics, mean±SD		
RA duration (years)	2.7±5.1	4.4±7.3
Swollen joint count (66)	16.1±8.0	19.8±9.8
Tender joint count (68)	21.1±10.2	29.0±13.7
C reactive protein (mg/L)	20.1±23.7	27.7±33.6
Physician's global assessment of disease activity*	58.7±18.5	64.5±17.2
Patient's global assessment of disease activity*	55.0±25.6	60.6±23.3
Radiographic findings, mean±SD		
Erosion score	12.6±18.5	20.9±25.0
Joint space narrowing	8.3±14.3	16.3±21.4
Components of CDC achievement at week 26, n (%)		
Full achievement	200 (100.0)	0 (0.0)
Achieved HAQ <0.5 and DAS <2.6	0 (0.0)	58 (4.6)
Achieved HAQ <0.5 and ΔmTSS ≤0.5	0 (0.0)	216 (17.0)
Achieved DAS <2.6 and ΔmTSS ≤0.5	0 (0.0)	89 (7.0)
Achieved HAQ <0.5 only	0 (0.0)	95 (7.5)
Achieved DAS <2.6 only	0 (0.0)	33 (2.6)
Achieved ΔmTSS ≤0.5 only	0 (0.0)	495 (39.1)
Achieved none	0 (0.0)	281 (22.2)
ACR-EULAR remission achievement at week 26, n (%)		
Yes	65 (33.0)	29 (2.3)

*Measured from 0 to 100.

†Not measured in OPTIMA patients.

‡Measured from 0 to 52. Greater values indicate less fatigue.

§Among OPTIMA patients only.

¶Among PREMIER/DE032 patients only.

ACR-EULAR, American College of Rheumatology-European League Against Rheumatism; CDC, comprehensive disease control; DAS, Disease Activity Score; FACIT, Functional Assessment of Chronic Illness Therapy; HAQ-DI, Health Assessment Questionnaire disability index; RA, rheumatoid arthritis; SF-36, Short Form 36; VAS, Visual Analogue Scale; mTSS, modified total Sharp score; ΔmTSS, change in modified total Sharp score.

Table 2 Comparison of patient-reported and work-related outcomes by week 26 CDC achievement*†

Patient-reported outcomes in the pooled population				
	CDC achievers (A) N=200 Mean (95% CI)	Non-achievers (B) N=1267 Mean (95% CI)	Difference (C)=(A)-(B) Mean (95% CI)	p Value
Week 26				
VAS-Pain change, N=1463	-46.91 (-49.83 to -43.98) n=200	-26.88 (-28.04 to -25.72) n=1263	-20.03 (-23.18 to -16.88)‡	<0.0001
FACIT-Fatigue change, N=1452	13.29 (12.03 to 14.54) n=196	7.51 (7.02 to 8.00) n=1256	5.78 (4.42 to 7.13)‡	<0.0001
SF-36 mental change, N=1086§	8.06 (6.45 to 9.66) n=131	4.98 (4.39 to 5.57) n=955	3.07 (1.36 to 4.78)‡	0.0004
SF-36 physical change, N=1086§	19.66 (18.13 to 21.19) n=131	8.91 (8.35 to 9.47) n=955	10.76 (9.12 to 12.39)‡	<0.0001
Week 52				
VAS-Pain change, N=1306	-45.68 (-48.76 to -42.61) n=189	-28.83 (-30.08 to -27.57) n=1117	-16.86 (-20.18 to -13.53)‡	<0.0001
FACIT-Fatigue change, N=1326	12.98 (11.66 to 14.30) n=186	8.05 (7.52 to 8.57) n=1140	4.93 (3.51 to 6.36)‡	<0.0001
SF-36 mental change, N=1001§	7.26 (5.61 to 8.92) n=126	5.12 (4.49 to 5.74) n=875	2.15 (0.38 to 3.91)	0.0174
SF-36 physical change, N=1001§	19.37 (17.73 to 21.01) n=126	10.03 (9.42 to 10.65) n=875	9.34 (7.57 to 11.10)‡	<0.0001
Work-related outcomes in PREMIER/DE032††				
	CDC achievers (A) N=98 Mean (95% CI)	Non-achievers (B) N=563 Mean (95% CI)	Difference (C)=(A)-(B) Mean (95% CI)	p Value
Week 26				
P-HEQ absenteeism among employed patients, N=284	9.69 (3.36 to 16.02) n=54	11.87 (8.82 to 14.93) n=230	-2.19 (-9.22 to 4.85)	0.5414
P-HEQ presenteeism among employed patients, N=278	-44.23 (-50.30 to -38.17) n=52	-29.94 (-32.84 to -27.04) n=226	-14.30 (-21.03 to -7.56)	<0.0001
P-HEQ absenteeism among homemakers, N=182	2.30 (-5.88 to 10.48) n=24	8.51 (5.32 to 11.69) n=158	-6.21 (-14.99 to 2.58)	0.1650
P-HEQ presenteeism among homemakers, N=179	-52.00 (-61.89 to -42.11) n=22	-31.52 (-35.16 to -27.87) n=157	-20.49 (-31.07 to -9.90)	0.0002
Week 52				
P-HEQ absenteeism among employed patients, N=239	6.80 (-2.71 to 16.32) n=51	14.03 (9.09 to 18.98) n=188	-7.23 (-17.96 to 3.51)	0.1860
P-HEQ presenteeism among employed patients, N=237	-46.29 (-51.49 to -41.10) n=50	-33.65 (-36.33 to -30.97) n=187	-12.64 (-18.50 to -6.78)	<0.0001
P-HEQ absenteeism among homemakers, N=154	2.94 (-12.62 to 18.49) n=23	12.97 (6.46 to 19.47) n=131	-10.03 (-26.90 to 6.84)	0.2420
P-HEQ presenteeism among homemakers, N=152	-51.12 (-60.83 to -41.41) n=22	-34.16 (-38.09 to -30.24) n=130	-16.96 (-27.49 to -6.42)	0.0018
Work-related outcomes in OPTIMA**				
	CDC achievers (A) N=58 Mean (95% CI)	Non-achievers (B) N=255 Mean (95% CI)	Difference (C)=(A)-(B) Mean (95% CI)	p Value
Week 26				
WPAI activity impairment, N=308	-28.89 (-34.12 to -23.67) n=58	-10.94 (-13.37 to -8.51) n=250	-17.96 (-23.81 to -12.10)	<0.0001
WPAI work impairment, N=131	-25.47 (-32.88 to -18.06) n=37	-7.77 (-12.30 to -3.24) n=94	-17.70 (-26.59 to -8.80)	0.0001
WPAI absenteeism, N=131	-8.20 (-14.42 to -1.99) n=37	-1.53 (-5.42 to 2.35) n=94	-6.67 (-14.02 to 0.68)	0.0750
WPAI presenteeism, N=144	-23.19 (-29.04 to -17.34) n=41	-9.41 (-12.99 to -5.82) n=103	-13.78 (-20.82 to -6.74)	0.0002
RA-WIS, N=153	-5.72 (-7.45 to -3.99) n=34	-2.09 (-3.00 to -1.18) n=119	-3.63 (-5.61 to -1.65)	0.0004

Continued

Table 2 Continued

Work-related outcomes in OPTIMA**

	CDC achievers (A) N=58 Mean (95% CI)	Non-achievers (B) N=255 Mean (95% CI)	Difference (C)=(A)–(B) Mean (95% CI)	p Value
Week 52				
WPAI activity impairment, N=284	–29.85 (–35.69 to –24.01) n=55	–10.60 (–13.37 to –7.84) n=229	–19.24 (–25.81 to –12.68)	<0.0001
WPAI work impairment, N=125	–26.09 (–32.86 to –19.33) n=38	–11.63 (–16.00 to –7.27) n=87	–14.46 (–22.72 to –6.20)	0.0007
WPAI absenteeism, N=125	–8.93 (–14.10 to –3.76) n=38	–4.56 (–7.96 to –1.15) n=87	–4.37 (–10.59 to 1.84)	0.1659
WPAI presenteeism, N=134	–24.31 (–30.07 to –18.56) n=39	–9.81 (–13.39 to –6.23) n=95	–14.50 (–21.48 to –7.53)	0.0001
RA-WIS, N=146	–6.38 (–8.40 to –4.36) n=33	–1.74 (–2.81 to –0.67) n=113	–4.64 (–6.96 to –2.32)	0.0001

Boldface indicates statistical significance at the 0.05 level.

*The adjusted analysis compared the mean change from baseline for each outcome by CDC achievement using an analysis of covariance model in which the impact of CDC on change in the outcome was estimated adjusting for the baseline value of the outcome.

†For VAS-Pain, negative change from baseline indicates an improvement in the outcome. For FACIT-Fatigue, SF-36 Mental Health and SF-36 Physical Health, positive change from baseline indicates an improvement in the outcome.

‡Indicates difference exceeds the respective MCID for the outcome.

§SF-36 Physical and Mental Health Component Scores were not measured in OPTIMA.

¶For P-HEQ absenteeism, the mean cumulative number of work days missed from baseline to week 26 and from baseline to week 52 was compared by CDC achievement. For P-HEQ presenteeism, the mean change from baseline to week 26 and from baseline to week 52 was compared by CDC achievement adjusting for the baseline P-HEQ presenteeism.

**For WPAI measures and RA-WIS, the mean change from baseline to week 26 and from baseline to week 52 was compared by CDC achievement adjusting for the baseline value.

CDC, comprehensive disease control; FACIT, Functional Assessment of Chronic Illness Therapy; MCID, minimal clinically important difference; P-HEQ, Patient Health Economic Questionnaire; RA-WIS, Rheumatoid Arthritis-Work Instability Scale; SF-36, Short Form 36; VAS, Visual Analogue Scale; WPAI, Work Productivity and Activity Impairment.

outcome. To assess differences between CDC achievement and ACR-EULAR remission achievement, the differences in SF-36 PCS, MCS, FACIT-F and VAS-Pain were compared between patients achieving CDC compared with those achieving ACR-EULAR remission (Boolean-based definition⁴) but not CDC adjusting for the baseline value of each outcome.

RESULTS

A total of 1467 patients were included in the pooled trial sample (493 (33.6%) from DE019, 661 (45.1%) from PREMIER and 313 (21.3%) from OPTIMA) of whom 200 (13.6%) achieved CDC. Overall, patients who achieved CDC at week 26 were younger, had lower mean DAS28(CRP), HAQ-DI, mTSS and VAS-Pain, and higher mean FACIT-F, SF-36 PCS and SF-36 MCS scores (table 1).

Pooled RA population

After adjusting for the baseline value of the outcome, CDC at week 26 was associated with the following improvements in PROs at week 26: a 20.0 (95% CI 16.9 to 23.2) point improvement in change from baseline in VAS-Pain; a 5.8 (4.4 to 7.1) point improvement in FACIT-F; a 10.8 (9.1 to 12.4) point improvement in SF-36 PCS and a 3.1 (1.4 to 4.8) point improvement in SF-36 MCS. Further, after adjusting for the baseline value of the outcome, CDC at week 26 was associated with the following improvements in PROs at week 52: a 16.9 (13.5–20.2) point improvement in VAS-Pain; a 4.9 (3.5–6.4) point improvement in FACIT-F; a 9.3 (7.6–11.1) point improvement in SF-36 PCS and a 2.2 (0.4–3.9) point improvement in SF-36 MCS (table 2). All improvements were statistically significant. In addition, the differences in SF-36 PCS, VAS-Pain and FACIT-F exceeded their respective MCIDs at both week 26 and 52. The difference in SF-36 MCS only exceeded the MCID at week 26.

Early RA population

Among patients with early RA enrolled in PREMIER, CDC achievement was associated with statistically significant improvements in P-HEQ presenteeism at weeks 26 and 52 (table 2). Achievement of CDC at week 26 was also associated with statistically significant and clinically meaningful differences in SF-36 PCS, VAS-Pain and FACIT-F at both week 26 and 52 and statistically significant and clinically meaningful differences in SF-36 MCS at week 26 (figures 1 and 2).

Early RA MTX-failure population

Among all patients with early RA enrolled in OPTIMA who failed to achieve LDA with MTX, achievement of CDC at week 26 was associated with statistically significant improvements in change in WPAI activity impairment, WPAI work impairment, WPAI presenteeism and RA-WIS at weeks 26 and 52 after adjusting for baseline values (table 2). Among all MTX-failure patients, achievement of CDC at week 26 was associated with statistically significant differences in VAS-Pain and FACIT-F at weeks 26 and 52 after adjusting for baseline values. The differences in VAS-Pain exceeded the MCID at both week 26 and 52 (figures 1 and 2).

Established RA population

Among patients with established RA enrolled in DE019, achievement of CDC at week 26 was associated with statistically significant improvements in change in SF-36 PCS, FACIT-F and VAS-Pain at weeks 26 and 52 after adjusting for the baseline scores (figures 1 and 2).

Incremental value of CDC

Compared with patients achieving DAS remission alone, CDC achievement was associated with statistically significant and clinically meaningful differences of 7.0 (4.0–10.0) and 6.9 (3.5–10.4) in SF-36 PCS at weeks 26 and 52, respectively. CDC




Figure 1 Comparison of patient-reported outcomes by CDC achievement and trial population at week 26.¹ FACIT, Functional Assessment of Chronic Illness Therapy; SF-36, Short Form 36; VAS, Visual Analogue Scale; RA, rheumatoid arthritis; MCS, mental component score; PCS, physical component score; CDC, comprehensive disease control; MTX, methotrexate.

achievement was also associated with a 9.0 (1.6–16.3) point decrease in VAS-Pain compared with DAS remission alone. When compared with patients achieving normal physical function alone, CDC achievement was associated with statistically significant differences of 4.0 (2.0–6.0) and 2.9 (0.6–5.2) in SF-36 PCS and 7.6 (3.1–12.1) and 5.5 (0.5–10.6) reductions in VAS-Pain at weeks 26 and 52, respectively. CDC achievement was associated with statistically significant and clinically meaningful improvements in SF-36 PCS, MCS, FACIT-F and VAS-Pain at both week 26 and 52 compared with patients with the absence of radiographic progression alone (table 3).

Comparison with ACR-EULAR remission

There were no statistically significant differences in VAS-Pain, FACIT-Fatigue, SF-36 PCS and SF-36 MCS between patients achieving CDC and patients achieving ACR-EULAR remission but not CDC at weeks 26 and 52 (table 4).

DISCUSSION

Clinical, functional and structural efficacy outcomes are the three aspects against which therapies are evaluated in the disease management for RA. In addition, simultaneous achievement of all three is proposed as the long-term treatment




Figure 2 Comparison of patient-reported outcomes by CDC achievement and trial population at week 52.¹ FACIT, Functional Assessment of Chronic Illness Therapy; SF-36, Short Form 36; VAS, Visual Analogue Scale; RA, rheumatoid arthritis; MCS, mental component score; PCS, physical component score; CDC, comprehensive disease control; MTX, methotrexate.

Table 3 Change in patient-reported outcomes from baseline to week 26 and week 52 by week 26 CDC components achieved (DE019, PREMIER and OPTIMA)

Continuous outcomes*†	CDC achievers (A) N=200 Mean (95% CI)	Achieved DAS remission (B) N=380 Mean (95% CI)	Achieved normal physician function (C) N=569 Mean (95% CI)	Achieved non-radiographic progression (D) N=1000 Mean (95% CI)	Difference (E)=(A)–(B) Mean (95% CI)	Difference (F)=(A)–(C) Mean (95% CI)	Difference (G)=(A)–(D) Mean (95% CI)
Week 26							
VAS-Pain change, N=1463	–47.49 (–50.07 to –44.91) 200	–42.21 (–48.53 to –35.90) 378	–39.89 (–43.62 to –36.16) 568	–18.32 (–19.95 to –16.68) 997	–5.27 (–12.09 to 1.55)	–7.60 (–12.11 to –3.08)	–29.17 (–32.24 to –26.11)
FACIT-Fatigue change, N=1452	13.65 (12.49 to 14.81) 196	10.81 (8.02 to 13.60) 373	13.92 (12.27 to 15.56) 563	4.42 (3.70 to 5.15) 988	2.84 (–0.18 to 5.86)	–0.27 (–2.27 to 1.74)	9.23 (7.84 to 10.61)
SF-36 mental change, N=1086	8.08 (6.50 to 9.67) 131	8.96 (5.55 to 12.38) 246	7.97 (5.94 to 9.99) 420	3.54 (2.56 to 4.52) 680	–0.88 (–4.64 to 2.89)	0.12 (–2.45 to 2.68)	4.54 (2.68 to 6.41)
SF-36 physical change, N=1086	20.36 (19.09 to 21.62) 131	13.39 (10.70 to 16.08) 246	16.33 (14.73 to 17.93) 420	4.81 (4.04 to 5.59) 680	6.97 (3.99 to 9.95)	4.03 (2.01 to 6.05)	15.55 (14.05 to 17.04)
Week 52							
VAS-Pain change, N=1306	–46.11 (–48.95 to –43.27) 189	–37.16 (–43.92 to –30.40) 350	–40.61 (–44.80 to –36.42) 516	–21.02 (–22.87 to –19.17) 900	–8.95 (–16.28 to –1.62)	–5.50 (–10.55 to –0.45)	–25.09 (–28.49 to –21.68)
FACIT-Fatigue change, N=1326	13.27 (12.01 to 14.52) 186	10.35 (7.43 to 13.27) 348	13.61 (11.82 to 15.40) 521	5.42 (4.63 to 6.22) 912	2.92 (–0.26 to 6.09)	–0.35 (–2.52 to 1.83)	7.84 (6.34 to 9.34)
SF-36 mental change, N=1001	7.28 (5.64 to 8.92) 126	8.52 (4.91 to 12.13) 236	6.93 (4.80 to 9.06) 393	3.95 (2.92 to 4.99) 630	–1.24 (–5.21 to 2.73)	0.35 (–2.34 to 3.04)	3.33 (1.38 to 5.27)
SF-36 physical change, N=1001	20.01 (18.57 to 21.45) 126	13.07 (9.96 to 16.18) 236	17.12 (15.29 to 18.96) 393	6.42 (5.52 to 7.31) 630	6.94 (3.51 to 10.37)	2.89 (0.57 to 5.20)	13.59 (11.88 to 15.30)

Boldface indicates statistical significance at the 0.05 level.

*The adjusted analysis compared the mean change from baseline for each outcome by CDC component achievement using an analysis of covariance model in which the impact of component achievement on change in the outcome was estimated adjusting for the baseline value of the outcome.

†For VAS-Pain, negative change from baseline indicates an improvement in the outcome. For FACIT-Fatigue, SF-36 Mental Health and SF-36 Physical Health, positive change from baseline indicates an improvement in the outcome. CDC, comprehensive disease control; DAS, Disease Activity Score; FACIT, Functional Assessment of Chronic Illness Therapy; SF-36, Short Form 36; VAS, Visual Analogue Scale.

Table 4 Change in patient-reported outcomes from baseline to week 26 and week 52 by week 26 CDC/ACR-EULAR remission achievement (DE019, PREMIER and OPTIMA)

Continuous outcomes*†	CDC achievers (A)	ACR-EULAR remission but not CDC (B)	Neither ACR-EULAR remission nor CDC (C)	Difference (D)=(A)-(B)	
	N=200 Mean (95% CI)	N=29 Mean (95% CI)	N=1226 Mean (95% CI)	Mean (95% CI)	p Value
Week 26					
VAS-Pain change, N=1451	-47.07 (-49.96 to -44.19) 200	-51.59 (-59.12 to -44.05) 29	-26.45 (-27.61 to -25.29) 1222	4.51 (-3.55 to 12.57)	0.2722
FACIT-Fatigue change, N=1440	13.31 (12.06 to 14.56) 196	12.82 (9.62 to 16.01) 29	7.42 (6.92 to 7.91) 1215	0.50 (-2.93 to 3.92)	0.7764
SF-36 mental change, N=1085	8.06 (6.46 to 9.67) 131	6.05 (2.52 to 9.57) 27	4.95 (4.35 to 5.55) 927	2.02 (-1.85 to 5.89)	0.3065
SF-36 physical change, N=1085	19.69 (18.19 to 21.20) 131	19.17 (15.91 to 22.44) 27	8.59 (8.03 to 9.15) 927	0.52 (-3.07 to 4.11)	0.7754
Week 52					
VAS-Pain change, N=1294	-45.84 (-48.87 to -42.80) 189	-51.22 (-59.36 to -43.09) 26	-28.52 (-29.79 to -27.26) 1079	5.39 (-3.29 to 14.07)	0.2234
FACIT-Fatigue change, N=1314	13.00 (11.69 to 14.31) 186	14.16 (10.64 to 17.68) 25	7.97 (7.44 to 8.50) 1103	-1.16 (-4.90 to 2.59)	0.5441
SF-36 mental change, N=1000	7.27 (5.62 to 8.93) 126	8.19 (4.48 to 11.89) 25	5.03 (4.39 to 5.66) 849	-0.91 (-4.97 to 3.14)	0.6580
SF-36 physical change, N=1000	19.38 (17.77 to 21.00) 126	20.54 (16.97 to 24.11) 25	9.72 (9.11 to 10.34) 849	-1.16 (-5.08 to 2.76)	0.5626

*The adjusted analysis compared the mean change from baseline for each outcome by CDC/ACR-EULAR achievement using an analysis of covariance model in which the impact of achievement on change in the outcome was estimated adjusting for the baseline value of the outcome.

†For VAS-Pain, negative change from baseline indicates an improvement in the outcome. For FACIT-Fatigue, SF-36 Mental Health and SF-36 Physical Health, positive change from baseline indicates an improvement in the outcome.

ACR-EULAR, American College of Rheumatology-European League Against Rheumatism; CDC, comprehensive disease control; FACIT, Functional Assessment of Chronic Illness Therapy; SF-36, Short Form 36; VAS, Visual Analogue Scale.

goal with implied benefits on patients. While correlated, these three aspects have respective variations and each of them adds value while being achieved simultaneously. The purpose of this study is to assess the impact of such achievement, referred to as CDC, on short-term as well as long-term patient outcomes including fatigue, pain, work outcomes and general physical and mental health. CDC includes DAS remission of <2.6, normal physical function (as measured by HAQ-DI) and absence of radiographic progression (as measured by mTSS).

In the present study, CDC was associated with significant improvements in patient's work-related outcomes, HRQoL, pain and fatigue. These improvements were observed both at the time of CDC achievement and 26 weeks later. The improvements in HRQoL are especially pronounced, as patients who achieve CDC are 2.5 times more likely to have SF-36 MCS score greater than the general population norm and 9.8 times more likely to have SF-36 PCS greater than the general population norm at week 26. Similarly, CDC achievers are 2.3 times more likely to have SF-36 MCS score greater than the general population norm and 6.7 times more likely to have SF-36 PCS greater than the general population norm at week 52. These improvements are associated with reduced healthcare use and costs and reduced mortality,^{38 39} further demonstrating that CDC achievement is associated with broad and significant improvements in patient outcomes. Also, the differences between the CDC achievers and non-achievers for all the measured outcomes were greater than the clinically meaningful thresholds for each of the outcome. The simultaneous achievement of all three CDC domains was also demonstrated to be associated with significant improvements in physical function

and pain compared with achieving the individual domains in isolation. This result suggests there may be an incremental benefit related to achieving all three components. An adjusted correlation analysis of the individual components showed that nearly 50%–55% of the variations in the outcomes at week 26 (SF-36 and VAS-Pain) were explained by improvements in DAS28 remission, 35%–47% by improvements in HAQ-DI score and 2%–9% by mTSS scores.

Current recommendations state that treatment of RA should focus on achieving clinical remission to inhibit progression of joint damage and improve physical function and quality of life.³ The ACR and EULAR also recognise the importance of remission for predicting improvement in physical function and halting radiographic progression. It is noted in the EULAR 2013 recommendation update that ACR-EULAR remission should serve as the optimal treatment target to be used in routine clinical practice, while LDA is a good alternative for the many patients who cannot attain remission. Within the pooled population in this study, a total of 94 patients experienced ACR-EULAR remission, of whom 65 (69.1%) also achieved CDC. The presence of radiographic progression was the most common reason for discordance (17 patients). There were no significant differences in SF-36 PCS, MCS, FACIT-F or VAS-Pain between patients achieving CDC and those achieving ACR-EULAR remission (but not CDC), although this analysis was limited by small sample size (table 4). Regardless of which treatment target is selected, the long-term goal remains to maximise long-term HRQoL through control of symptoms, prevention of structural damage, normalisation of function and social participation. Discussions around the simultaneous achievement of clinical, physical and structural efficacy can also engage

patients and encourage patients to involve their observation around physical function.

This analysis has several strengths. First, we pooled data from three large RCTs of early RA, established RA and MTX non-responders and found that within trial, analyses were consistent with the results from the pooled data. So while patients with experienced disease may achieve CDC at a lower rate, there is no differential impact of CDC achievement on quality of life by RA duration. Further, improvements in PROs observed at 52 weeks were similar to those observed at 26 weeks, demonstrating that achievement of CDC may have a long-standing effect on quality of life.

This analysis has limitations. First, patient characteristics were not balanced between CDC achievers and non-achievers. However, adjustment for these factors may not account for all baseline differences between CDC achievers and non-achievers. An observed case analysis was performed, which led to 17% of patients being dropped from the analysis. However, imputation procedures require assumptions about the underlying missing data mechanism and may not be appropriate for this analysis. Last, not all outcomes, such as SF-36, were measured in all trials making it difficult to assess the full impact of CDC achievement on SF-36 in patients who were non-responders to MTX.

This post hoc analysis of three individual clinical trials demonstrated the benefits of CDC at the time of achievement and 6 months postachievement. This finding can be strengthened if supported by the results of a future prospective randomised controlled trial using CDC as a ranked secondary endpoint. Future research into whether the benefits of CDC are maintained over a longer-term is also warranted.

CDC achievement was associated with clinically meaningful improvements in work-related outcomes, HRQoL, pain and fatigue compared with those who did not achieve CDC. These results demonstrate that simultaneous improvement in all three RA domains can provide meaningful benefits to patients.

Contributors PE, AK and PM conceptualised the research and contributed to the interpretation of the data. YB and AG wrote the statistical analysis plan and analysed the data. AG, PE, AK, PM and YB contributed to the drafting of the paper and approved the final version. Significant contribution towards medical writing and analytical support was provided by Keith Betts, PhD, of Analysis Group; this support was funded by AbbVie. AbbVie sponsored the studies, participated in study design, data collection, analysis and interpretation, and in the writing, reviewing and approval of the final version.

Funding AbbVie funded the studies DE019 (NCT00195702), PREMIER (NCT00195663) and OPTIMA (NCT00420927), contributed to their design and was involved in the collection, analysis and interpretation of the data, and in the writing, review and approval of the manuscript.

Competing interests PE—Grant/research support: AbbVie, Pfizer, Novartis, BMS and Roche; Consultant for: AbbVie, Pfizer, MSD, UCB, Roche, Novartis and BMS. AK—Grant/research support: AbbVie, Amgen, Astra-Zeneca, BMS, Celgene, Centocor-Janssen, Pfizer, Roche and UCB; Consultant for: AbbVie, Amgen, Astra-Zeneca, BMS, Celgene, Centocor-Janssen, Pfizer, Roche and UCB. AG—Shareholder of: AbbVie; Employee of: AbbVie. YB—Shareholder of: AbbVie; Employee of: AbbVie. PM—Shareholder of: AbbVie; Employee of: AbbVie.

Provenance and peer review Not commissioned; externally peer reviewed.

Open Access This is an Open Access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 3.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/3.0/>

REFERENCES

1 Lindqvist E, Jonsson K, Saxne T, *et al.* Course of radiographic damage over 10 years in a cohort with early rheumatoid arthritis. *Ann Rheum Dis* 2003;62:611–16.

- 2 van der Heijde DM, van Leeuwen MA, van Riel PL, *et al.* Radiographic progression on radiographs of hands and feet during the first 3 years of rheumatoid arthritis measured according to Sharp's method (van der Heijde modification). *J Rheumatol* 1995;22:1792–6.
- 3 Smolen JS, Aletaha D, Bijlsma JW, *et al.* Treating rheumatoid arthritis to target: recommendations of an international task force. *Ann Rheum Dis* 2010;69:631–7.
- 4 Felson DT, Smolen JS, Wells G, *et al.* American College of Rheumatology/European League against Rheumatism provisional definition of remission in rheumatoid arthritis for clinical trials. *Ann Rheum Dis* 2011;70:404–13.
- 5 Smolen JS, Landewe R, Breedveld FC, *et al.* EULAR recommendations for the management of rheumatoid arthritis with synthetic and biological disease-modifying antirheumatic drugs: 2013 update. *Ann Rheum Dis* 2014;73:492–509.
- 6 Prevoo ML, van 't Hof MA, Kuper HH, *et al.* Modified disease activity scores that include twenty-eight-joint counts. Development and validation in a prospective longitudinal study of patients with rheumatoid arthritis. *Arthritis Rheum* 1995;38:44–8.
- 7 Mittendorf T, Dietz B, Sterz R, *et al.* Personal and economic burden of late-stage rheumatoid arthritis among patients treated with adalimumab: an evaluation from a patient's perspective. *Rheumatology (Oxford)* 2008;47:188–93.
- 8 Pollard LC, Choy EH, Gonzalez J, *et al.* Fatigue in rheumatoid arthritis reflects pain, not disease activity. *Rheumatology (Oxford)* 2006;45:885–9.
- 9 Cohen SB, Strand V, Aguilar D, *et al.* Patient- versus physician-reported outcomes in rheumatoid arthritis patients treated with recombinant interleukin-1 receptor antagonist (anakinra) therapy. *Rheumatology (Oxford)* 2004;43:704–11.
- 10 Strand V, Cohen S, Crawford B, *et al.* Patient-reported outcomes better discriminate active treatment from placebo in randomized controlled trials in rheumatoid arthritis. *Rheumatology (Oxford)* 2004;43:640–7.
- 11 Hazes JM, Geuskens GA, Burdorf A. Work limitations in the outcome assessment of rheumatoid arthritis. *J Rheumatol* 2005;32:980–2.
- 12 Brouwer WB, Meerding WJ, Lamers LM, *et al.* The relationship between productivity and health-related QOL: an exploration. *Pharmacoeconomics* 2005;23:209–18.
- 13 *The Burden of Musculoskeletal Conditions at the Start of the New Millennium.* World Health Organization, 2003. http://whqlibdoc.who.int/trs/WHO_TRS_919.pdf (accessed 13 Dec 2013).
- 14 Carr A, Hewlett S, Hughes R, *et al.* Rheumatology outcomes: the patient's perspective. *J Rheumatol* 2003;30:880–3.
- 15 Kirwan JR, Minnock P, Adebajo A, *et al.* Patient perspective: fatigue as a recommended patient centered outcome measure in rheumatoid arthritis. *J Rheumatol* 2007;34:1174–7.
- 16 Heiberg T, Kvien TK. Preferences for improved health examined in 1,024 patients with rheumatoid arthritis: pain has highest priority. *Arthritis Rheum* 2002;47:391–7.
- 17 Breedveld FC, Weisman MH, Kavanaugh AF, *et al.* The PREMIER study: a multicenter, randomized, double-blind clinical trial of combination therapy with adalimumab plus methotrexate versus methotrexate alone or adalimumab alone in patients with early, aggressive rheumatoid arthritis who had not had previous methotrexate treatment. *Arthritis Rheum* 2006;54:26–37.
- 18 Kavanaugh A, Fleischmann RM, Emery P, *et al.* Clinical, functional and radiographic consequences of achieving stable low disease activity and remission with adalimumab plus methotrexate or methotrexate alone in early rheumatoid arthritis: 26-week results from the randomized, controlled OPTIMA study. *Ann Rheum Dis* 2013;72:64–71.
- 19 Keystone EC, Kavanaugh AF, Sharp JT, *et al.* Radiographic, clinical, and functional outcomes of treatment with adalimumab (a human anti-tumor necrosis factor monoclonal antibody) in patients with active rheumatoid arthritis receiving concomitant methotrexate therapy: a randomized, placebo-controlled, 52-week trial. *Arthritis Rheum* 2004;50:1400–11.
- 20 Smolen J, Fleischmann R, Emery P, *et al.* The OPTIMA Study of Methotrexate and Adalimumab: 78-Week Outcomes in Early Rheumatoid Arthritis Patients Based on Achieving a Low DAS28 Target After 26 Weeks. *Ann Rheum Dis* 2011;70:259.
- 21 Franssen J, van Riel PL. The Disease Activity Score and the EULAR response criteria. *Clin Exp Rheumatol* 2005;23:593–9.
- 22 Fries JF, Spitz PW, Young DY. The dimensions of health outcomes: the health assessment questionnaire, disability and pain scales. *J Rheumatol* 1982;9:789–93.
- 23 Hochberg MC, Chang RW, Dwosh I, *et al.* The American College of Rheumatology 1991 revised criteria for the classification of global functional status in rheumatoid arthritis. *Arthritis Rheum* 1992;35:498–502.
- 24 Plant MJ, Saklatvala J, Borg AA, *et al.* Measurement and prediction of radiological progression in early rheumatoid arthritis. *J Rheumatol* 1994;21:1808–13.
- 25 Sharp JT, Lidsky MD, Collins LC, *et al.* Methods of scoring the progression of radiologic changes in rheumatoid arthritis. Correlation of radiologic, clinical and laboratory abnormalities. *Arthritis Rheum* 1971;14:706–20.
- 26 van der Heijde D. How to read radiographs according to the Sharp/van der Heijde method. *J Rheumatol* 1999;26:743–5.
- 27 van der Heijde D, Simon L, Smolen J, *et al.* How to report radiographic data in randomized clinical trials in rheumatoid arthritis: guidelines from a roundtable discussion. *Arthritis Rheum* 2002;47:215–18.

- 28 van Vollenhoven RF, Cifaldi MA, Ray S, *et al.* Improvement in work place and household productivity for patients with early rheumatoid arthritis treated with adalimumab plus methotrexate: work outcomes and their correlations with clinical and radiographic measures from a randomized controlled trial companion study. *Arthritis Care Res (Hoboken)* 2010;62:226–34.
- 29 Reilly MC, Zbrozek AS, Dukes EM. The validity and reproducibility of a work productivity and activity impairment instrument. *Pharmacoeconomics* 1993;4:353–65.
- 30 Gilworth G, Chamberlain MA, Harvey A, *et al.* Development of a work instability scale for rheumatoid arthritis. *Arthritis Rheum* 2003;49:349–54.
- 31 Ware JE Jr, Sherbourne CD. The MOS 36-item short-form health survey (SF-36). I. Conceptual framework and item selection. *Med Care* 1992;30:473–83.
- 32 Kosinski M, Zhao SZ, Dedhiya S Jr, *et al.* Determining minimally important changes in generic and disease-specific health-related quality of life questionnaires in clinical trials of rheumatoid arthritis. *Arthritis Rheum* 2000;43:1478–87.
- 33 Brazier JE, Harper R, Jones NM, *et al.* Validating the SF-36 health survey questionnaire: new outcome measure for primary care. *BMJ* 1992;305:160–4.
- 34 Ware JE, Kosinski M, Keller SD. *SF-36 physical and mental health summary scales: a user's manual*. Boston: The Health Institute, New England Medical Center, 1994.
- 35 Lubeck DP. Patient-reported outcomes and their role in the assessment of rheumatoid arthritis. *Pharmacoeconomics* 2004;22:27–38.
- 36 Wells GA, Tugwell P, Kraag GR, *et al.* Minimum important difference between patients with rheumatoid arthritis: the patient's perspective. *J Rheumatol* 1993;20:557–60.
- 37 Yellen SB, Cella DF, Webster K, *et al.* Measuring fatigue and other anemia-related symptoms with the Functional Assessment of Cancer Therapy (FACT) measurement system. *J Pain Symptom Manage* 1997;13:63–74.
- 38 Ethgen O, Kahler KH, Kong SX, *et al.* The effect of health related quality of life on reported use of health care resources in patients with osteoarthritis and rheumatoid arthritis: a longitudinal analysis. *J Rheumatol* 2002;29:1147–55.
- 39 Singh JA, Nelson DB, Fink HA, *et al.* Health-related quality of life predicts future health care utilization and mortality in veterans with self-reported physician-diagnosed arthritis: the veterans arthritis quality of life study. *Semin Arthritis Rheum* 2005;34:755–65.