Separately tackling the development of erosions with denosumab: ultimately closing a gap in the treatment of patients with rheumatoid arthritis or trying too hard too late?

P Verschueren,^{1,2} R Westhovens^{1,2}

Patients with rheumatoid arthritis (RA) have benefited from an unprecedented treatment improvement over the last two decades. Taking the disease seriously, with earlier and more intensive treat-to-target strategies and new treatment options with biologicals, has led to a better long-term disease control. Classical antirheumatic therapies, primarily directed against the inflammatory/immunological processes that determine RA disease activity, are successfully improving pain and the severe clinical and functional burden patients are confronted with. Moreover, one of the most important previous hallmarks of this disease, severe joint destruction with subsequent deformations, can now in most cases be prevented thanks to the development of new treatment strategies and the introduction of novel therapeutics. The Cobra trial¹ demonstrated the existence of a window of opportunity for preventing joint damage in early RA and the Attract trial² showed an arrest in further X-ray damage—at least at the group level—in severe methotrexate (MTX) refractory disease treated with infliximab, which was unprecedented at that time. Thanks to more effective treatments prevention of further damage was reported and even, in a (small) number of cases, what could be considered as healing of erosions.^{3–4} Other publications focused on a potential uncoupling between controlling inflammation and damage. Prevention of X-ray progression appeared possible independent from successfully reducing disease activity, at least with certain therapeutics.⁵ Despite the fact that today joint damage can be prevented in the large majority of cases, it is still an important determinant in the

evaluation of RA treatment efficacy given the link between X-ray progression and future physical disability.⁶

Takeuchi *et al*⁷ demonstrate an additional inhibitory effect on the progression of bone erosions (not joint space narrowing) of three different dosages of denosumab on top of the traditional RA treatment over a 12-month period in Japanese patients with RA, confirming results reported already 8 years ago by Cohen *et al*⁸ in North America. Although the erosive changes are relatively small and a certain ‘hang-over’ of MTX is possible, this study has clear strengths like the placebo-controlled design, the relatively early study population, the similar evolution of disease activity markers across treatment arms and the documentation of improved bone-turnover and not cartilage-turnover markers, confirming the X-ray results. The results are convincing even though the study population consisted of patients with relatively low disease activity. One wonders what this would have been in more active patients or with less glucocorticoid use. On the other hand, the treat-to-target approach was not applied, making the relevance of the study results for daily practice somewhat uncertain. Nevertheless, this study confirms the predefined proof of concept that denosumab can be of help in preventing joint erosions in RA.

Earlier attempts to prevent (cartilage) damage in RA with therapeutic agents not specifically targeting joint inflammation were already initiated >15 years ago, focusing mainly on metalloproteinase inhibitors. A nice overview on this topic was written by Close⁹ in this journal in 2001. Questions about representativeness of the animal models the drugs were tested in, lack of sensitivity in outcome measurements, and specificity and safety issues of the drugs tested, were raised. Inefficacy of minocycline—known also for its collagenase inhibitory properties—in preventing X-ray damage in RA¹⁰ and the preliminary discontinuation of the

Trocade trial with a MMP-1 inhibitor despite promising explorative findings,¹¹ precluded further research along this therapeutic pathway in RA because of an apparent unfavourable risk–benefit profile. Currently, matrix metalloproteinase-3 is still under investigation although not so much as a direct therapeutic target but rather as a predictive biomarker for joint damage and therefore as an additional measurement instrument in treat-to-target strategies.^{12–13} In the meantime, attempts to separately tackle bony erosions were continued at a smaller scale and with mixed success, with denosumab and with bisphosphonates¹⁴ and parathormone (PTH),¹⁵ but apparently without real breakthrough and without any further implementation in daily practice.

While the idea of additional prevention of cartilage and/or bone destruction with agents not directly targeting inflammation is being investigated already for more than two decades and is again brought to our attention with this denosumab trial, one could question the relevance of this complementary approach given the successfulness of current RA treatment standards.

There is no doubt that avoiding the occurrence of joint erosions is still an important issue. Joint erosions historically have always been perceived to be a critical indicator of permanent future disability in patients with RA, although data suggest also that joint space narrowing, occurring early in disease process, may be a more important determinant of irreversible physical disability.^{6–16} On the other hand, erosions are more prevalent than joint space narrowing in early RA and a recent study suggested existing erosions in a single joint are associated with future joint space narrowing making it clear that tackling both bone and cartilage damage is important, especially in early disease. Interestingly, these data derived from the PREMIER study¹⁷ also suggested that the effect of pre-existing erosions on future joint space narrowing, documented in the MTX+adalimumab (ADA) and the ADA monotherapy group, was at least partly independent from the presence of clinical synovitis.

Thanks to the availability of effective therapeutic strategies we are more successful in controlling disease activity than ever before, but we should not forget to evaluate joint damage as an objective and cumulative readout of treatment effectiveness. Swollen joint counts and all disease activity measurement instruments including swollen joints, as well as acute phase reactants, are still robustly associated with radiographic damage.¹⁸ This highlights

¹Department of Development and Regeneration KU Leuven, Skeletal Biology and Engineering Research Center, Leuven, Belgium; ²Department of Rheumatology, University Hospitals Leuven, Leuven, Belgium

Correspondence to Dr R Westhovens, Department of Rheumatology, UZ KU Leuven, Herestraat 49, Leuven 3000, Belgium; rene.westhovens@uzleuven.be

the importance of the ongoing discussion about the definition of 'real remission' and how to evaluate this in daily practice¹⁹ as well as further research on how to implement effective intensive treatment strategies as quickly as possible, especially in early disease.²⁰

Perhaps not surprisingly, in trials with the more recently developed tumour necrosis factor blockers like golimumab it appeared more difficult to demonstrate a significant radiographic benefit from the study drug because X-ray progression in the so-called MTX-refractory study population was globally lower than expected.²¹ The latter findings led to an important viewpoint paper by Landewé *et al*²² focusing on trial design and statistical challenges when evaluating patients with RA and measuring treatment efficacy, especially efficacy based on X-ray outcomes. Dr Landewé correctly states that we can no longer hang on to the classical enrichment techniques selecting very active patients with a severe risk profile for inclusion in trials, with the aim of demonstrating superiority in damage control of one treatment over another, without serious ethical problems in an era where many effective alternative therapies are available. He proposes to replace 'inhibition of radiographic progression' by 'maintaining structural integrity' as an outcome parameter in clinical trials. While the focus of this paper was on methodological issues as a consequence of the limited radiographic progression in clinical trials, this seems perfectly in accordance with our own experiences in daily practice, where early intensive step down and treat-to-target strategies are capable of bringing RA disease activity under control in the vast majority of patients, making X-ray damage little by little a minor problem.²³ For the subset of patients refractory to optimal initial treatment, today several additional treatments are available to control disease activity and in this way avoid radiographic progression.

In view of preventing radiographic damage and future disability it is clear that aspects of timely treatment and avoiding treatment delay,²⁴ as well as personalised communication with individual patients about their fears and beliefs, and issues related to the implementation of early intensive treatments in daily practice^{25, 26} are at least as important as the introduction of new and sophisticated medication. Some of us might still tend to adapt their treatment choices based on markers of a bad prognosis in terms of future X-ray damage. Unfortunately

however, current prediction matrices combining several of these markers in order to improve their predictive capacity still do not perform adequately when used in daily practice²⁷ probably as a consequence of the growing application, although flexibly, of the treat-to-target principle. From our point of view, pending the discovery of more adequate prognostic biomarkers allowing personalised treatment stratification, all patients with RA should be treated intensively and any symptoms and signs pointing to remaining disease activity should be taken seriously, irrespective of the patient's traditional risk profile. We agree with the mantra of 'maintaining structural integrity' also in the present treatment era, but would like to expand this towards 'for as many patients as possible' and hopefully allowing 'maintenance of optimal societal participation'.

A last and very important issue to consider is the eventual additional cost for the introduction of denosumab as an adjuvant therapy for RA in an era where societal expenditures for biologicals are already very high and even utopic in many countries of our world. It is questionable if cost-effectivity would ever be reached taking into consideration the treatment goals already achieved using the present armamentarium. Probably, for most patients a more adequate implementation of the current knowledge on how to induce and maintain disease control would be more cost-effective than trying to tackle additionally the progression of erosions with denosumab. Of course besides improving inflammation and preventing joint damage, adequate RA management involves also the prevention and treatment of associated co-morbidities and safety issues that might occur with any additional therapy. The proposed additional denosumab treatment could, apart from preventing joint erosions, in certain patients also constitute a therapy for RA-associated osteoporosis, as suggested in the current paper⁷ reporting also increased bone density. One could therefore consider to reserve such therapy for those patients with RA with a particularly high fracture risk, in order to improve the cost-effectiveness.

Competing interests RW: consultant for Janssen, speakers bureau for BMS, research money Roche and principal investigator Galapagos. All money to his institution. PV holds the Pfizer Early RA Chair at the KU Leuven.

Provenance and peer review Commissioned; externally peer reviewed.

To cite Verschueren P, Westhovens R. *Ann Rheum Dis* 2016;**75**:947–949.

Received 4 January 2016

Accepted 17 January 2016

Published Online First 9 February 2016

► <http://dx.doi.org/10.1136/annrheumdis-2015-208052>

Ann Rheum Dis 2016;**75**:947–949.

doi:10.1136/annrheumdis-2015-208863

REFERENCES

- Boers M, Verhoeven AC, Markusse HM, *et al*. Randomised comparison of combined step-down prednisolone, methotrexate and sulphasalazine with sulphasalazine alone in early rheumatoid arthritis. *Lancet* 1997;350:309–18.
- Lipsky PE, van der Heijde DM, St Clair EW, *et al*. Infliximab and methotrexate in the treatment of rheumatoid arthritis. Anti-Tumor Necrosis Factor Trial in Rheumatoid Arthritis with Concomitant Therapy Study Group. *N Engl J Med* 2000;343:1594–602.
- Sharp JT, Van Der Heijde D, Boers M, *et al*. Repair of erosions in rheumatoid arthritis does occur. Results from 2 studies by the OMERACT Subcommittee on Healing of Erosions. *J Rheumatol* 2003;30:1102–7.
- Møller Døhn U, Boonen A, Hetland ML, *et al*. Erosive progression is minimal, but erosion healing rare, in patients with rheumatoid arthritis treated with adalimumab. A 1 year investigator-initiated follow-up study using high-resolution computed tomography as the primary outcome measure. *Rheum Dis* 2009;68:1585–90.
- Landewé R, van der Heijde D, Klareskog L, *et al*. Disconnect between inflammation and joint destruction after treatment with etanercept plus methotrexate: results from the trial of etanercept and methotrexate with radiographic and patient outcomes. *Arthritis Rheum* 2006;54:3119–25.
- van der Heijde D, Landewé R, van Vollenhoven R, *et al*. Level of radiographic damage and radiographic progression are determinants of physical function: a longitudinal analysis of the TEMPO trial. *Ann Rheum Dis* 2008;67:1267–70.
- Takeuchi T, Tanaka Y, Ishiguro N, *et al*. Effect of denosumab on Japanese patients with rheumatoid arthritis: a dose-response study of AMG 162 (Denosumab) in patients with rheumatoid arthritis on methotrexate to validate inhibitory effect on bone erosion (DRIVE)-a twelve-month, multicenter, randomized, double-blind, placebo-controlled, phase II clinical trial. *Ann Rheum Dis* 2016;**75**: 983–90.
- Cohen SB, Dore RK, Lane NE, *et al*. Denosumab Rheumatoid Arthritis Study Group. Denosumab treatment effects on structural damage, bone mineral density, and bone turnover in rheumatoid arthritis: a twelve-month, multicenter, randomized, double-blind, placebo-controlled, phase II clinical trial. *Arthritis Rheum* 2008;58:1299–309.
- Close DR. Matrix metalloproteinase inhibitors in rheumatic diseases. *Ann Rheum Dis* 2001;60(Suppl 3):iii62–7.
- Bluhm GB, Sharp JT, Tilley B, *et al*. Radiographic results from the Minocycline in Rheumatoid Arthritis (MIRA) trial. *J Rheumatol* 1997;24:1295–302.
- Hemmings FJ, Farhan M, Rowland J, *et al*. Tolerability and pharmacokinetics of the collagenase-selective inhibitor Trocade™ in patients

- with rheumatoid arthritis. *Rheumatology (Oxford)* 2001;40:537–43.
- 12 Ma JD, Wei XN, Zheng DH. Continuously elevated serum matrix metalloproteinase-3 for 3 ~ 6 months predict one-year radiographic progression in rheumatoid arthritis: a prospective cohort study. *Arthritis Res Ther* 2015;17:289.
 - 13 Urata Y, Uesato R, Tanaka D, *et al.* Treating to target matrix metalloproteinase 3 normalisation together with disease activity score below 2.6 yields better effects than each alone in rheumatoid arthritis patients: T-4 Study. *Ann Rheum Dis* 2012;71:534–40.
 - 14 Jarrett SJ, Conaghan PG, Sloan VS, *et al.* Preliminary evidence for a structural benefit of the new bisphosphonate zoledronic acid in early rheumatoid arthritis. *Arthritis Rheum* 2006;54:1410–4.
 - 15 Migliore A, Massafra U, Bizzi E, *et al.* May etanercept and PTH (1–34) association heal erosions in early rheumatoid arthritis? A pilot study. *Eur Rev Med Pharmacol Sci* 2012;16:363–9.
 - 16 Aletaha D, Funovits J, Smolen JS. Physical disability in rheumatoid arthritis is associated with cartilage damage rather than bone destruction. *Ann Rheum Dis* 2011;70:733–9.
 - 17 Landewé R, Smolen JS, Florentinus S, *et al.* Existing joint erosions increase the risk of joint space narrowing independently of clinical synovitis in patients with early rheumatoid arthritis. *Arthritis Res Ther* 2015;17:133.
 - 18 Navarro-Compán V, Gherghe AM, Smolen JS, *et al.* Relationship between disease activity indices and their individual components and radiographic progression in RA: a systematic literature review. *Rheumatology (Oxford)* 2015;54:994–1007.
 - 19 Salomon-Escoto KI, Gravalles EM, Kay J. Assessment of control of rheumatoid arthritis disease activity. *Best Pract Res Clin Rheumatol* 2011;25:497–507.
 - 20 De Cock D, Van der Elst K, Meyfroidt S, *et al.* The optimal combination therapy for the treatment of early rheumatoid arthritis. *Expert Opin Pharmacother* 2015;16:1615–25.
 - 21 Rahman MU, Buchanan J, Doyle MK, *et al.* Changes in patient characteristics in anti-tumour necrosis factor clinical trials for rheumatoid arthritis: results of an analysis of the literature over the past 16 years. *Ann Rheum Dis* 2011;70:1631–40.
 - 22 Landewé R, Strand V, van der Heijde D. From inhibition of radiographic progression to maintaining structural integrity: a methodological framework for radiographic progression in rheumatoid arthritis and psoriatic arthritis clinical trials. *Ann Rheum Dis* 2013;72:1113–17.
 - 23 Verschueren P, De Cock D, Corluy L, *et al.* Remission induction with DMARD combination and glucocorticoids is not superior to remission induction with MTX monotherapy and glucocorticoids: 52 week results of the high risk group of the CareRA trial. *Ann Rheum Dis* 2015;74 (Suppl 2):139.
 - 24 Raza K, Stack R, Kumar K, *et al.* Delays in assessment of patients with rheumatoid arthritis: variations across Europe. *Ann Rheum Dis* 2011;70:1822–5.
 - 25 Van der Elst K, Meyfroidt S, De Cock D, *et al.* Unraveling patient-preferred health and treatment outcomes in early rheumatoid arthritis: A longitudinal qualitative study. *Arthritis Care Res (Hoboken)* 2015. doi:10.1002/acr.22824 [epub ahead of print 29 Dec 2015].
 - 26 Meyfroidt S, van Hulst L, De Cock D, *et al.* Factors influencing the prescription of intensive combination treatment strategies for early rheumatoid arthritis. *Scand J Rheumatol* 2014;43:265–72. doi:10.3109/03009742.2013.863382
 - 27 De Cock D, Vanderschueren G, Meyfroidt S, *et al.* The performance of matrices in daily clinical practice to predict rapid radiologic progression in patients with early RA. *Semin Arthritis Rheum* 2014;43:627–31.