



OPEN ACCESS

CONCISE REPORT

Sustained improvements in MRI outcomes with abatacept following the withdrawal of all treatments in patients with early, progressive rheumatoid arthritis

Charles Peterfy,¹ Gerd R Burmester,² Vivian P Bykerk,³ Bernard G Combe,⁴ Julie C DiCarlo,¹ Daniel E Furst,⁵ Tom W J Huizinga,⁶ Dennis A Wong,⁷ Philip G Conaghan,^{8,9} Paul Emery^{8,9}

Handling editor Tore K Kvien

► Additional material is published online only. To view please visit the journal online (<http://dx.doi.org/10.1136/annrheumdis-2015-208258>).

For numbered affiliations see end of article.

Correspondence to

Dr Charles Peterfy, Spire Sciences, Inc., 5314 Boca Marina Cir N, Boca Raton, FL 33487-5221, USA; charles.peterfy@spiresciences.com

PGC and PE are joint senior authors.

Received 16 July 2015
Revised 30 November 2015
Accepted 16 January 2016
Published Online First
10 February 2016

ABSTRACT

Objectives To assess structural damage progression with subcutaneous abatacept (ABA) in the Assessing Very Early Rheumatoid arthritis Treatment (AVERT) trial following abrupt withdrawal of all rheumatoid arthritis (RA) medication in patients achieving Disease Activity Score (DAS)-defined remission or low disease activity.

Methods Patients with early, active RA were randomised to ABA plus methotrexate (ABA/MTX) 125 mg/week, ABA 125 mg/week or MTX for 12 months. All RA treatments were withdrawn after 12 months in patients with DAS28 (C reactive protein (CRP)) <3.2. Adjusted mean changes from baseline in MRI-based synovitis, osteitis and erosion were calculated for the intention-to-treat population.

Results 351 patients were randomised and treated: ABA/MTX (n=119), ABA (n=116) or MTX (n=116). Synovitis and osteitis improved, and progression of erosion was statistically less with ABA/MTX versus MTX at month 12 (−2.35 vs −0.68, −2.58 vs −0.68, 0.19 vs 1.53, respectively; p<0.01 for each) and month 18 (−1.34 vs −0.49, −2.03 vs 0.34, 0.13 vs 2.0, respectively; p<0.01 for erosion); ABA benefits were numerically intermediate to those for ABA/MTX and MTX.

Conclusions Structural benefits with ABA/MTX or ABA may be maintained 6 months after withdrawal of all treatments in patients who have achieved remission or low disease activity.

Trial registration number NCT01142726; Results.

INTRODUCTION

Abatacept is a cytotoxic T lymphocyte-associated antigen-4 (CTLA-4)–immunoglobulin G1 fusion protein¹ that selectively modulates the CD80/CD86:CD28 costimulatory pathway required for T-cell activation.² In people with rheumatoid arthritis (RA), treatment with abatacept has been shown to normalise levels of many inflammatory mediators associated with disease activity and progression.¹

The Assessing Very Early Rheumatoid arthritis Treatment (AVERT) trial evaluated clinical remission at 12 months with subcutaneous abatacept plus methotrexate (MTX), abatacept monotherapy or MTX alone in patients with early RA and maintenance of remission following rapid withdrawal of

all RA treatments.³ Patients treated with abatacept in combination with MTX achieved higher rates of protocol-defined Disease Activity Score (DAS) remission (DAS28 C reactive protein (CRP) <2.6) versus MTX alone, and a small but significantly higher number of patients achieved sustained DAS-defined remission following withdrawal of all RA treatments.³ The safety profile of abatacept, in combination with MTX or as monotherapy, was comparable with that of MTX alone.³

The AVERT trial evaluated clinical and also MRI outcomes. MRI is increasingly being used in clinical trials to assess the therapeutic efficacy and can detect early changes in inflammation—including synovitis, osteitis and joint damage—before radiographic joint erosion occurs;⁴ in clinical studies, MRI measures of inflammation have even been shown to predict radiographic progression.^{5–7} In patients with established RA, the benefits of abatacept in reducing synovitis and osteitis have previously been demonstrated using MRI.^{8,9}

In this substudy of the AVERT trial, we report MRI measures of disease in patients with DAS-defined remission (DAS28 (CRP) <2.6) following 12 months of treatment and subsequent withdrawal of all RA medication, for up to 18 months.

METHODS**Study design and patient population**

The trial design and primary results for AVERT (NCT01142726) have been described previously.³ Briefly, patients were randomised (1:1:1) to receive abatacept 125 mg/week plus MTX, abatacept monotherapy (125 mg/week) or MTX alone for 12 months. At month 12, patients with DAS28 (CRP) <3.2 could enter the withdrawal period and all treatments for RA were stopped, with MTX and corticosteroids being tapered during the first month.

Assessments

All patients underwent contrast-enhanced 1.5 T MRI of the clinically most active wrist and hand (metacarpophalangeal joints 1–5) at baseline and at 6-month intervals thereafter; the same wrist/hand was imaged at each designated visit (see online



Open Access
Scan to access more
free content



CrossMark

To cite: Peterfy C, Burmester GR, Bykerk VP, et al. *Ann Rheum Dis* 2016;**75**:1501–1505.

supplementary material). MRI assessments were evaluated at baseline and up to month 18; only patients with baseline and postbaseline MRI assessments were included (see online supplementary material).

Statistical analysis

Adjusted mean changes from baseline in MRI scores for synovitis, osteitis and erosion were calculated at months 12 and 18 in the intention-to-treat population, and analysed using a longitudinal repeated measures model adjusting for MRI score and corticosteroid use (yes/no) at baseline and presented by treatment group; observed data at all time points were included. For all mean response rates, SE, 95% CIs and p values were calculated (see online supplementary material).

The proportion of patients achieving DAS-defined remission and MRI non-progression (defined as change from baseline \leq smallest detectable change separately for erosion, osteitis and synovitis) was calculated at months 6, 12 and 18 as an exploratory endpoint (see online supplementary material). The percentage change in CRP from baseline was evaluated over time. Additionally, *post hoc* analyses included the number of patients achieving DAS-defined remission and MRI progression (defined as change from baseline $>$ smallest detectable change) at month 18; the proportion of patients with a synovitis score >5 at baseline and at months 6, 12 and 18; and adjusted mean changes from baseline in MRI scores in the subgroup of patients who had DAS28 (CRP) <2.6 at both months 12 and 18.

RESULTS

Patient disposition and baseline characteristics

A total of 511 patients were enrolled, and 351 patients at 72 worldwide sites were randomly assigned (1:1:1) to treatment with abatacept plus MTX (n=119), abatacept monotherapy (n=116) or MTX (n=116). The proportion of patients with MRI assessments at baseline was 95.8% (114/119), 96.6% (112/116) and 95.7% (111/116), respectively; at baseline and month 12, it was 76.5% (91/119), 69.8% (81/116) and 72.4% (84/116), respectively; and at baseline, month 12 and month 18, it was 31.9% (38/119), 30.2% (35/116) and 25.0% (29/116), respectively. The average rate of missing MRI scores for those who had assessments at baseline was 16.9% over 12 months and 69.9% at 18 months. A total of 129/202 patients (63.9%) who entered the withdrawal period and who had MRI assessments at baseline and at month 12 discontinued during the withdrawal period (abatacept plus MTX, n=48 (37.2%); abatacept monotherapy, n=40 (31.0%); MTX, n=41 (31.8%)). The main reason for discontinuation was lack of efficacy (n=123/202; 60.9%). Clinical baseline characteristics for patients with MRI assessments at baseline and at month 12 were similar across treatment groups and were comparable with the overall AVERT population (table 1).

Efficacy

Abatacept plus MTX resulted in significantly greater decreases from baseline in synovitis and osteitis scores on-treatment, and significantly less progression of erosion score on-treatment and following withdrawal of all therapies than MTX alone (figure 1). While mean MRI synovitis and osteitis scores increased following withdrawal of treatment in all three groups, the adjusted mean reductions from baseline with abatacept plus MTX at month 18 were still numerically greater than those with MTX alone (figure 1A, B). Changes in erosion score showed minimal difference between months 6 and 18 (figure 1C). Benefits of abatacept monotherapy were numerically intermediate to those of

Table 1 Baseline characteristics for patients with MRI assessments at baseline and month 12

Characteristic	Abatacept plus MTX (n=91)	Abatacept monotherapy (n=81)	MTX (n=84)
Synovitis score	5.6 (4.2)	5.3 (3.8)	5.7 (4.2)
Osteitis score	4.4 (6.9)	4.3 (6.7)	3.4 (6.4)
Erosion score	7.2 (7.0)	5.1 (4.7)	6.3 (7.8)
Symptom duration, years	0.6 (0.5)	0.6 (0.5)	0.5 (0.5)
DAS28 (CRP)	5.5 (1.3)	5.5 (1.2)	5.3 (1.3)
HAQ-DI	1.5 (0.7)	1.4 (0.6)	1.4 (0.6)
Tender Joint Count (28 joints)	13.8 (7.8)	14.3 (7.7)	12.5 (7.7)
Swollen Joint Count (28 joints)	11.2 (7.1)	12.4 (7.6)	10.2 (6.9)
CRP, mg/L	17.5 (23.2)	16.6 (25.6)	16.0 (21.2)

Data are presented as mean (SD).

CRP, C reactive protein; DAS, Disease Activity Score; HAQ-DI, Health Assessment Questionnaire-Disability Index; MTX, methotrexate.

abatacept plus MTX and MTX alone. Cumulative probability plots for changes in MRI scores from baseline at month 12 are shown in online supplementary figure S1.

During study treatment, there was a reduction in the proportion of patients with active synovitis (score >5 ; indicative of active disease resulting in erosion progression¹⁰) in the abatacept plus MTX group versus that in the MTX alone group; at month 18, there was a numerical increase versus MTX alone (see online supplementary figure S1).

During study treatment, a statistically higher percentage of patients receiving abatacept plus MTX achieved DAS-defined remission together with MRI non-progression in synovitis, osteitis and erosion compared with those receiving MTX alone (table 2). Fewer patients in all three treatment groups achieved DAS-defined remission together with MRI non-progression after withdrawal of study drug compared with on-treatment (table 2). However, the percentages of patients in the abatacept plus MTX group were still approximately twice those of the MTX-alone group. The effect of abatacept monotherapy was numerically intermediate to that of abatacept plus MTX and MTX alone. In all treatment groups, a small number of patients still had MRI progression (see online supplementary material).

Post hoc analyses revealed that, in the small proportion of patients with DAS-defined remission at both months 12 and 18, MRI benefits achieved at month 12 were maintained up to month 18 (ie, after withdrawal of all therapies) in all three groups (see online supplementary figure S3).

Abatacept plus MTX treatment resulted in greater decreases from baseline in the inflammatory marker, CRP, during the treatment period and following the withdrawal of all therapies compared with MTX alone. Abatacept monotherapy and MTX alone had similar effects on CRP (see online supplementary table S1).

DISCUSSION

In this analysis of tissue effects, abatacept reduced inflammation and structural damage progression as assessed by changes in MRI scores (synovitis, osteitis and bone erosions) in patients with early and progressive RA. Following withdrawal of all therapies in patients with clinical disease activity, there was some reappearance of inflammatory activity in the joints. However, in those patients who maintained clinical remission or low disease




Figure 1 Adjusted mean change from baseline in MRI scores (ITT population). Change from baseline in (A) synovitis scores, (B) osteitis and (C) erosion at 6, 12 and 18 months. *p<0.05, **p<0.01 and ***p<0.001 for adjusted mean difference versus MTX at given time point. Error bars represent SEs. ITT, intention-to-treat; MTX, methotrexate.

activity following withdrawal of all RA therapy, this remission/low disease activity correlated with maintenance of inflammation control and structural benefits irrespective of initial treatment group. During both the treatment and withdrawal periods, improvements as assessed by MRI were greatest for abatacept plus MTX, followed by abatacept monotherapy and MTX monotherapy.

The improvements in MRI scores observed in the present study are consistent with previous abatacept studies^{9, 11} and recent randomised controlled trials of other biologics in patients with early or established RA (see online supplementary material).^{12–16} The

data presented here are also consistent with the maintenance of MRI benefits for 6 months following the withdrawal of abatacept in patients with early RA reported in the ADJUST trial.⁹ Only a few studies of biologics have evaluated treatment withdrawal or de-escalation and imaging outcomes (MRI or radiography) in patients with RA.^{17–20} Consistent with our study, Smolen *et al* showed that most patients with early RA (<1 year disease duration) who achieved a stable low-dose disease activity target at ~6 months maintained structural benefits, despite withdrawal of adalimumab. Similarly, induction treatment with MTX plus adalimumab was shown to result in fewer erosions at 1 year versus

Table 2 Proportion of patients with DAS28 (CRP) remission and MRI non-progression per MRI pathology over time (ITT population)

MRI pathology	Study time point (months)	Patients with DAS28 (CRP) remission* and MRI non-progression†			Abatacept plus MTX vs MTX estimate of difference (95% CI)	Abatacept monotherapy vs MTX estimate of difference (95% CI)
		Abatacept plus MTX (n=119)	Abatacept monotherapy (n=116)	MTX (n=116)		
Synovitis—n‡ (%) (95% CI)§	6	44 (37.0) (28.30 to 45.65)	32 (27.6) (19.45 to 35.72)	28 (24.1) (16.35 to 31.93)	12.84 (0.33 to 25.34); p=0.046	3.45 (−8.67 to 15.57); p=0.653
	12	59 (49.6) (40.60 to 58.56)	43 (37.1) (28.28 to 45.86)	41 (35.3) (26.65 to 44.04)	14.24 (0.88 to 27.59); p=0.038	1.72 (−11.50 to 14.95); p=0.891
	18	18 (15.1) (8.69 to 21.56)	11 (9.5) (4.15 to 14.81)	9 (7.8) (2.89 to 12.63)	7.37 (−1.55 to 16.29); p=0.117	1.72 (−6.36 to 9.81); p=0.815
Osteitis—n‡ (%) (95% CI)§	6	43 (36.1) (27.50 to 44.77)	32 (27.6) (19.45 to 35.72)	27 (23.3) (15.6 to 30.97)	12.86 (0.45 to 25.27); p=0.044	4.31 (−7.75 to 16.37); p=0.546
	12	58 (48.7) (36.76 to 57.72)	42 (36.2) (27.46 to 44.95)	41 (35.3) (26.65 to 44.04)	13.39 (0.04 to 26.75); p=0.052	0.86 (−12.34 to 14.06); p=1.000
	18	18 (15.1) (8.69 to 21.56)	12 (10.3) (4.80 to 15.89)	9 (7.8) (2.89 to 12.63)	7.37 (−1.55 to 16.29); p=0.117	2.59 (−5.65 to 10.82); p=0.647
Erosion—n‡ (%) (95% CI)§	6	38 (31.9) (23.56 to 40.31)	30 (25.9) (17.89 to 33.83)	23 (19.8) (12.57 to 27.08)	12.11 (0.17 to 24.04); p=0.049	6.03 (−5.60 to 17.67); p=0.348
	12	51 (42.9) (33.97 to 51.75)	36 (31.0) (22.62 to 39.45)	33 (28.4) (20.24 to 36.66)	14.41 (1.46 to 27.36); p=0.030	2.59 (−10.04 to 15.21); p=0.774
	18	15 (12.6) (6.64 to 18.57)	11 (9.5) (4.15 to 14.81)	8 (6.9) (2.29 to 11.51)	5.71 (−2.68 to 14.10); p=0.210	2.59 (−5.32 to 10.50); p=0.632

*DAS-defined remission was defined as DAS28 (CRP) <2.6.

†MRI non-progression was defined as change from baseline \leq smallest detectable change. Smallest detectable change values used for non-progression: synovitis (2.01), osteitis (2.81) and erosion (2.29).

‡n=number of patients with DAS28 (CRP) remission and MRI non-progression.

§Normal approximation is used if the number of DAS28 (CRP) remission and MRI non-progression for all treatment arms was at least 5; otherwise an exact method was used. CRP, C reactive protein; DAS, Disease Activity Score; ITT, intention-to-treat; MTX, methotrexate.

MTX alone in patients who discontinued adalimumab after 6 months' treatment.^{17–20}

Some limitations of the present study should be taken into account. The *post hoc* analysis of change from baseline in MRI scores in patients with DAS28 (CRP) <2.6 at both months 12 and 18 had a small sample size, and therefore additional studies are needed to confirm these results, despite the sensitivity of MRI for detecting change. Additionally, the study was not powered to compare abatacept combination therapy with monotherapy.

In conclusion, these data from the AVERT study demonstrate that abatacept reduces inflammation and structural damage progression as assessed by MRI, and extends the current understanding of the effect of abatacept on synovium and bone. The withdrawal of abatacept therapy after achieving remission in some patients with early, progressive RA appears to be possible without an associated risk of joint damage progression.

Author affiliations

¹Spire Sciences, Inc., Boca Raton, Florida, USA

²Department of Rheumatology and Clinical Immunology, Charité—University Medicine Berlin, Berlin, Germany

³Department of Rheumatology, Hospital for Special Surgery, Weill Cornell Medical College, New York, New York, USA

⁴Department of Rheumatology, Service d'Immuno-Rheumatologie, Montpellier, France

⁵Department of Medicine, University of California Los Angeles, Los Angeles, California, USA

⁶Department of Rheumatology, Leiden University Medical Center, Leiden, The Netherlands

⁷Bristol-Myers Squibb, Princeton, New Jersey, USA

⁸Leeds Institute of Rheumatic and Musculoskeletal Medicine, University of Leeds, Leeds, UK

⁹NIHR Leeds Musculoskeletal Biomedical Research Unit, Leeds Teaching Hospitals NHS Trust, Leeds, UK

Correction notice This article has been corrected since it was published Online First. In table 1 the units for CRP have been corrected to mg/L.

Acknowledgements The first draft of the manuscript was prepared by academic and industry authors, with professional medical writing and editorial assistance provided by Stacey Reeber, PhD, at Caudex and funded by Bristol-Myers Squibb. The academic authors vouch for the completeness and accuracy of the data and data analyses, and for the fidelity of the study to the protocol.

Contributors CP and PGC contributed to the MRI study design and data acquisition/interpretation. GRB, VPB, BGC, DEF, TWJH and PE contributed to data acquisition and interpretation. JCD and DAW contributed to data analysis and interpretation. All authors assisted in the drafting and critical revision of the manuscript and agree to be accountable for all aspects of the work.

Funding The study was funded by Bristol-Myers Squibb.

Competing interests CP: consulting fees: AbbVie, Acerta, Amgen, AstraZeneca, Bristol-Myers Squibb, Centrexion, Daiichi Sankyo, Five Prime Therapeutics, Genentech, Hoffmann-La Roche, Janssen, Lilly USA, MedImmune, Merck, Novartis, Plexikon, Pfizer, Sanofi, Salix-Santarus, Samsung; speakers bureau: Amgen; Founder and CEO of Spire Sciences, which provides imaging services to multiple pharmaceutical companies. GRB: grant/research support: Bristol-Myers Squibb, AbbVie, Pfizer, MedImmune, Novartis, Roche, UCB, Lilly; consultant: Bristol-Myers Squibb, AbbVie, Pfizer, MSD, MedImmune, Roche, UCB; speakers bureau: Bristol-Myers Squibb, AbbVie, Pfizer, MSD, Roche, UCB. VPB: Consultant: Amgen, Antares, Bristol-Myers Squibb, Genentech, Pfizer, UCB. BGC: grant/research support: Pfizer, Roche-Chugai, UCB; speakers bureau: Bristol-Myers Squibb, Janssen, Lilly, Merck, Novartis, Pfizer, Roche-Chugai, UCB. JCD: consulting: AbbVie, Acerta, Amgen, AstraZeneca, Bristol-Myers Squibb, Centrexion, Daiichi Sankyo, Five Prime Therapeutics, Genentech, Hoffmann-La Roche, Janssen, Lilly USA, MedImmune, Merck, Novartis, Plexikon, Pfizer, Sanofi, Salix-Santarus, Samsung; Employee of Spire Sciences, Inc., which provides imaging services to multiple pharmaceutical companies. DEF: grant/research support: AbbVie, Actelion, Amgen, Bristol-Myers Squibb, Gilead, GSK, NIH, Novartis, Pfizer, Roche/Genentech, UCB; consultant: AbbVie, Actelion, Amgen, Bristol-Myers Squibb, Cytos, Janssen, Gilead, GSK, NIH, Novartis, Pfizer, Roche/Genentech, UCB; speakers bureau (CME only): AbbVie, Actelion, UCB. TWJH: consulting fees: Abbott Laboratories, Biotest AG, Bristol-Myers Squibb, Crescendo Bioscience, Inc., Novartis Pharmaceuticals Corporation, Pfizer, Inc., Roche, Sanofi-Aventis, Schering-Plough, UCB, Inc., Eli Lilly; non-remunerative positions of influence: Meteor Board; research grants: EU & Dutch Arthritis

Foundation; speakers bureau: Abbott Laboratories, Biotest AG, Bristol-Myers Squibb, Novartis, Pfizer, Roche, Sanofi-Aventis, Schering-Plough; travel support: Abbott Laboratories, Roche. DAW: shareholder and former employee of Bristol-Myers Squibb. PGC: speakers bureau or consultancies: AbbVie, Bristol-Myers Squibb, Janssen, Novartis, Roche. PE: consulting fees: AbbVie, Bristol-Myers Squibb, Lilly, Merck, Novartis, Pfizer, Roche, Samsung, Takeda, UCB; grant/research support: AbbVie, Merck, Pfizer, Roche.

Ethics approval The AVERT study protocol was approved by the Institutional Review Board or Independent Ethics Committee at each site.

Provenance and peer review Not commissioned; externally peer reviewed.

Open Access This is an Open Access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

REFERENCES

- Herrero-Beaumont G, Martínez Calatrava MJ, Castañeda S. Abatacept mechanism of action: concordance with its clinical profile. *Rheumatol Clin* 2012;8:78–83.
- Lenschow DJ, Walunas TL, Bluestone JA. CD28/B7 system of T cell costimulation. *Annu Rev Immunol* 1996;14:233–58.
- Emery P, Burmester GR, Bykerk VP, et al. Evaluating drug-free remission with abatacept in early rheumatoid arthritis: results from the phase 3b, multicentre, randomised, active-controlled AVERT study of 24 months, with a 12-month, double-blind treatment period. *Ann Rheum Dis* 2015;74:19–26.
- McQueen FM. Imaging in early rheumatoid arthritis. *Best Pract Res Clin Rheumatol* 2013;27:499–522.
- McQueen FM, Benton N, Perry D, et al. Bone edema scored on magnetic resonance imaging scans of the dominant carpus at presentation predicts radiographic joint damage of the hands and feet six years later in patients with rheumatoid arthritis. *Arthritis Rheum* 2003;48:1814–27.
- Hetland ML, Ejlertsen B, Hørslev-Petersen K, et al. MRI bone oedema is the strongest predictor of subsequent radiographic progression in early rheumatoid arthritis. Results from a 2-year randomised controlled trial (CIMESTRA). *Ann Rheum Dis* 2009;68:384–90.
- Baker JF, Ostergaard M, Emery P, et al. Early MRI measures independently predict 1-year and 2-year radiographic progression in rheumatoid arthritis: secondary analysis from a large clinical trial. *Ann Rheum Dis* 2014;73:1968–74.
- Conaghan PG, Durez P, Alten RE, et al. Impact of intravenous abatacept on synovitis, osteitis and structural damage in patients with rheumatoid arthritis and an inadequate response to methotrexate: the ASSET randomised controlled trial. *Ann Rheum Dis* 2013;72:1287–94.
- Emery P, Durez P, Dougados M, et al. Impact of T-cell costimulation modulation in patients with undifferentiated inflammatory arthritis or very early rheumatoid arthritis: a clinical and imaging study of abatacept (the ADJUST trial). *Ann Rheum Dis* 2010;69:510–16.
- Gandjbakhch F, Haavardsholm EA, Conaghan PG, et al. Determining a magnetic resonance imaging inflammatory activity acceptable state without subsequent radiographic progression in rheumatoid arthritis: results from a followup MRI study of 254 patients in clinical remission or low disease activity. *J Rheumatol* 2014;41:398–406.
- Mease P, Genovese MC, Gladstein G, et al. Abatacept in the treatment of patients with psoriatic arthritis: results of a six-month, multicenter, randomized, double-blind, placebo-controlled, phase II trial. *Arthritis Rheum* 2011;63:939–48.
- Peterfy C, Haraoui B, Durez P, et al. Decreased incidence of synovitis, osteitis, and erosion in early RA patients treated with adalimumab plus methotrexate compared to those with methotrexate alone: high-field MRI analysis from OPTIMA (Abstract 123). *Arthritis Rheum* 2010;62:S51.
- Østergaard M, Emery P, Conaghan PG, et al. Significant improvement in synovitis, osteitis, and bone erosion following golimumab and methotrexate combination therapy as compared with methotrexate alone: a magnetic resonance imaging study of 318 methotrexate-naïve rheumatoid arthritis patients. *Arthritis Rheum* 2011;63:3712–22.
- Conaghan PG, Peterfy C, Olech E, et al. The effects of tocilizumab on osteitis, synovitis and erosion progression in rheumatoid arthritis: results from the ACT-RAY MRI substudy. *Ann Rheum Dis* 2014;73:810–16.
- Peterfy C, Emery P, Tak PP, et al. MRI assessment of suppression of structural damage in patients with rheumatoid arthritis receiving rituximab: results from the randomised, placebo-controlled, double-blind RA-SCORE study. *Ann Rheum Dis* 2016;75:170–7.
- Peterfy C, Emery P, Genovese M, et al. Magnetic resonance imaging substudy in a phase 2b dose-ranging study of baricitinib, an oral janus kinases 1/janus kinase 2 inhibitor, in combination with traditional disease-modifying antirheumatic drugs in patients with rheumatoid arthritis. *Arthritis Rheum* 2012;64 (Suppl 10):1050.
- Smolen JS, Emery P, Fleischmann R, et al. Adjustment of therapy in rheumatoid arthritis on the basis of achievement of stable low disease activity with adalimumab plus methotrexate or methotrexate alone: the randomised controlled OPTIMA trial. *Lancet* 2014;383:321–32.
- Quinn MA, Conaghan PG, O'Connor PJ, et al. Very early treatment with infliximab in addition to methotrexate in early, poor-prognosis rheumatoid arthritis reduces magnetic resonance imaging evidence of synovitis and damage, with sustained benefit after infliximab withdrawal: results from a twelve-month randomised, double-blind, placebo-controlled trial. *Arthritis Rheum* 2005;52:27–35.
- Emery P, Hammoudeh M, FitzGerald O, et al. Sustained remission with etanercept tapering in early rheumatoid arthritis. *N Engl J Med* 2014;371:1781–92.
- Detert J, Bastian H, Listing J, et al. Induction therapy with adalimumab plus methotrexate for 24 weeks followed by methotrexate monotherapy up to week 48 versus methotrexate therapy alone for DMARD-naïve patients with early rheumatoid arthritis: HIT HARD, an investigator-initiated study. *Ann Rheum Dis* 2013;72:844–50.

SUPPLEMENTARY MATERIALS

Assessments The wrist and hand were scanned separately with surface coils using a positioning frame and biplanar slice alignment to ensure reproducibility. Pulse sequences included coronal 2-dimensional short-tau inversion recovery (STIR) and coronal fat-saturated, T1-weighted 3-dimensional (3D) gradient echo (3D GRE) with and without intravenous gadolinium-based contrast. Voxel dimensions were $469\mu \times 625\mu \times 3,000\mu$ for STIR and $203\mu \times 625\mu \times 1,500\mu$ for 3D GRE.

MRI images were evaluated at a central reading facility by two independent radiologists using the Outcome Measures in Rheumatology (OMERACT) Rheumatoid Arthritis Magnetic Resonance Imaging Scoring (RAMRIS) system [1]: synovitis (scale, 0–3 in 8 joints), osteitis (scale, 0–3 in 25 bones) and bone erosion (scale, 0–10 in 25 bones). All time points for an individual patient were viewed simultaneously but in random order and with the dates masked. The scores of the two radiologists were averaged, and the top 5% of discrepancies for total change scores relative to baseline for each RAMRIS feature were adjudicated by consensus review. The interclass correlation coefficients (ICCs) for synovitis, osteitis and erosion were 0.94, 0.96 and 0.91 at baseline, respectively. ICC values for change from baseline to the last time point available for each patient were 0.89, 0.95 and 0.82, respectively.

Statistical analysis

Adjusted mean changes from baseline in magnetic resonance imaging (MRI) scores were calculated at Months 12 and 18 and analysed in the intent-to-treat (ITT) population (all randomised patients who received at least one dose of study medication in the treatment period, with patients analysed according to randomised treatment). If the score for a given parameter (erosion, osteitis or synovitis) was missing for >20% of joints, the MRI score for each parameter was considered missing. A linear mix model was used to handle missing data when a series of outcomes were measured repeatedly over time. Under the Missing At Random (MAR) assumption, the methods above provided an unbiased estimate of the treatment effect that would have been observed if all patients had continued on treatment for the full study duration.

Adjusted mean differences between treatment groups and associated standard errors (SEs) and two-sided 95% confidence intervals (CIs) were also calculated for abatacept plus MTX and abatacept monotherapy versus MTX alone using the same model. Although no formal statistical testing was performed for these secondary MRI efficacy endpoints, p values were calculated.

Response rates were calculated as point estimates with associated 95% CI for treatment difference. The 95% CIs for a response rate within treatment group were based on normal approximation, provided there were at least five events in each treatment group; otherwise, the exact method was used. The 95% CI for treatment difference in response rate was constructed using the continuity correction.

For the proportion of patients achieving DAS-defined remission (DAS28 [CRP] <2.6) and MRI non-progression, missing DAS-defined remission not due to premature discontinuation and not at treatment period Day 1 or withdrawal period Day 169 was imputed as DAS-defined remission if the missing value was between two observed DAS-defined remissions.

Results

A small number of patients who achieved DAS-defined remission at Month 18 still had MRI progression in synovitis, osteitis and erosion, respectively: abatacept plus MTX (0, 0, 1 patients, respectively); abatacept monotherapy (1, 0, 1 patients) and MTX alone (1, 1, 2 patients).

Discussion

MRI assessments following abatacept treatment have previously been reported in the ADJUST trial in MTX-naïve patients with undifferentiated inflammatory arthritis or very early RA, in the ASSET trial in patients with established RA and an inadequate response to MTX and in patients with psoriatic arthritis.[2-4] A notable difference between ASSET and this study was that the low baseline mean synovitis scores in the ASSET cohort (due to the use of wrist synovitis scores alone being included) made it difficult to show a difference in synovitis outcomes.[4] Data from the current analysis show that abatacept does reduce joint inflammation as indicated by osteitis in patients with early RA.

References

1. Ostergaard M, Peterfy C, Conaghan P, et al. OMERACT Rheumatoid Arthritis Magnetic Resonance Imaging Studies. Core set of MRI acquisitions, joint pathology definitions, and the OMERACT RA-MRI scoring system. *J Rheumatol* 2003;30:1385-6.




2. Mease P, Genovese MC, Gladstein G, et al. Abatacept in the treatment of patients with psoriatic arthritis: results of a six-month, multicenter, randomized, double-blind, placebo-controlled, phase II trial. *Arthritis Rheum* 2011;63:939-48.
3. Emery P, Durez P, Dougados M, et al. Impact of T-cell costimulation modulation in patients with undifferentiated inflammatory arthritis or very early rheumatoid arthritis: a clinical and imaging study of abatacept (the ADJUST trial). *Ann Rheum Dis* 2010;69:510-6.
4. Conaghan PG, Durez P, Alten RE, et al. Impact of intravenous abatacept on synovitis, osteitis and structural damage in patients with rheumatoid arthritis and an inadequate response to methotrexate: the ASSET randomised controlled trial. *Ann Rheum Dis* 2013;72:1287-94.

Supplementary Table 1 Mean (SE) change in CRP from baseline

	Abatacept plus MTX (n=119)	Abatacept monotherapy (n=116)	MTX (n=116)
6 months	-11.99 (2.08) (-16.12, -7.87)	-9.68 (2.84) (-15.31, -4.06)	-9.14 (2.23) (-13.57, -4.72)
12 months	-12.73 (2.24) (-17.18, -8.27)	-11.63 (2.49) (-16.58, -6.68)	-8.01 (2.27) (-12.51, -3.50)
18 months (withdrawal period)	-6.72 (3.06) (-12.87, -0.56)	-2.07 (2.63) (-8.01, 2.62)	-4.37 (4.02) (-12.54, 3.81)


Months 6 and 12 represent study treatment period. 95% confidence intervals are presented.

Supplementary Figure 1 Cumulative probability plots of change from baseline in MRI scores. Change from baseline in (A) synovitis (B) osteitis (C) erosion scores at Month 12. SDC values were 2.01, 2.81 and 2.29, respectively. MRI, magnetic resonance imaging; SDC, smallest detectable change.

A**B****C**

Supplementary Figure 2 Proportion of patients with a synovitis score >5 at 6, 12 and 18 months.

Error bars represent 95% CIs. All randomized and treated patients (ITT population) with synovitis score measurement at baseline. The total population is presented; n, includes all patients irrespective of remission status. CI, confidence interval; ITT, intent-to-treat; MTX, methotrexate.



Supplementary Figure 3 Adjusted mean change from baseline in MRI scores; *post hoc* analysis in patients with DAS28 (CRP) <2.6 both at Month 12 and at Month 18. There were no significant differences versus MTX. MRI scores were adjusted for baseline and corticosteroid use at baseline (yes/no). Error bars represent standard errors; n=number of patients who are in remission. CRP, C-reactive protein; DAS, Disease Activity Score; MRI, magnetic resonance imaging; MTX, methotrexate.

