



OPEN ACCESS

CONCISE REPORT

A rare coding allele in *IFIH1* is protective for psoriatic arthritis

Ashley Budu-Aggrey,^{1,2} John Bowes,¹ Philip E Stuart,³ Matthew Zawistowski,⁴ Lam C Tsoi,^{3,4} Rajan Nair,³ Deepak Rohit Jadon,^{5,6} Neil McHugh,^{5,7} Eleanor Korendowych,⁵ James T Elder,^{3,8} Anne Barton,^{1,2} Soumya Raychaudhuri^{1,2,9,10}

► Additional material is published online only. To view, please visit the journal online (<http://dx.doi.org/10.1136/annrheumdis-2016-210592>).

For numbered affiliations see end of article.

Correspondence to

Professor Soumya Raychaudhuri, Arthritis Research UK Centre for Genetics and Genomics, Centre for Musculoskeletal Research, Institute of Inflammation and Repair, University of Manchester, Manchester Academic Health Science Centre, Stopford Building, Oxford Road, Manchester M13 9PT, UK; soumya@broadinstitute.org

Received 26 September 2016

Revised 1 March 2017

Accepted 12 March 2017

Published Online First

16 March 2017

ABSTRACT

Objectives Psoriatic arthritis (PsA) is an inflammatory arthritis associated with psoriasis. While many common risk alleles have been reported for association with PsA as well as psoriasis, few rare coding alleles have yet been identified.

Methods To identify rare coding variation associated with PsA risk or protection, we genotyped 41 267 variants with the exome chip and investigated association within an initial cohort of 1980 PsA cases and 5913 controls. Genotype data for an independent cohort of 2234 PsA cases and 5708 controls was also made available, allowing for a meta-analysis to be performed with the discovery dataset.

Results We identified an association with the rare variant rs35667974 ($p=2.39 \times 10^{-6}$, OR=0.47), encoding an Ile923Val amino acid change in the *IFIH1* gene protein product. The association was reproduced in our independent cohort, which reached a high level of significance on meta-analysis with the discovery and replication datasets ($p=4.67 \times 10^{-10}$). We identified a strong association with *IFIH1* when performing multiple-variant analysis ($p=6.77 \times 10^{-6}$), and found evidence of independent effects between the rare allele and the common PsA variant at the same locus.

Conclusion For the first time, we report a rare coding allele in *IFIH1* to be protective for PsA. This rare allele has also been identified to have the same direction of effect on type I diabetes and psoriasis. While this association further supports existing evidence for *IFIH1* as a causal gene for PsA, mechanistic studies will need to be pursued to confirm that *IFIH1* is indeed causal.

INTRODUCTION

In approximately 6%–42% of cases psoriasis is associated with an inflammatory arthritis,¹ psoriatic arthritis (PsA), which includes joint inflammation, involving primarily the distal interphalangeal (DIP) joints, entheses and sacroiliac joints. As patients typically test negative for rheumatoid factor, PsA is considered one of the spondyloarthropathies.¹ Patients with PsA have a worse quality of life than patients with psoriatic skin disease alone.² PsA may demonstrate a greater genetic component than psoriasis; a recent family study among first-degree relatives estimated a sibling recurrence risk (λ_s) between 30 and 40, greater than that of psoriasis, where the recurrence risk is between 4 and 10.^{3,4}

Genome-wide association studies have identified common risk alleles that are associated with

both psoriasis and PsA.^{5–10} Familial studies¹¹ and case-control studies¹² have reported rare missense alleles in *CARD14* that confer strong risk of psoriasis, but to date no rare coding alleles have been identified for PsA risk. As rare coding variants are more likely to result in an interpretable functional effect, they might more easily provide insight into the mechanism of disease than known common variants associated with psoriasis and PsA. If rare alleles are protective, have a strong effect and result in loss of gene function, they could be flagging promising drug targets.¹³ For example, the Y142X, C679X and R46L alleles in *PCSK9* identified this gene as a promising drug target for lipid modification.^{14,15}

To identify rare alleles that are associated with PsA (MAF<0.05), we queried the exome by analysing 1980 PsA cases and 5913 healthy controls that were genotyped using the Illumina HumanExome chip and HumanCoreExome chip. In doing so, we report an association with a rare coding variant at the *IFIH1* locus.

METHODS

Study sample description

We recruited a total of 2384 patients with PsA and 5946 healthy individuals in this study following approval of the research ethics committee (MRES 99/8/84), as described previously.¹⁶ All participants were the UK Caucasians and had provided written informed consent.

Genotyping

We genotyped DNA samples using the Infinium HumanExome-12 BeadChip (V.1-0) (Illumina) and the Infinium HumanCoreExome-24 BeadChip (V.1-0) (Illumina) (see online supplementary table 1), where 691 cases and all control samples were genotyped with the HumanExome chip, and the remaining 1693 cases were genotyped with the HumanCoreExome chip within the Arthritis Research UK Centre for Genetics and Genomics at The University of Manchester.

Genotype calling and quality control

We performed genotype calling and clustering with the GenomeStudio Data Analysis software (Genotyping Module V.1.8.4). We carried out initial genotype clustering to identify and exclude samples with a genotyping call rate <98%. We then performed automated reclustering of the remaining samples to



CrossMark

To cite: Budu-Aggrey A, Bowes J, Stuart PE, et al. *Ann Rheum Dis* 2017;**76**:1321–1324.

obtain more accurate genotype clusters. As we were concerned that rare variants may be incorrectly clustered, we conducted an extensive manual review of clusters. We reviewed variants that were filtered based on cluster separation (<0.4), signal intensity (<1.0) and allele frequency, where we identified variants that had failed genotyping, and those that had been incorrectly called as 'missing'.

Statistical quality control

We identified duplicated samples and related individuals using identity-by-descent with PLINK software (see online supplementary table 2). We also performed principal component analysis with the EIGENSOFT software package (V.6.0.1), to identify and exclude samples with divergent ancestry (see online supplementary figure 1, supplementary table 2). Principal components were calculated with a set of linkage disequilibrium (LD)-pruned single nucleotide polymorphisms (SNPs) ($n=13\,444$). We excluded monomorphic, Y chromosome, pseudoautosomal, mitochondrial and variants within the human leukocyte antigen (HLA) region, as well as variants with a call rate $<100\%$ in order to ensure high quality of the data. We also excluded SNPs found to significantly deviate from Hardy-Weinberg equilibrium ($<1.0 \times 10^{-6}$) among the control samples.

To prevent any bias due to differences in genotyping between the two platforms used in this study, we excluded all variants with evidence of case allele frequency differences between the two platforms ($p < 0.05$) (see online supplementary figure 2). We also excluded variants with an allele count of 0 among either of the case cohorts or the control samples. Furthermore, we excluded rare variants showing evidence of genotyping bias by assessing the allele counts of cases and controls for each genotyping platform used.

Replication dataset

We used an independent cohort of 2234 PsA cases and 5708 controls to replicate our findings. Briefly, people of European Caucasian ancestry were collected in North America and Sweden, and were genotyped using the Affymetrix Axiom Biobank Plus Genotyping Array. More details of this cohort, as well as quality control procedures applied to the genotype data, have been described previously.¹⁷

Association testing

We performed single-point analysis using the Fisher's exact test in PLINK with all SNPs passing quality control, and separately for rare variants. We calculated genomic inflation for all variants, and separately for rare and common variants alone, where we created a null SNP set ($n=13\,106$) with the common variants and excluded previously reported psoriasis loci (see online supplementary figures 3–5). We performed multiple-variant testing with the rare variants using SKAT analysis in R. We used the Firth logistic regression test (with four principal components as covariates) for association analysis of the replication cohort. In our replication cohort, principal components had been calculated with a set of LD-pruned SNPs, where known psoriasis loci had been excluded ($n=11\,005$). We performed a meta-analysis with the discovery and independent cohort data assuming an inverse-variance fixed-effects model. The choice of methods to perform the single-variant analysis was based on their suitability for analysing rare variants.

IFIH1 conditional analysis and conditional haplotype analysis

At the *IFIH1* locus, we performed conditional logistic regression in PLINK to test for independence between the rare (Ile923Val)

and common (rs984971) variants using PsA Immunochip genotype data (1962 cases, 8923 controls). We also used this dataset to construct haplotypes in PLINK to test for association with PsA, where we calculated OR and 95% CIs. We performed conditional haplotype analysis based on a likelihood ratio, to test for an effect of Ile923Val independent from that of rs984971.

RESULTS

After quality control, we examined genotype data for 41 267 non-HLA variants in 1980 cases and 5913 controls (see online supplementary table 3). We excluded more case samples compared with controls due to the overlap of case samples that were genotyped on both platforms. We carried out single-point analysis using the Fisher's exact test to identify associations with PsA. Unsurprisingly, we observed evidence of association in our cohort at known PsA alleles (see online supplementary table 4), including the rs33980500 SNP at *TRAF3IP2* ($p=3.9 \times 10^{-18}$). After removing these known associations, we observed only modest genomic inflation ($\lambda=1.06$, $\lambda_{1000,1000}=1.02$), suggesting little evidence of population stratification (see online supplementary figure 4).

We conducted a single-point analysis for a total of 27 066 rare variants alone ($MAF < 0.05$, $\lambda=0.79$) (see online supplementary figure 5). The strongest association we found was with the SNP rs35667974 ($p=2.39 \times 10^{-6}$, $OR=0.47$) (see online supplementary figures 6 and 7, supplementary table 5), mapping to *IFIH1*; this variant causes a missense mutation, resulting in an amino acid substitution of valine for isoleucine (Ile923Val). Of the 18 variants taken forward for replication ($p < 1.0 \times 10^{-3}$), we only found the Ile923Val association to replicate within our independent cohort ($p=3.5 \times 10^{-5}$, $OR=0.49$), based on a Bonferroni-corrected p value threshold of 2.78×10^{-3} ($=0.05/18$, see online supplementary tables 6 and 7). On meta-analysis with the discovery and replication datasets, we found the association with the Ile923Val allele to exceed genome-wide significance ($p=4.78 \times 10^{-10}$). We performed multiple-variant analysis to assess allele counts among genes and obtained further evidence to support the association of *IFIH1* with PsA ($p=6.77 \times 10^{-6}$, Bonferroni-corrected p value threshold $=5.60 \times 10^{-6}$, based on the number of genes tested) (see online supplementary table 8).

Independent effects at the *IFIH1* locus

A common variant at *IFIH1* has been previously reported as associated with PsA in our recent Immunochip study.⁹ When performing single-point analysis with the Immunochip dataset, we found evidence for an independent signal within the locus. In that study, we observed that Ile923Val obtained nominal significance after conditional analysis (figure 1). We also observed this association to be stronger than that previously reported for the common variant at the same locus, rs984971 ($p=3.62 \times 10^{-6}$). When we performed conditional logistic regression, the Ile923Val variant remained associated with PsA ($p < 0.05$) while conditioning on rs984971 ($p_{\text{cond}}=4.0 \times 10^{-6}$) (figure 1). Likewise, the SNP rs984971 remained associated with PsA when we conditioned on the Ile923Val variant ($p_{\text{cond}}=1.6 \times 10^{-4}$) (figure 1). This gives evidence of the independent effects of the two variants. We performed conditional haplotype analysis using haplotypes constructed with both variants (figure 2). Here, we found that the derived haplotype containing Ile923Val (CG) has a significantly greater protective effect compared with that containing the derived allele at rs984971 (TA), relative to the ancestral haplotype (TG) ($p=6.36 \times 10^{-11}$ vs $p_{\text{cond}}=2.95 \times 10^{-7}$). This confirms that the association with the rare *IFIH1* variant is not driven by the common rs984971 variant.




Figure 1 Conditional logistic regression within the *IFIH1* region using psoriatic arthritis (PsA) Immunochip genotype data (Bonferroni-corrected $p < 0.025$). (A) Association of the rare (Ile923Val) and common variant (rs984971) with PsA (Fisher's exact test). (B) The rare variant remains associated when rs984971 is conditioned on. (C) The common variant remains associated when the Ile923Val variant is conditioned on. ↓ Position of the SNP that has been conditioned on.

DISCUSSION

We have identified a PsA association with the rare coding allele Ile923Val within the *IFIH1* gene. This rare allele was found to be protective for PsA, and also replicated within an independent cohort. We found the association signal to reach genome-wide significance during meta-analysis, and confirmed the association with *IFIH1* when performing multiple-variant analysis. The Ile923Val allele has been previously reported to be protective for type 1 diabetes¹⁸ and psoriasis.¹⁹ Associations within the *IFIH1*




Figure 2 Frequencies and psoriatic arthritis association OR of the haplotypes for Ile923Val and rs984971. OR and their 95% CIs are shown relative to the most frequent haplotype on a natural log scale (ln). 1; Ile923Val, 2; rs984971, Freq; frequency.

locus have also been reported to increase PsA risk.^{10 20} However, until now no association with the Ile923Val allele has been reported to have a protective effect on PsA alone. We found the Ile923Val variant to be independent of the common SNP previously identified at the *IFIH1* locus (rs984971). This SNP is highly correlated ($r^2 = 0.64$) with an additional common *IFIH1* variant (rs1990760) that has been recently reported for PsA.¹⁰ We also found the Ile923Val variant to have an independent effect from this variant ($P_{\text{cond}} = 4.39 \times 10^{-6}$). Furthermore, we also found the Ile923Val variant to be independent of the common rs3747517 *IFIH1* variant previously reported for PsA ($P_{\text{cond}} = 7.73 \times 10^{-7}$).¹⁰ These findings highlight the relevance of *IFIH1* in immune-related disorders, especially given the recent association reported for the rare *IFIH1* allele with ankylosing spondylitis, Crohn's disease, psoriasis, primary sclerosing cholangitis and ulcerative colitis.²¹ The frequency of the Ile923Val variant in our control population (0.021) is slightly higher than observed in European populations in the 1000 genomes project and the exome sequencing project (0.01–0.018).^{22 23} This should be taken into consideration when interpreting the results of this study.

As the rare allele identified lies within a coding region, this provides strong evidence to suggest that *IFIH1* is a causal gene for PsA. As discussed by Plenge *et al*,¹³ a rare and independent variant in a causal gene for disease could provide a potential therapeutic target for treatment. However, linking human genetics with therapeutic targets requires the biological function of the causal gene and variant to be known.

IFIH1 encodes a cytoplasmic RNA helicase that recognises viral RNA and mediates an immune response on viral infection.¹⁸ The protective effect of the rare *IFIH1* allele observed with PsA suggests that the variant results in a loss of function phenotype, where the production or activity of *IFIH1* is decreased. Increased expression levels of *IFIH1* have been observed in psoriasis skin lesions compared with healthy skin,¹⁰ suggesting that pharmacological inhibition of this gene could be effective in treating psoriatic disease. However, when investigating the effect of Ile923Val in knockout mice, mutants showed no change in their double-stranded RNA binding activity compared with the wild-type.²⁴ This indicates the requirement for further investigation into the biological effect before any conclusions can be drawn from our current study regarding the impact of this allele on disease or the potential for treatment of PsA. Mechanistic studies will also be required to confirm that *IFIH1* is causal for PsA.

In conclusion, we report for the first time an association with a rare coding allele in *IFIH1* that is protective for PsA, and independent of associations that have previously been reported within this locus.

Author affiliations

¹Arthritis Research UK Centre for Genetics and Genomics, Centre for Musculoskeletal Research, University of Manchester, Manchester, UK

²NIHR Manchester Musculoskeletal Biomedical Research Unit, Central Manchester Foundation Trust and University of Manchester, Manchester Academy of Health Sciences, Manchester, UK

³Department of Dermatology, University of Michigan, Ann Arbor, Michigan, USA

⁴Department of Biostatistics, Center for Statistical Genetics, University of Michigan, Ann Arbor, Michigan, USA

⁵Royal National Hospital for Rheumatic Diseases, University of Bath, Bath, UK

⁶Department of Rheumatology, Cambridge University Hospitals NHSFT, Cambridge, UK

⁷Department of Pharmacy and Pharmacology, University of Bath, Bath, UK

⁸Ann Arbor Veterans Affairs Hospital, Ann Arbor, Michigan, USA

⁹Department of Medicine, Brigham and Women's Hospital and Harvard Medical School, Boston, Massachusetts, USA

¹⁰Program in Medical and Population Genetics, Broad Institute of MIT and Harvard, Cambridge, Massachusetts, USA

Contributors AB, SR, JB and JTE devised the study concept and design, and managed the project. AB-A performed genotyping and statistical analysis. AB-A, SR, AB, JB and JTE were responsible for the interpretation of the data. PS, MZ, LT and RN were responsible for the analysis of the Michigan cohort which was directed and coordinated by JTE. DP, NM and EK contributed data towards the discovery phase. All authors contributed to and approved the final version of the manuscript.

Funding We thank Arthritis Research UK for their support (grant ref 20 385). This report includes independent research funded by the NIHR Manchester Musculoskeletal BRU. AB-A is funded by a studentship from the MRC. This work was supported by awards from the National Institutes of Health (R01AR042742, R01AR050511, R01AR063611, R01AR065183) to JTE. JTE is supported by the Ann Arbor Veterans Affairs Hospital. LCT is supported by the Dermatology Foundation, National Psoriasis Foundation and the Arthritis National Research Foundation. This work was part-funded by an investigator-initiated research grant from Pfizer Ltd., awarded to DRJ and NMH.

Disclaimer The views expressed in the publication are those of the authors and not necessarily those of the National Health Service, the National Institute for Health Research or the Department of Health.

Competing interests None declared.

Provenance and peer review Not commissioned; externally peer reviewed.

Open Access This is an Open Access article distributed in accordance with the terms of the Creative Commons Attribution (CC BY 4.0) license, which permits others to distribute, remix, adapt and build upon this work, for commercial use, provided the original work is properly cited. See: <http://creativecommons.org/licenses/by/4.0/>

© Article author(s) (or their employer(s) unless otherwise stated in the text of the article) 2017. All rights reserved. No commercial use is permitted unless otherwise expressly granted.

REFERENCES

- Gladman DD, Antoni C, Mease P, *et al.* Psoriatic arthritis: epidemiology, clinical features, course, and outcome. *Ann Rheum Dis* 2005;64(Suppl 2):ii14–ii17.
- Rosen CF, Mussani F, Chandran V, *et al.* Patients with psoriatic arthritis have worse quality of life than those with psoriasis alone. *Rheumatology* 2012;51:571–6.
- Chandran V, Schentag CT, Brockbank JE, *et al.* Familial aggregation of psoriatic arthritis. *Ann Rheum Dis* 2009;68:664–7.
- Karason A, Love TJ, Gudbjornsson B. A strong heritability of psoriatic arthritis over four generations—the Reykjavik psoriatic arthritis study. *Rheumatology* 2009;48:1424–8.
- Ellinghaus E, Stuart PE, Ellinghaus D, *et al.* Genome-wide meta-analysis of psoriatic arthritis identifies susceptibility locus at REL. *J Invest Dermatol* 2012;132:1133–40.
- Hüffmeier U, Uebe S, Ekici AB, *et al.* Common variants at TRAF3IP2 are associated with susceptibility to psoriatic arthritis and psoriasis. *Nat Genet* 2010;42:996–9.
- Bowes J, Loehr S, Budu-Aggrey A, *et al.* PTPN22 is associated with susceptibility to psoriatic arthritis but not psoriasis: evidence for a further PsA-specific risk locus. *Ann Rheum Dis* 2015;74:1882–5.
- Zuo X, Sun L, Yin X, *et al.* Whole-exome SNP array identifies 15 new susceptibility loci for psoriasis. *Nat Commun* 2015;6:6793.
- Bowes J, Budu-Aggrey A, Hüffmeier U, *et al.* Dense genotyping of immune-related susceptibility loci reveals new insights into the genetics of psoriatic arthritis. *Nat Commun* 2015;6:6046.
- Stuart PE, Nair RP, Tsoi LC, *et al.* Genome-wide association analysis of psoriatic arthritis and cutaneous psoriasis reveals differences in their genetic architecture. *Am J Hum Genet* 2015;97:816–36.
- Jordan CT, Cao L, Roberson ED, *et al.* PSORS2 is due to mutations in CARD14. *Am J Hum Genet* 2012;90:784–95.
- Jordan CT, Cao L, Roberson ED, *et al.* Rare and common variants in CARD14, encoding an epidermal regulator of NF-kappaB, in psoriasis. *Am J Hum Genet* 2012;90:796–808.
- Plenge RM, Scolnick EM, Altshuler D. Validating therapeutic targets through human genetics. *Nat Rev Drug Discov* 2013;12:581–94.
- Cohen JC, Boerwinkle E, Mosley TH, *et al.* Sequence variations in PCSK9, low LDL, and protection against coronary heart disease. *N Engl J Med* 2006;354:1264–72.
- Stein EA, Mellis S, Yancopoulos GD, *et al.* Effect of a monoclonal antibody to PCSK9 on LDL cholesterol. *N Engl J Med* 2012;366:1108–18.
- Bowes J, Eyre S, Flynn E, *et al.* Evidence to support IL-13 as a risk locus for psoriatic arthritis but not psoriasis vulgaris. *Ann Rheum Dis* 2011;70:1016–9.
- Tsoi LC, Stuart PE, Tian C, *et al.* Large scale meta-analysis characterizes genetic architecture for common psoriasis-associated variants. *Nat Commun*. In Press.
- Nejentsev S, Walker N, Riches D, *et al.* Rare variants of IFIH1, a gene implicated in antiviral responses, protect against type 1 diabetes. *Science* 2009;324:387–9.
- Li Y, Liao W, Cargill M, *et al.* Carriers of rare missense variants in IFIH1 are protected from psoriasis. *J Invest Dermatol* 2010;130:2768–72.
- Julia A, Tortosa R, Hernanz JM, *et al.* Risk variants for psoriasis vulgaris in a large case-control collection and association with clinical subphenotypes. *Hum Mol Genet* 2012;21:4549–57.
- Ellinghaus D, Jostins L, Spain SL, *et al.* International IBD Genetics Consortium (IBDGC), International Genetics of Ankylosing Spondylitis Consortium (IGAS), International PSC Study Group (IPSCSG), Genetic Analysis of Psoriasis Consortium (GAPC), Psoriasis Association Genetics Extension (PAGE). Analysis of five chronic inflammatory diseases identifies 27 new associations and highlights disease-specific patterns at shared loci. *Nat Genet* 2016;48:510–8.
- Auton A, Brooks LD, Durbin RM, Abecasis GR, Altshuler DM, *et al.* 1000 Genomes Project Consortium. A global reference for human genetic variation. *Nature* 2015;526:68–74.
- Lek M, Karczewski KJ, Minikel EV, *et al.* Exome Aggregation Consortium. Analysis of protein-coding genetic variation in 60,706 humans. *Nature* 2016;536:285–91.
- Shigemoto T, Kageyama M, Hirai R, *et al.* Identification of loss of function mutations in human genes encoding RIG-I and MDA5: implications for resistance to type 1 diabetes. *J Biol Chem* 2009;284:13348–54.