tology Department, we retrospectively selected 35 patients that within one year from the first visit had a defined diagnosis of RA (according with the 2010 ACR/EULAR criteria) or of axial/peripheral SpA (according with the 2009 ASAS criteria). Demographic, clinical parameters and MCPs' MSUS assessment at baseline were recorded during the patients' first EA clinic visit by an experienced Rheumatologist and by a blinded skilled sonographer. ESAOTE MyLAB 70 with 6–18 MHz linear array transducer was used for all patients scanning and US were scored according with OMERACT guidelines.

Results: MSUS data of twenty RA patients (17 F – 3 M; median age of 59 yrs with range 35–83 yrs, median time for defined diagnosis of 2.4 months with range 1–11 months) and 15 SpA patients (9F–6M; median age of 53 yrs with range 18–78 yrs, median time for defined diagnosis of 1.7 months with range 1–8 months) were retrospectively analyzed. At the first EA clinic visit, all patients showed at least one joint with synovial fluid and/or synovial hypertrophy in Grey scale (GS) with a score >1 and no statistically significant differences were found in the percentage of patients that presented at least one joint with power Doppler (PD) positivity (55% RA Vs 53% SpA, $p=0.92$) and that presented PD positivity tenosynovitis of the flexor tendons in at least one finger (10% RA Vs 33% SpA, $p=0.08$). A statistically significant difference was found in the percentage of patients with erosions in at least one MCP (25% RA Vs 0% SpA, $p=0.036$) and in the percentage of patients with PD positive paratenonitis of the extensor tendons in at least one finger (30% RA Vs 80% SpA pts, $p=0.003$).

Conclusions: Early RA patients showed a statistically higher percentage of erosions at MCPs' MSUS evaluation of the first visit of the EA clinic, when compared with early SpA group and early SpA patients presented at the level of the extensor tendons a higher percentage of PD positive paratenonitis when compared with the early RA group.

References:

- [1] Lage-Hansen PR. et al. The role of ultrasound in diagnosing rheumatoid arthritis, what do we know? An updated review. *Rheumatol Int.* 2017Feb;37(2):179–187.
- [2] Gutierrez M. et al. Differential diagnosis between rheumatoid arthritis and psoriatic arthritis: the value of ultrasound findings at metacarpophalangeal joints level. *Ann Rheum Dis.* 2011 Jun;70.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2017-eular.5658

SAT0621 AGREEMENT BETWEEN SEMIQUANTITATIVE AND QUANTITATIVE DOPPLER SCORING SYSTEMS FOR THE ASSESSMENT OF SYNOVIAL PATHOLOGICAL VASCULARIZATION IN RHEUMATOID ARTHRITIS

E. Naredo^{1,2}, I.M. Rutigliano³, I. Janta¹, C. Scirocco³, L. Valor¹, F. Figus⁴, L. Carreño¹, J. Garrido⁵, A. Iagnocco⁶. ¹Department of Rheumatology, Hospital General Universitario Gregorio Marañón; ²Department of Rheumatology, Hospital Universitario Fundación Jiménez Díaz. Autónoma University, Madrid, Spain; ³Dipartimento Medicina Interna e Specialità Mediche, Sapienza Università di Roma, Rome; ⁴Dipartimento Reumatologia, Università degli Studi di Cagliari, Cagliari, Italy; ⁵Department of Social Psychology and Methodology, Faculty of Psychology, Autónoma University of Madrid, Madrid, Spain; ⁶Dipartimento Scienze Cliniche e Biologiche, Università degli Studi di Torino, Turin, Italy

Objectives: To compare power Doppler (PD) vs colour Doppler (CD) semiquantitative and quantitative scoring of synovial vascularization (RA) and to evaluate the relationship between semiquantitative and quantitative scores in patients with rheumatoid arthritis (RA).

Methods: One hundred RA patients underwent B-mode, PD, and CD assessments of 12 joints at two European centres. Each joint with synovial hypertrophy detected on B-mode was semiquantitatively scored (0–3) for PD (SPD score) and CD (SCD score) synovial signal. PD and CD synovial signal were also quantitatively scored (0–100%) (QPD and QCD scores, respectively) using a software for counting the colour fraction.

Results: We found SH in 184 joints. SPD and SCD agreed in 92.3% (95% CI: 88.4; 96.2%) of paired scores, with Kendall rank correlation coefficient tau-b=0.95. Significant differences between marginal distributions of SPD and SCD were not found ($p=0.565$). QPD and QCD scores were highly correlated (Pearson's coefficient=0.70) but Bland-Altman plot showed insufficient agreement, being the QCD scores systematically slightly higher than the QPD scores. The distribution of QPD and QCD values between SPD and SCD scores, respectively, showed significant differences between grade 0 and grade 1 ($p<0.001$), and grade 2 and grade 3 ($p=0.042$ and $p=0.007$, respectively) but not between grade 1 and 2 ($p=0.154$ and $p=0.150$, respectively).

Conclusions: SPD and SCD scores were concordant and QPD and QCD scores highly correlated although were not concordant. There was consistency between SPD and SCD moderate and severe scores and QPD and QCD scores. There was an overlapping between SPD and SCD mild and moderate scores regarding QPD and QCD scores.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2017-eular.3251

SAT0622 MULTICENTER 14-3-3ETA BIOMARKER REPRODUCIBILITY; THE JAPANESE EXPERIENCE

S. Hirata^{1,2}, M. Kaneda³, T. Isayama³, Y. Gui⁴, A. Marotta⁴, W.P. Maksymowych⁵, Y. Tanaka². ¹Hiroshima University Hospital, Hiroshima; ²University of Occupational and Environmental Health, Kitakyushu; ³Medical & Biological Laboratories, Nagano-ken, Japan; ⁴Augurex Life Sciences Corp, Vancouver; ⁵University of Alberta, Edmonton, Canada

Background: The soluble biomarker sub-committee of OMERACT has published validation criteria for biomarkers reflecting structural damage end-points. Within these, categories of Discrimination and Feasibility relate to assay reproducibility and performance. 14–3–3eta (η) is a joint-derived biomarker that drives joint damage processes and informs radiographic progression independently of acute phase reactants. As an ELISA assay, 14–3–3 η is currently available for clinical use as a laboratory developed test (LDT) in the United States (US) and as an in-vitro diagnostic (IVD) in Canada. The ELISA is (CE) marked for Europe and is Therapeutics Good Administration (TGA) approved for Australia. Previous studies have described 14–3–3eta assay equivalence at independent laboratories in the US and Canada.

Objectives: The purpose of this study was to demonstrate the reproducibility of 14–3–3eta measurements at two independent laboratories in Japan and one in Canada.

Methods: A total of 212 samples from the University of Occupational and Environmental Health were provided for this study. Serum 14–3–3eta measurements were performed using the 14–3–3eta ELISA provided by Augurex. Testing in Canada occurred in August 2015; these 14–3–3eta values were set as the "standard". The 212 samples were shipped from Canada to Japan in January 2016 with testing being performed in February 2016. Testing at MBL was performed on a blinded-basis. Upon completion of testing, results were sent to the investigators at which point in time, MBL was unblinded. 14–3–3eta positivity was defined at ≥ 0.19 ng/ml. Contingency and Spearman analyses were performed to assess the strength of the results between the two testing centres. Values above the linear range of the assay i.e. ≥ 20 ng/ml were excluded for the Spearman analyses and determination of the median (IQR). A p-value of <0.05 established statistical significance.

Results: In Canada, of the 212 samples tested, 146 (68.8%) were 14–3–3eta positive and in Japan, 147 (69.3%) were positive. In Canada, 187 patients had reportable values in the linear range and in Japan 186 did. Median 14–3–3eta levels in Canada were 0.51 ng/ml (IQR: 0.11–2.09) and in Japan they were 0.58 ng/ml (IQR: 0.02–2.68), respectively. Spearman correlation analysis revealed a highly significant correlation between the two testing sites, $r=0.92$; $p<0.00001$. Contingency analysis revealed a strong concordance between the two sets of results delivering a likelihood ratio (LR) of 156, $p<0.0001$. Between the two sites, 58 of the samples were 14–3–3eta negative and 139 were positive; there was agreement in 197 (93%) of the 212 samples.

Conclusions: As defined by the soluble biomarker sub-committee of OMERACT, reliability and standardization of biomarker assessments in routine clinical practice across the globe and in clinical studies is of high importance. The results presented herein demonstrate that the 14–3–3eta assay is highly reliable, that testing/delivery of results can be standardized between two independent laboratories, and that it is very stable over time and when shipped between continents.

Disclosure of Interest: S. Hirata: None declared, M. Kaneda Employee of: Medical & Biological Laboratories, T. Isayama Employee of: Medical & Biological Laboratories, Y. Gui Employee of: Augurex Life Sciences Corp., A. Marotta Employee of: Augurex Life Sciences Corp., W. Maksymowych Consultant for: Augurex Life Sciences Corp, Y. Tanaka: None declared

DOI: 10.1136/annrheumdis-2017-eular.5543

SAT0623 CORRELATIONS BETWEEN CLINICAL, ULTRASOUND AND DISEASE ACTIVITY SCORES OF PERIPHERAL ENTHESESITIS IN SPONDYLOARTHRITIS (SPA)

A. Haddouche, S. Haid, S. Bencheikh, S. Slimani, A. Abdessemed, N. Brahimi, A. Ladjouze. *Medicine, University of Algiers 2, Algiers, Algeria*

Objectives: To look for correlations between clinical, ultrasound (US) and disease activity scores of peripheral enthesitis in an SpA cohort.

Methods: A prospective study of all SpAs meeting SpA ASAS criteria followed at EHS Ben Aknoun, over a period from January 2015 to April 2016. Seventeen enthesal sites were assessed bilaterally: insertions of supra-spinatus, infra-spinatus, sub-scapular, medial and lateral epicondylars, triceps brachialis, gluteus medius and minimus, quadriceps, proximal and distal insertion (patellar ligament, medial and lateral collateral ligament), Achilles tendon and plantar aponeurosis. Peripheral entheses were assessed by the following clinical scores: Enthesitis Peripheral Score (PES= Sum of symptomatic peripheral entheses sites on clinical examination), Visual Analog Scale of peripheral enthesitis (VAS), Spondyloarthritis Research Consortium of Canada score (SPARCC) as well as the following US enthesitis scores: Acute Enthesitis score (Sum of acute enthesitis US scores for each site), Chronic Enthesitis score (Sum of US chronic enthesitis scores for each site), Global Enthesitis score (Sum of the acute and chronic US scores of enthesitis), Doppler signal Enthesitis score (Sum of Doppler signal scores less than 2mm from cortical bone for each site), Madrid Sonography Enthesitis Index (MASEI), Simplified Echographic Score (SES) which assesses only the Achilles

tendon and the plantar aponeurosis. Correlations between clinical, activity and US scores were investigated by SPSS software.

Results: A total of 208 patients were included. At examination 88.9% had an active disease and 64.4% of SpA were taking NSAIDs. 6240 entheses were assessed clinically and with US. A positive correlation was found between activity scores and two clinical scores (Peripheral Enthesitis Score, SPARCC). A good correlation was found between Activity scores and all US scores on one hand and with SPARCC scores and PES scores on the other hand. For US and clinical correlations, only Doppler scores was positively correlated with 2 clinical scores (SPARCC and PES scores), Acute Enthesitis scores was correlated with the PES scores. No correlation was found between clinical scores and MASEI or SES scores. No correlation was found with the mean enthesitis VAS scores.

Table 1. Correlations between clinical scores and activity scores with ultrasound scores

scores	PES	VASm Enthesitis	SPARCC	ASDAS-vs	ASDAS-crp
Acute Enthesitis	r: 0,14 p: 0,04	r: -0,03 p: 0,72	r: 0,16 p: 0,07	r: 0,43 p<0,001	r: 0,52 p<0,001
Chronic Enthesitis	r: 0,07 p: 0,30	r: -0,15 p: 0,03	r: 0,06 p: 0,41	r: 0,19 p: 0,006	r: 0,22 p: 0,002
Global Enthesitis	r: 0,15 p: 0,03	r: -0,08 p: 0,24	r: 0,13 p: 0,07	r: 0,43 p<0,001	r: 0,52 p<0,001
Doppler Enthesitis	r: 0,16 p: 0,02	r: -0,001 p: 0,99	r: 0,18 p: 0,01	r: 0,34 p<0,001	r: 0,36 p<0,001
MASEI	r: 0,09 p: 0,19	r: -0,08 p: 0,23	r: 0,12 p: 0,07	r: 0,44 p<0,001	r: 0,34 p<0,001
SES	r: 0,04 p: 0,55	r: -0,04 p: 0,59	r: 0,09 p: 0,18	r: 0,31 p<0,001	r: 0,38 p<0,001

Table 2. Correlations between clinical scores and activity scores

	PES	VASm Enthesitis	SPARCC
ASDAS-vs	r: 0,32 p<0,001	r: -0,04 p: 0,62	r: 0,27 p<0,001
ASDAS-crp	r: 0,31 p<0,001	r: -0,03 p: 0,62	r: 0,27 p<0,001

Conclusions: All US enthesitis scores were correlated with disease activity scores but those correlated with the clinical symptoms and not with its intensity were: Acute Enthesitis Scores, Global Enthesitis Scores and especially Doppler signal Enthesitis Scores.

Disclosure of Interest: None declared
DOI: 10.1136/annrheumdis-2017-eular.1968

SAT0624   QUANTITATIVE 3D IMAGING OF TENOSYNOVITIS AND BONE MARROW EDEMA BY DCE-MRI IS A SENSITIVE MEASURE OF RESPONSE TO THERAPY IN RHEUMATOID ARTHRITIS

C. Roberts¹, G. Guillard², M.A. Bowes², J. Burlison², A. Khan², N. Maguire¹, S. Gotla¹, A. Morgan¹, G.J. Parker^{1,3}, R. Hodgson^{3,4}, J. Freeston⁴, E.M. Vital⁴, P. Bird⁵, P. Emery⁴, P.G. Conaghan⁴. ¹Bioxydyn; ²Imorphics; ³Centre for Imaging Sciences, University of Manchester, Manchester; ⁴University of Leeds, Leeds, United Kingdom; ⁵Combined Rheumatology Practice, Sydney, Australia

Background: Quantitative analysis of tissue microvascular function using dynamic contrast-enhanced MRI (DCE-MRI) shows promise for improved understanding of synovial pathophysiology in rheumatoid arthritis (RA). Assessing tenosynovitis and bone marrow edema (BME) using physiologically-interpretable measures such as K^{trans} may offer additional insights; this will require increased precision in quantification.

Objectives: To apply a 3D DCE-MRI method to quantify capillary permeability (K^{trans}) in tendons, bone marrow edema, and to explore muscle involvement. To generate precise and consistent 3D regions of interest (ROI) of tenosynovitis and BME using active appearance models (AAMs)? and evaluate response to therapy within these regions.

Methods: MR images of the hand were acquired in 27 patients with established RA who had recently commenced the same biological therapy. Subjects were imaged at 0,3,6 months. The MRI protocol included pre- and post-contrast high-resolution 3D FLASH acquisitions. The DCE-MRI scan protocol included T1 mapping sequences followed by 15 sequential volumes acquired over 5 minutes, during which a bolus of gadolinium (0.1 mmol/kg) was administered at 2 ml/s at the beginning of the 5th volume measurement. The Extended Kety model was applied to each voxel concentration-time series within ROIs, allowing voxelwise estimates of K^{trans} . Pre-contrast T1-weighted images were searched using 3D active appearance models (AAMs) to reliably identify tendons and their sheath in the carpal tunnel region as well as bones and marrow area. Images from the dynamic series were registered to the high-resolution pre-contrast images, providing standardised 3D ROIs for each of the regions. Median K^{trans} was summarised in each patient for each ROI. T-test ($p<0.05$) determined significant differences from baseline.

Results: Differences in baseline K^{trans} were observed in each tissue type and demonstrated that highest levels of inflammation within the synovium, followed by tenosynovitis, BME and detection of low grade inflammation in the muscle (Fig. 1). Significant post-therapy responses in K^{trans} , indicating a reduction in perfusion and capillary permeability associated with reduced inflammation, were seen in each of these tissue types (Fig.1). RAMRIS scoring showed no significant

change at either 3 or 6 months. 3D visualisation of K^{trans} in tenosynovitis revealed additional spatial response to the biological treatment (Fig. 2) – most of the remaining tenosynovitis at 6 months had lower permeability with almost all of the higher permeability class of synovitis disappearing after therapy.

Figure 1: K^{trans} measured in each of 3D segmented regions of interest. Group mean and standard error shown.

Figure 2: K^{trans} maps overlaid on post-contrast T₁-weighted images from an example patient imaged at baseline (left), 3 months post-therapy (centre) and 6 months (right).

Conclusions: 3D DCE-MRI measures of tenosynovitis and bone marrow edema are practical in RA MR imaging trials, and offer sensitivity to change and differential tissue response not visible with other methods. Therapy-induced response in muscle suggests that there may be an inflammatory process in RA affecting local muscle groups.

Disclosure of Interest: None declared
DOI: 10.1136/annrheumdis-2017-eular.6584

SAT0625   ULTRASOUND ASSESSMENT OF JOINTS AND ENTHESIS IN A COHORT OF PATIENTS AFFECTED BY OCHRONOSIS: HOW COMMON IS INFLAMMATION?

C. Toscano¹, V. Di Sabatino¹, A. Adinolfi¹, M. Bardelli¹, V. Picerno¹, E. Selvi¹, A. Santucci², M. Galeazzi¹, B. Frediani¹, G. Filippou¹. ¹Department of Medicine, Surgery and Neuroscience; ²Department of Biotechnology, Chemistry and Pharmacy, University of Siena, Siena, Italy

Background: Ochronosis, the musculoskeletal manifestation of alcaptonuria (AKU), is characterized by alterations of the spine and large joints of the limbs similar to those of osteoarthritis. However, some cases of spinal involvement that resembles spondyloarthritis (SpA) have been describe, suggesting a prevalent inflammatory involvement of the joints.

Objectives: To evaluate the prevalence of inflammatory abnormalities in peripheral joints and entheses of a cohort of patients affected by AKU.

Methods: consecutive patients with definite diagnosis of AKU (with or without clinical manifestations) referred at our clinic from 2014 to 2016 were enrolled. All patients underwent a US examination of the following sites bilaterally: metacarpophalangeal joints (MCP), proximal interphalangeal joints (PIP), radiocarpal/mid carpal joints, elbow, gleno-humeral, hip, knee, ankle and metatarso-phalangeal (MTP) joints; flexor and extensor tendons of fingers and wrist and the ankle tendons. Further, the entheses of the rotator cuff of the shoulder, triceps, quadriceps, patellar and Achilles tendon were assessed. Joints and tendons with a synovial sheath were assessed for effusion, synovial hypertrophy and power Doppler (PD) signal while entheses were evaluated for the presence of PD signal, enthesophytes and calcifications. All the US lesions were scored using a dichotomous scale (presence/absence). All US exams were performed by an expert sonographer blind to clinical history, using an Esaote MyLab70 scanner equipped with high resolution linear probes.

Results: 11 patients (6 women) were enrolled in this study with a mean age of 57 yo (SD±11,50). the mean number of joints with effusion was 3,9 for each patient (median 3, range 2–8) while the mean number of joints with synovial hypertrophy was of 2,9 (median 2, range 2–7). =0,18 joints (median 0, range 0–2) presented also PD signal. 1 The mean number of exudative tenosynovitis was 0,81 (median 2, range 0–3) while proliferative tenosynovitis (mean 0,54, median 0, range 0–2) and PD in tendons with sheaths (mean 0,27, median 0, range 0–2) were rare. Finally, the mean number of entheses with PD was 1,27 (median 1, range 0–7), the mean number of enthesophytes was 0,63 (median 0, range 0–3) and for calcifications 4,27 (median 5, range 1–8).

Conclusions: Ochronotic arthropathy is believed to be characterized by a widespread articular damage, correlated mainly to degenerative processes due to the deposition of Homogentisinic Acid in the joints. The results of this US study