Results: Carotid intima media thickness was positively associated with 1-demographic characteristics of the participants such as age of the participants ($p < 0.001$), and age at RA symptoms onset ($p = 0.001$). 2-traditional cardiovascular risk factors such as systolic blood pressure ($p < 0.001$), diastolic blood pressure ($p = 0.016$), triglyceride level ($p = 0.016$), and low density lipoprotein (LDL) ($p = 0.001$). 3-inflammatory markers such as erythrocytes sedimentation rate (ESR) ($p = 0.020$) and c-reactive protein (CRP) ($p = 0.020$), and 4-renal function parameters such as uric acid level ($p = 0.006$), urine microalbumin level ($p = 0.030$). cIMT negatively associated with high density lipoprotein (HDL) ($p = 0.037$), 24 hours urine creatinine level ($p = 0.020$) and glomerular filtration rate ($p = 0.008$).

Conclusions: Subclinical renal function in conjunction with traditional and non-traditional cardiovascular risk factors work synergistically to accelerate atherosclerosis in RA population.

References:

- [1] Maradit-Kremser H, Crowson, Nicola PJ, Ballman KV, Roger VL, Jacobsen, Gabriel SE. Increased unrecognized coronary heart disease and sudden deaths in rheumatoid arthritis: a population-based cohort study. *Arthritis Rheum*, 2005;52:402–411.
- [2] Roman MJ, Moeller E, Davis A, Paget SA, Crow MK, Lockshin MD, Sammaritano L. Preclinical carotid atherosclerosis in patients with rheumatoid arthritis: prevalence and associated factors. *Ann Intern Med*, 2006; 144:249–256.
- [3] van Sijl AM, van den Oever IA, Peters MJ, et al. Subclinical renal dysfunction is independently associated with cardiovascular events in rheumatoid arthritis: the CARRE Study. *Ann Rheum Dis*. 2012;71(3):341–344.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2017-eular.1664

SAT0711 CHECKPOINT INHIBITOR THERAPY IN PATIENTS WITH ADVANCED MALIGNANCIES AND PREEXISTING RHEUMATOLOGIC DISEASE: THE MAYO CLINIC EXPERIENCE

U. Thanarajasingam¹, O. Pinkston², L. Kottschade³, H. Finnes³, S. Markovic³.
¹Rheumatology, Mayo Clinic, Rochester; ²Rheumatology, Mayo Clinic, Jacksonville; ³Oncology, Mayo Clinic, Rochester MN, Rochester, United States

Background: Immune checkpoint inhibitors (specifically the anti-CTLA-4 antibody ipilimumab, and the anti-PD-1 antibodies nivolumab and pembrolizumab, have revolutionized the treatment of advanced malignancies. However, by virtue of their mechanism of action – that is, loss of T-cell inhibition and impaired self-tolerance, patients with underlying autoimmune and rheumatologic diseases were typically excluded from the clinical trials leading to the approval of these agents. Limited data, typically of single checkpoint inhibitors, exist for their safety in patients with autoimmune diseases. However, the risk for rheumatologic disease flare in patients exposed to checkpoint inhibitors is unknown.

Objectives: To determine the risk of rheumatologic disease flare in patients receiving checkpoint inhibitor therapy.

Methods: We retrospectively studied all patients who had received a checkpoint inhibitor (i.e. ipilimumab, nivolumab, pembrolizumab, or any combination thereof) for any malignancy on the Mayo Clinic Rochester, Minnesota campus between January 1st, 2011 and May 16th, 2016 (approximately 5,200 patients.) Of these patients, we identified those with preexisting rheumatologic disease according to specific diagnostic codes.

Results: Of the 16 patients identified (13 [81%] female; median [range] age, 68.5 [34–86]y), 6 had inflammatory arthritis, 3 had polymyalgia rheumatica, 2 had lupus, 2 had Sjogren's syndrome, 1 had temporal arteritis, 1 had IBD-associated spondyloarthritis and 1 had gout. Patients were treated for the following metastatic/advanced malignancies: melanoma (9 [56%]), lung (5 [31%]) and lymphoma (2 [12%]) with: ipilimumab (4 [25%]), nivolumab (7 [44%]), pembrolizumab (5 [31.2%]) and ipilimumab/nivolumab (1 [6%]). Notably, in all cases, checkpoint inhibitor therapy was offered after failure of numerous other chemotherapies. Ten (62.5%) patients were on immunosuppressive therapy (mainly low dose prednisone and methotrexate) at the time of cancer diagnosis, and the majority (10 [62.5%]) were well-controlled/in remission from the standpoint of their rheumatologic disease. Three (19%) patients had flares of their rheumatologic disease after treatment for the following cancers as noted: Patient 1 – temporal arteritis – nivolumab – non-small cell lung cancer, Patient 2 – Sjogren's (severe sicca) – pembrolizumab – melanoma, and Patient 3 – spondylitis-ipilimumab/pembrolizumab combination – melanoma. All flares responded to steroids or supportive therapy. Checkpoint inhibitor therapy was discontinued in all three patients for the following reasons: Patient 1 – flare of temporal arteritis, Patients 2 & 3 – cancer progression.

Conclusions: To our knowledge, we have identified the largest cohort (16) of patients from a single academic with preexisting rheumatologic disease who had been exposed to checkpoint inhibitor therapy for advanced malignancy. Of these patients, only a minority experienced a flare of their disease during cancer treatment, and responded to standard therapies. With close monitoring and in the appropriate clinical context, checkpoint inhibitor therapies should be considered and made available to patients with preexisting rheumatologic disease who develop advanced malignancies.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2017-eular.5461

SAT0712 SELF-REPORTED SEDENTARY BEHAVIOUR IS ADVERSELY ASSOCIATED WITH MICROVASCULAR ENDOTHELIAL FUNCTION IN PATIENTS WITH RHEUMATOID ARTHRITIS

S.A.M. Fenton^{1,2}, J.J. Veldhuijzen van Zanten^{1,2}, A. Sandoo³, J.L. Duda¹, G.S. Metsios^{2,4}, G.D. Kitas². ¹School of Sport Exercise and Rehabilitation Sciences, University of Birmingham, Birmingham; ²Rheumatology, Russells Hall Hospital, Dudley; ³School of Sport, Health and Exercise Sciences, Bangor University, Gwynedd, Wales; ⁴Faculty of Education Health & Wellbeing, University of Wolverhampton, Wolverhampton, United Kingdom

Background: Patients with Rheumatoid Arthritis (RA) are at increased risk for cardiovascular disease (CVD). Research suggests impaired vascular function contributes to this heightened risk. At present, little is known regarding factors associated with vascular function in RA. Epidemiological evidence demonstrates sedentary behaviour (i.e., waking behaviour ≤ 1.5 metabolic equivalents whilst sitting or lying), to be adversely linked to CVD risk in the general population. Whilst the biological processes underlying this relationship are not understood, vascular dysfunction may play a role (1, 2). However, research is yet to examine the association between sedentary behaviour and vascular function in healthy adults and/or clinical populations. Studies investigating this relationship in RA, will help to determine the extent to which sedentary behaviour may represent a modifiable risk factor for CVD in these patients.

Objectives: To investigate the cross-sectional associations between sedentary behaviour and microvascular and large vessel endothelial function among patients with RA.

Methods: Fifty-three patients with RA participated in the study (M age=52.9 $\pm$ 12.8, 72% female). Laser Doppler imaging with iontophoresis was used to assess microvascular endothelium-dependent (acetylcholine, ACh) and endothelium-independent (sodium nitroprusside, SNP) function. Large vessel endothelium-dependent and endothelium-independent functions were measured via flow-mediated dilation (FMD) and glyceryl trinitrate dilation (GTN), respectively. Sedentary behaviour was self-reported via the International Physical Activity Questionnaire (hours/week sitting). Data were analysed using multiple linear regressions adjusted for traditional CVD risk factors; age, gender, total cholesterol, smoking status, family history of CVD, hypertension and body-mass-index.

Results: Sitting time (hours/week, $M = 39.2 \pm 17.9$) was significantly negatively related to % increase in perfusion in response to ACh ($\beta = -0.30$, $p < 0.05$) and SNP ($\beta = -0.37$, $p < 0.01$) after adjustment for traditional CVD risk factors. Sitting time accounted for 8% and 12% of the variance in microvascular endothelium-dependent function (ACh) and endothelium-independent function (SNP), respectively (traditional CVD risk factors, $R^2 = 0.3$). No significant associations were observed between self-reported sitting time and large vessel endothelium-dependent vasodilation (FMD, $\beta = 0.16$, $p = 0.29$) or independent vasodilation (GTN, $\beta = -0.08$, $p = 0.55$).

Conclusions: Sedentary behaviour appears to adversely affect microvascular endothelial function, but not large vessel function in patients with RA. It may therefore represent a modifiable risk factor for CVD in this population. Experimental studies employing objective measures of sedentary behaviour are necessary to confirm these findings, and to determine the utility of sedentary behaviour interventions for improving vascular function and reducing CVD risk in RA.

References:

- [1] Thosar SS, Bielko SL, Mather KJ, et al. Effect of Prolonged Sitting and Breaks in Sitting Time on Endothelial Function. *Med Sci Sports Exerc* 2015;47(4):843–9.
- [2] Fenton SAM & Kitas GD. Rheumatoid Arthritis: Sedentary behaviour in RA – a new research agenda. *Nat Rev Rheumatol* 2016;12(12):698–700.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2017-eular.4821

SAT0713 SURVEY ON PREVALENCE OF RHEUMATIC DISORDERS IN BANGLADESHI ADULTS

S. Ahmed¹, S.A. Haq¹, A.Z. Al-qadir¹, M.M. Rahman², S. Paul³.
¹Rheumatology, Bangabandhu Sheikh Mujib Medical University; ²Medicine, Dhaka Medical College, Dhaka; ³Medicine, Chittagong Medical College, Chittagong, Bangladesh

Background: From Community based COPCORD (community oriented program for control of rheumatic diseases) study in Bangladesh about a quarter of people is suffering from musculoskeletal disorders. But the study was conducted over a small number of samples and some specific localities of the country. So further study was required, covering the whole Bangladesh to justify this high prevalence.

Objectives: To determine the prevalence of musculoskeletal symptoms and specific rheumatic disorders in adult population of Bangladesh.

Methods: In this survey, a total of 2000 individuals aged 18 years or older were selected in twenty clusters (primary sample unit) from the seven divisions of the country. Modified COPCORD (Community Oriented Program for Control of Rheumatic Disorders) questionnaire was used to detect positive respondents. Standard criteria were used for diagnosing rheumatic disorders. Clinical judgment was used to solve diagnostic problems.

Results: In total 1843 individuals were interviewed with a response rate of 92.1%. The point prevalence of musculoskeletal pain was 33.7%. It was higher in women (38.7%) than men (28.4%) and higher in rural (34.5%) than that in urban (32.4%) areas. Higher prevalence rates were observed in homemakers (16.0%), laborers