root diameter/BSA ratio was significantly higher in HLA-B27 positive patients compared to HLA-B27 negative patients: 1.75 ± 0.22 mm vs. 1.61 ± 0.14 mm ($p=0.001$). Eight AS patients (6%) had aortic root dilatation (>2.1 mm corrected for BSA), who were all HLA-B27 positive. Patients with and without aortic root dilatation did not significantly differ in age or disease duration. The median aortic root diameter/BSA ratio was correlated with disease duration ($r=0.229$, $p=0.012$), but not with inflammatory biomarkers.

Conclusions: HLA-B27 positive AS patients with a long disease duration have an increased aortic root diameter and a higher risk for aortic root dilatation compared to HLA-B27 negative AS patients. The risk of developing aortic root dilatation in AS patients is comparable to patients with essential hypertension (6%)[2].

Future prospective studies should assess which AS patients phenotype are at the highest risk of this AS specific cardiac disease, thus candidates for regular echocardiographic screening.

References:

- [1] Klingberg et al. Aortic Regurgitation Is Common in AS: Time for Routine Echocardiography Evaluation? *Am J Med* 2015Nov;128(11):1244–1250.
- [2] Cuspidi C et al. Prevalence and correlates of aortic root dilatation in patients with essential hypertension. *J Hypertens* 2006 Mar;24(3):573–80.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2017-eular.4616

AB0117 CADHERIN-11 MRNA EXPRESSION IS INCREASED IN THE PERIPHERAL BLOOD OF PATIENTS WITH ACTIVE SPONDYLARTHROPATHY: A PILOT STUDY

N.I. Vlachogiannis, P.F. Christopoulos, G. Bamias, K. Fragiadaki, M. Tektonidou, P.P. Sfikakis. *Joint Academic Rheumatology Program, National and Kapodistrian University of Athens Medical School, Athens, Greece*

Background: Cadherin-11 is a key regulator of synovial architecture and has a central role in the formation of the rheumatoid pannus. Immunohistochemical studies have shown upregulation of cadherin-11 in the synovium of patients with seronegative spondylarthropathies (SpA), comparable to that in rheumatoid arthritis (RA), as well as in the intestinal tissue of patients with inflammatory bowel disease (IBD). Moreover, cadherin-11 mRNA transcripts have been identified in the peripheral blood of patients with RA and independently associated, among various disease characteristics, with the presence of active inflammation in multiple joints [1].

Objectives: To test the hypothesis that cadherin-11 mRNA transcripts are increased in the peripheral blood of patients with active SpA and search for possible associations with clinical features.

Methods: Fifteen patients with active SpA (BASDAI >4), aged between 21 and 71 years, 11 men, and 30 age- and gender-matched healthy controls were examined. Peripheral whole blood samples (3 ml) were subjected to cDNA synthesis and cadherin-11 mRNA expression was quantified by real-time PCR. Available cDNA from 33 IBD patients without axial or peripheral active arthritis served as disease control and were studied in parallel.

Results: Cadherin-11 mRNA was detected in the peripheral blood of 9/15 (60%) patients with SpA versus 5/30 (17%) healthy controls and 10/33 (30%) patients with IBD (SpA vs healthy controls $p: 0.006$, SpA vs IBD $p: 0.06$). Notably, cadherin-11 was not associated with BASDAI or skin/nail psoriasis present in 10 patients, but was detected with increased frequency among SpA patients with clinically active peripheral arthritis at the time of sampling (7 out of 10, 70%) than the remaining patients (2 out of 5, 40%). Moreover, cadherin-11 positivity associated significantly with increased erythrocyte sedimentation rate in SpA but not in IBD.

Conclusions: Cadherin-11 mRNA is upregulated in the peripheral blood of patients with SpA and may correlate with "spreading" of inflammation in peripheral joints. Since an anti-cadherin-11 mAb is in early clinical development for RA, further studies in patients with inflammatory arthritis are warranted.

References:

- [1] Sfikakis PP, Christopoulos PF, Vaiopoulos AG, et al. Cadherin-11 mRNA transcripts are frequently found in rheumatoid arthritis peripheral blood and correlate with established polyarthritis. *Clin Immunol* 2014;155:33–41. doi:10.1016/j.clim.2014.08.008.

Disclosure of Interest: N. Vlachogiannis: None declared, P. Christopoulos: None declared, G. Bamias: None declared, K. Fragiadaki: None declared, M. Tektonidou: None declared, P. Sfikakis Grant/research support from: Educational grant from Janssen

DOI: 10.1136/annrheumdis-2017-eular.3650

AB0118 THE LYMPHATIC SYSTEM: A GATEKEEPER FOR MIGRATION OF PATHOGENIC T-CELLS TOWARDS SYNOVIAL JOINTS AND ENTHESES IN PSORIASIS

R. Bisoendial¹, E. Prens², A. Mus³, P. Asmawidjaja³, N. Davelaar³, A. Hofman⁴, J.-B. Jaquet⁴, J. Hazes³, M. Kok⁵, E. Lubberts³. ¹Department of Rheumatology, Maasstad hospital and Erasmus Medical Center; ²Department of Dermatology; ³Department of Rheumatology, Erasmus University Medical Center; ⁴Department of Plastic surgery; ⁵Department of Rheumatology, Maasstad hospital, Rotterdam, Netherlands

Background: The factors underlying the transition of psoriasis (PsO) to psoriatic

arthritis (PsA) are poorly understood. The lymphatic system may control the homing of disease-associated T-cells to skin and extra-cutaneous sites like synovial joints and entheses.

Objectives: To study the capacity of lymphatic endothelial cells (LEC) to regulate T-cell homing capabilities in PsA.

Methods: Human dermal LEC (0.5×10^4), and fibroblast-like synoviocytes of a patient with PsA (PsA-FLS; 1.0×10^4) were pre-incubated for 3 days with PsA synovial fluid (PsA-SF; 0/10/20% v/v) and co-cultured with 2.5×10^4 allogeneic CD4+CD45RO+CD25- T-cells that were sorted from peripheral blood mononuclear cells of 3 donors (with or without stimulation with α CD3/ α CD28). After 72 h, T-cells were analyzed by flow cytometry. The CCR6+ T-helper (Th) subsets Th17.1 (CCR4-/CXCR3+), Th17/Th22 (CCR4+/CXCR3-), Th17 (CCR4+/CXCR3-/CCR10-) and Th22 (CCR4+/CXCR3-/CCR10+), and CCR6- subsets Th1 (CCR4-/CXCR3+), and Th2 (CCR4+/CXCR3-) were identified. We also looked at cutaneous lymphocyte-associated antigen (CLA). IL-17A, IL-22, and TNF protein levels were determined by ELISA. Statistical analysis included unpaired t-test (two-sided) for two-group comparison or one-way ANOVA with the Tukey-Kramer post hoc test for multi-group comparisons.

Results: Stimulation of CD4+CD45RO+ T-cells in co-culture with PsA-FLS skewed towards the CCR6+ subset Th17/Th22, which were predominantly Th17 cells. Th17 differentiation upon stimulation was suppressed in co-culture with LEC, even when LEC were pre-incubated with PsA-SF. T-cell stimulation in co-culture with LEC, as compared to PsA-FLS, promoted the generation of the Th22 subset. Upon co-culture with untreated LEC, stimulated T-cells showed higher expression of the skin homing receptor CLA than those that were co-cultured with PsA-FLS. The proportional reduction in CLA expression on T-cells in the co-cultures with LEC pre-incubated with PsA-SF 20% was comparable to PsA-FLS, however LEC conserved CLA expression on CD4+CD45RO T-cells at a higher level than PsA-FLS; this phenomenon particularly affected the CCR6+ T-cells. In line, a trend towards lower IL-17A and higher IL-22 levels were observed in the co-cultures with LEC that were pretreated with PsA-SF 20%, as compared to the co-culture with PsA-FLS. No differences were seen for TNF protein levels.

Conclusions: LECs are directly involved in T-cell differentiation and homing capabilities, as shown by suppression of Th17 differentiation upon stimulation, as compared to PsA-FLS. Also, LEC promoted Th22 generation, and conserved CLA expression in CCR6+ T-cells, even when LEC were preincubated with PsA-SF. Studies are underway to confirm that LECs from relevant biological tissues (e.g. synovium and lymph nodes) are critical for tissue-restricted T-cell migration to skin and synovial membranes in PsA.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2017-eular.5360

AB0119 ANTI-FIBROTICS TO PREVENT NEW BONE FORMATION IN SPONDYLOARTHRITIS: PROOF OF CONCEPT USING PIRFENIDONE IN CELL CULTURE MODELS

J. Laustsen^{1,2}, S. Lomholt³, P. Andersen³, J. Kelsen⁴, T.W. Kragstrup¹.

¹Department of Rheumatology, Aarhus University Hospital; ²Department of Biomedicine; ³Aarhus University; ⁴Aarhus University Hospital, Aarhus C, Denmark

Background: The pathogenesis of spondyloarthritis (SpA) involves both inflammation and new bone formation in the spine. In line with this, the disease has been characterized as both inflammatory and fibrotic. Current treatment including inhibitors of tumour necrosis factor alpha (TNF α) seem to dampen inflammation while new bone formation can progress. Therefore, there is an unmet therapeutic need for the treatment of new bone formation in SpA. Fibrosis is mediated by myofibroblasts and new bone formation is the result of increased osteoblast mineralization and decreased osteoclast bone degradation. Here, we evaluate the potential effect of the newly approved anti-fibrotic agent pirfenidone (Esbriet, Pirespa) on fibrosis and new bone formation in cell culture models of SpA.

Objectives: We hypothesized that pirfenidone inhibits SpA myofibroblast formation and activity and osteoblast mineralization.

Methods: Synovial fluid mononuclear cells from patients with SpA ($n=6$) were included for culturing fibroblast-like synovial cells (FLSs) while osteoblasts were purchased. The cells were cultured with pirfenidone in increasing concentrations (0.25, 0.5, and 1.0 mg/ml) with or without stimulation with tumor necrosis factor alpha (TNF α), transforming growth factor beta (TGF β), or interferon gamma (INF γ). The proliferation of FLSs was analyzed with light microscopy and flow cytometry using the marker Ki67. The differentiation and activation of FLSs was assessed with flow cytometry, a proteome profiler assay and enzyme-linked immunosorbent assays. The mineralization capacity of the osteoblasts was measured as deposition of hydroxyapatite.

Results: Pirfenidone reduced the Ki67 expression 7.1-fold in untreated FLSs ($p=0.001$) and 11.0-fold in FLSs stimulated with TGF β , TNF α , and INF γ ($p=0.022$). PFD further inhibited TGF β induced upregulation of α MA (Figure 2A) and INF γ induced upregulation of HLA-DR (Figure 2C) in all cultures. There was no difference between the percentage of ICAM-1 positive FLSs in cultures treated with or without pirfenidone. In supernatants from FLSs stimulated with TGF β , TNF α , and INF γ a total of 12 cytokines or chemokines had values above the detection limit in the membrane-based antibody array. Pirfenidone decreased the secretion of 3 of these 12 cytokines or chemokines more than 2-fold. The changes in secretion of monocyte chemoattractant protein 1 (MCP-1) and chitinase-3-like