MSCs transplantation. PBMCs from 8 patients were collected and co-cultured with UC MSCs at ratios of 1:1, 10:1 and 50:1, for 24 hours, 48 hours and 72 hours, respectively, to detect the levels of tolerogenic DCs. The FLT3L in the supernatant solution were determined. FLT3L siRNA was added to the co-culture system, and the level of tolerogenic DCs were detected.

Results: The levels of peripheral CD1c⁺ DCs and serum FLT3L were significantly decreased in SLE patients compared to healthy controls. After UC MSCs transplantation, the levels of CD1c⁺ DCs increased, along with an increase in serum FLT3L. In vitro studies showed that UC MSCs time-dependently up-regulated peripheral CD1c⁺ DCs, but not dose-dependently. The supernatant FLT3L level significantly increased after co-cultured with MSCs. However, the addition of FLT3L siRNA significantly abrogated the up-regulation of CD1c⁺ DCs by MSCs.

Conclusions: UC MSCs induce CD1c⁺ tolerogenic DCs through up-regulating FLT3L in lupus patients.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2017-eular.3334

Scleroderma, myositis and related syndromes - etiology, pathogenesis and animal models

AB0164 AMYLOIDOSIS IN PROGRESSIVE SYSTEMIC SCLEROSIS – A POSTMORTEM CLINICOPATHOLOGIC STUDY OF 12 PATIENTS

Á. Apáthy¹, M. Bély², ¹Department of Rheumatology, St. Margaret Clinic Budapest; ²Department of Pathology, Hospital of the Order of the Brothers of Saint John of God, Budapest, Hungary

Background: Systemic sclerosis (SSc), like all chronic autoimmune disorders, may be complicated by AA amyloidosis (AAa). An associated B-cell lymphoma my cause amyloid λ or κ light-chain deposition and AL amyloidosis (ALa) in SSc as well.

Objectives: The aim of this study was to determine the prevalence and type of amyloidosis in SSc patients, appraise the extent of amyloid deposits in various organs.

Methods: 12 patients (females 11, age: 54.82 years, range 66–32, onset of SSc: 48.86, disease duration: 6.43 years; one male, age: 65.0 years at death, onset of SSc and disease duration not known) were studied. All patients were autopsied. SSc was diagnosed clinically according to the criteria of the ACR [1].

Amyloid deposits on different tissue structures [arteriole, small artery, medium size artery, venule, small vein, medium size vein, interstitial collagen fiber, reticulin fiber (collagen IV), basal lamina, nerve, renal glomerulus] of 6 organs [heart, lungs, kidney, gastrointestinal tract, skin and brain] were determined histologically. The extent of amyloid A deposits was evaluated by semi-quantitative, visual estimation on a 0 to 3 plus scale, based on the number of involved tissue structures per light microscopic field [2] ("0": no amyloid deposits, "1": Sporadic, minimal amyloid deposits on different tissue structures, "2": less than five, "3": five or more involved tissue structures per microscopic field at objective magnification of x20)

The prevalence and extent of amyloid deposits in various organs were compared by Student (Welch) t-probe.

Results: Systemic AL-I light-chain amyloidosis was diagnosed in 1 (8.0%) 67 year old female patient (onset of SSc: 66 years, disease duration 1 year), and systemic AAa in 1 (8.0%) 53 year old female patient (onset of SSc: 41 years, disease duration 12 years).

The prevalence (in %) and the average extent of AL-I light-chain and amyloid A deposits (absolute value) in various organs of SSc patients are summarized in Table 1.

Table 1

Organs	SSc-λ Prevalence in %	SSc-AAa Prevalence in %	p<	SSc-λ Average extent	SSc-AAa Average extent	p<
Skin	80.00	0.00	0.0001	1.45	0.00	0.0002
Heart	72.73	63.64	0.3329	1.45	0.91	0.117
Kidney	63.64	63.64	0.1228	1.09	0.73	0.185
G-I Tract	41.67	66.67	0.1185	1.08	1.00	0.433
Lung	40.00	40.00	0.5000	1.00	0.50	0.077
Brain	0.00	0.00		0.00	0.00	–
Average/Organ	50.79	41.27	0.052	1.040	0.556	0.0036
Average/Patient	45.56	35.96	0.144	0.965	0.494	0.0006

Conclusions: AL-λ deposits were present earlier and were more prominent in the skin, heart and kidney, than in the, G-I tract and the lung. Amyloid A deposits were present earlier and were more prominent in the G-I tract, heart and kidney, than in the, lung and the skin. In the brain AL-λ and amyloid A deposits were absent in both diseases.

Higher prevalence (80.0% versus 0.0%; $p<0.0001$) and massive AL-λ deposition in the skin (1.45 versus 0.0; $p<0.0002$) may be explained by qualitative differences of AL-λ and amyloid A and by a diverse affinity of circulating amyloid precursors to the qualitative changed interstitial collagen and reticulin fibers. In systemic sclerosis patients qualitative change of collagens has been demonstrated [3–5].

References:

[1] van den Hoogen F et al: Ann Rheum Dis 2013; 72:1747–1755.
[2] Bély M, Apáthy Á: Clinical Pathology of rheumatoid arthritis. Akadémiai Kiadó, Budapest 2012 <http://www.akkrt.hu>.
[3] Istok R, et al. Annals Rheum Dis 1999; 58(Suppl1): 192.
[4] Istok R, et al: Reumatologia 2000; 2: 91.
[5] Istok R, et al: Exp Dermatol 2001; 26:545–547.
Disclosure of Interest: None declared
DOI: 10.1136/annrheumdis-2017-eular.1226

AB0165 CXCL4 MAY PLAY A KEY ROLE IN SYSTEMIC SCLEROSIS BY DRIVING CD4 T CELLS TO PRODUCE IL-17

A.J. Affandi^{1,2}, S.C. Silva Cardoso^{1,2}, S. Garcia-Perez^{1,2}, W. Marut^{1,2}, J.A.G. van Roon^{1,2}, T.R.D.J. Radstake^{1,2}. ¹Laboratory of Translational Immunology; ²Dept. of Rheumatology & Clinical Immunology, UMC Utrecht, Utrecht, Netherlands

Background: CXCL4 regulates multiple facets of immune response and its level is highly increased in various Th17-associated rheumatic diseases, including systemic sclerosis (SSc)^{1–4}. Recently, CXCL4 was shown to induce type I interferon production as well as endothelial activation in SSc patients¹. Th17 skewing has been demonstrated in SSc^{5–6}, however, whether CXCL4 plays a role in the induction of IL-17 production by human CD4 T cells is currently unclear.

Objectives: To investigate the effect of CXCL4 on human CD4 T cell phenotype in particular IL-17 production in the absence or presence of antigen presenting cells.

Methods: Blood was obtained from healthy donors and CD4 T cells, monocytes, dendritic cells, were isolated using magnetic-based sorting (n=20). In addition, CD4 T cells from SSc patients were isolated (n=10). CD4 T cells were activated using anti-CD3/CD28, or in co-cultures with antigen presenting cells, stimulated using superantigen Staphylococcal enterotoxin B. Exogenous recombinant human CXCL4 was added during (co-)culture in different concentrations. Cytokine production and proliferation were analyzed using Luminex immunoassays, intracellular cytokine staining, and flow cytometry.

Results: CXCL4 directly induced CD4 T cells secreting both IL-17 and IFN-γ, as well as IL-22, when costimulated with anti-CD3/CD28 ($p<0.05$). In many SSc patients, similar IL-17 increase upon CXCL4 treatment was observed, although this did not reach statistical significance. This might be due to the fact that CD4 T cells from SSc patients had already significant higher levels of IL-17 as compared to healthy donors (2182 ± 722.2 vs 1053 ± 263.6 pg/ml, mean±SEM, $p<0.05$). Furthermore, in co-culture system of CD4 T cells with monocytes or myeloid dendritic cells, CXCL4 treatment induced IL-17 production upon triggering by superantigen ($p<0.05$). Moreover, when monocyte-derived dendritic cells were differentiated in the presence of CXCL4, they orchestrated significantly increased levels of IL-17, IFN-γ, and proliferation by CD4 T cells (all $p<0.05$).

Conclusions: Altogether, we demonstrate that CXCL4 boosts pro-inflammatory cytokine production especially IL-17 by human CD4 T cells, either by acting directly or indirectly via antigen presenting cells. This indicates that targeting CXCL4 may potentially alleviate immune responses in Th17-mediated rheumatic diseases such as systemic sclerosis.

References:

[1] van Bon L, Affandi AJ, Broen J, et al. N Engl J Med 2014.
[2] Tamagawa-Mineoka R, et al. Allergol Int 2008.
[3] Vettori S, et al. Clin Rheumatol 2015.
[4] Yeo L, et al. Ann Rheum Dis 2015.
[5] Radstake TRDJ, van Bon L, et al. PLoS One 2009.
[6] Yang X, et al. Arthritis Res Ther 2014.

Acknowledgements: This study was supported by the Dutch Arthritis Association (Reumafonds) to A.J.A. and T.R.D.J.R., the Netherlands Organization for Scientific Research (NWO) to A.J.A. (Mosaic grant 017.008.014) and T.R.D.J.R. (Pre-Seed grant), the Portuguese Fundação para a Ciência e a Tecnologia to S.C.S.C. (SFRH/BD/89643/2012) and the European Research Council to T.R.D.J.R. (ERC Starting grant).

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2017-eular.6100

AB0166 INTERLEUKIN-6-POLARISED MACROPHAGES PROMOTE DERMAL MYOFIBROBLAST DIFFERENTIATION

A. Tam, X. Shi Wen, B. Ahmed-Abdi, C. Denton, D. Abraham, V. Ong. University College London, London, United Kingdom

Background: Macrophages and fibroblasts are key effector cell types present in scleroderma tissue¹. While the effect of scleroderma fibroblast conditioned medium on macrophages has been previously studied², less is known about the effect of scleroderma macrophages on fibroblasts. Interleukin-6 (IL-6) is an important mediator of fibrosis and is overexpressed in scleroderma sera and cells such as skin fibroblasts, monocytes and endothelial cells³. Given the increase in IL-6 levels in scleroderma and presence of the IL-6 receptor on the cell surface of macrophages, we are now investigating the phenotype of macrophages exposed to IL-6. Here we present our work into the paracrine function of IL-6-treated macrophages in stimulating fibroblasts using a media transfer approach.