of C-terminal telopeptide of type I collagen (CTX). Its place in the formation of osteoporosis in SLE patients is poorly understood, as well as its relationship with the course of the disease.

Objectives: The aim of study was to determine serum level of CTX in the SLE patients and its relationship with structural and functional state of bone tissue, course of the disease.

Methods: The study involved 58 SLE women (study group) and 29 healthy individuals (control group) representative by age and gender. The mean age of patients was 45.11 ± 1.03 years. For every patient data were recorded on age, body mass index (BMI), chronic SLE damage (SLICC/ACR DI) and disease activity score (SLEDAI), cumulative glucocorticoid dose, serum concentrations of interleukin-6 (IL-6) and C-reactive protein (CRP), bone resorption marker (CTX). Serum concentration of CTX was determined using ELISA test system "Nordic Bioscience Diagnostics A/S". Changes in BMD of the lumbar spine and proximal hip were determined by Dual-energy X-ray absorptiometry.

Results: It was established, that in patients with SLE serum level of C-terminal telopeptide of type I collagen was 1.15 ± 0.03 ng/ml, while in the control group – 0.83 ± 0.03 ng/ml, or was higher more than 19.0%. Violation of bone remodeling in patients with SLE was associated with reduced BMD, increased incidence of osteopenia and osteoporosis. Thus, the mean concentration of CTX in patients with osteoporosis was 1.63 ± 0.04 ng/ml, while in patients with normal bone – 1.0 ± 0.03 ng/ml. In SLE patients with osteopenia the level of C-terminal telopeptide of type I collagen was 1.23 ± 0.04 ng/ml (higher more than 23% compared with the control group). Increased CTX practically had no correlation with age, duration of the disease, smoking and BMI. At the same time the serum CTX was associated with chronic SLE damage index ($r = 0.51$), SLEDAI disease activity ($r = 0.41$), concentration of IL-6 ($r = 0.45$) and CRP ($r = 0.44$).

Conclusions: Alterations of bone metabolism were found in 19% female SLE patients in the form of increasing serum CTX and closely associated with the severity and activity of the disease, high levels of CRP and IL-6 and did not depend on the age, disease duration, smoking and body mass index.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2017-eular.5770

AB0515 SKIN MANIFESTATIONS AS INDEPENDENT PREDICTORS AND THE INITIAL RISK FACTORS FOR SYSTEMIC ANTIPHOSPHOLIPID EVENTS

L. Stojanovich, A. Djokovic, N. Stanisavljevic, M. Zdravkovic. *Internal Medicine Department, University Hospital Center Bezanijska Kosa, Belgrade, Serbia*

Background: Antiphospholipid syndrome (APS) patients express skin manifestations with the presence of various levels of antiphospholipid antibodies (aPL). Several studies have shown the frequency of dermatological manifestations with APS^{1,2,3}, including livedo reticularis, cutaneous ulcers, acrocyanosis, and other.

Objectives: Dermatological manifestations can be the initial clue in the diagnosis of this disease.

Methods: Our study includes a total of 508 APS patients; 360 were PAPS patients (283 female and 77 male, mean age 44.0 ± 12.9 years), 148 had APS associated with SLE/SAPS (133 female and 15 male, mean age 47.7 ± 14.8 years). aPL analysis included: LA, aCL (IgG/IgM), β_2 GPI (IgG/IgM). In all patients, we collected data considering frequently occurred skin lesions.

Results: Our results showed prevalence of skin manifestations in SAPS group of patinenst regading to PAPS (Table 1). Patients with skin manifestations overall had higher prevalence of thrombosis (Table 2).

Table 1. Prevalence of skin manifestations analyzed in PAPS and SAPS group

	PAPS (N=360)	SAPS (N=148)	p
Livedo	46 (12.8%)	86 (58.1%)	0.0001
Skin ulcerations	31 (8.6%)	47 (31.8%)	0.0001
Pseudovasculitis lesions	34 (9.4%)	72 (48.6%)	0.0001
Superficial cutaneous necrosis	9 (2.5%)	25 (16.9%)	0.0001
Digital gangrene	3 (0.8%)	14 (9.5%)	0.0001
Skin manifestations overall	98 (27.2%)	113 (76.4%)	0.0001

Table 2. Skin manifestations and thrombosis

	PAPS (p values)	SAPS (p values)
Livedo	0.038	0.062
Skin ulcerations	0.024	0.004
Pseudovasculitis lesions	0.007	0.019
Superficial cutaneous necrosis	0.005	0.380
Digital gangrene	1.000	0.008
Skin manifestations overall	0.0001	0.076

Conclusions: Dermatological manifestations can be very often the initial symptoms of severe manifestations of APS. Our study showed that patients with secondary APS had higher prevalence of skin lesions, and that some aPL types were risk factors for thrombotic manifestations in APS patients

References:

- [1] Toubi E, Krause I, Fraser A, Lev S, Stojanovich L, Rovinsky J, et al. Livedo reticularis is a marker for predicting multi-system thrombosis in antiphospholipid syndrome. *Clin Exp Rheumatol* 2005;23:499–504.

- [2] Weinstein S, Piette W. Cutaneous manifestations of antiphospholipid antibody syndrome. *Hematol Oncol Clin North Am* 2008;22:67–77.

- [3] Asherson RA, Cervera R, Klumb E, Stojanovich L, et al.: Amputation of Digits or Limbs in Patients with Antiphospholipid Syndrome. 21 January 2008. *Seminars in Arthritis and Rheumatism*. 2008; 38: 124–31.

- [4] Stojanovich L, Markovic O, Marisavljevic D et al. Influence of antiphospholipid antibody levels and type on thrombotic manifestations: results from the Serbian National cohort study. *Lupus* 2011;0: 1–8.

- [5] Stojanovich L, Kontic M, Djokovic A et al. Association between Systemic Non-criteria APS Manifestations and Antibody Type and Level: Results from the Serbian National Cohort Study. *Clinical and Experimental Rheumatology*, 2013,31(5):756–60.

Acknowledgements: This work was supported by research grant number 175041 for 2011 - 2017, issued by the Ministry of Science of the Republic of Serbia.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2017-eular.2759

AB0516 INCIDENCE OF CANCER IN A COHORT OF PATIENTS WITH PRIMARY SJÖGREN SYNDROME

M. Brom, S. Moyano, M. Scolnik, L.E. Pompermayer, I.J. Gandino, E.R. Soriano. *Rheumatology, Hospital Italiano de Buenos Aires, Ciudad Autónoma de Buenos Aires, Argentina*

Background: The most severe complication of Sjögren Syndrome is the development of lymphoproliferative processes. Several neoplasia have been associated with the disease, being non-Hodgkin lymphoma the most frequent one.

Objectives: Our objective was to evaluate incidence of cancer in a cohort of patients with primary Sjögren Syndrome.

Methods: A retrospective descriptive study was performed in a university hospital with its own health insurance and captive population. Using electronic medical records and laboratory database were review the entries performed between 01/01/2000 and 12/31/2015. We analyzed those patients with either diagnosis of Sjögren Syndrome, complain of dry mouth/eyes, or positive antibodies anti-Ro/SSA and anti-LA/SSB.

Among these patients, we included those fulfilling either ACR 2012 or EULAR 2002 Sjögren criteria, or those who were diagnosed as Primary Sjögren Syndrome by the treating rheumatologist even if they did not fulfill criteria.

We then proceeded to register and analyze demographic, clinical and histopathologic information available on their clinical records.

Results: One hundred fifty-seven patients with Primary Sjögren Syndrome were identified. Female accounted for 95.5% of the cohort; mean age at diagnosis was 49.4 years (SD 19). Median follow-up time was 7.7 years (IQR 8). The development rate and type of neoplasia was the following:

- Lymphomas: Three (Two MALT lymphomas of the parotid and one disseminated non-Hodgkin lymphoma). Density of Incidence 260/100,000 person/year (CI 95%: 50 – 750/100,000 person/year)
- Multiple Myeloma: One
- Skin (non-melanoma) neoplasia: Four
- Solid organ Neoplasia: Seven (Four breast cancer, one lung cancer, one uterus cancer, one tongue cancer). Density of Incidence 600/100,000 person/year (CI 95% 240 – 1240/100,000 person/year)

Univariate analysis showed association between lymphoma and cryoglobulinemia ($p=0.01$; OR=5.8), low C4 fraction of complement ($p=0.01$; OR=5.1), anemia ($p=0.02$; OR=1.96) and leucopenia ($p=0.03$; OR=1.67)

Conclusions: Development of cancer is a known complication of Primary Sjögren Syndrome. The association between lymphoma and cryoglobulinemia, low C4 fraction of complement, anemia and leucopenia enhances the importance of periodic screening for neoplasms among this subgroup of patients with Primary Sjögren Syndrome.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2017-eular.3195

AB0517 PREDICTIVE FACTORS FOR INFECTION IN SYSTEMIC LUPUS ERYTHEMATOSUS

M. Kechida¹, N. Lorenzo², R. Klii¹, S. Hammami¹, I. Khohtali¹. ¹Internal Medicine and Endocrinology Department, Fattouma Bourguiba Hospital, Monastir, Tunisia; ²Internal Medicine and cancerology Department, Saint-Cyr Hospital, Lyon, France

Background: it's known that infection could complicate the course of systemic lupus erythematosus (SLE) because of the immune status or the long term steroids and immunosuppressors.

Objectives: This study was aiming at determining the prevalence of infectious complications during SLE and their predictive factors.

Methods: A retrospective bi-centric analyzes of 289 patients diagnosed as SLE between January 2004 and December 2016 according to the ARA criteria of 1997 was conducted. A descriptive analysis of infectious complications was first made, then a comparative study between patients with (group 1) and without (group 2) infectious complications was performed to detect predictive factors.

Results: Mean age was 84.6 ± 13 years (14–72 years) with a sex ratio F/M=6.

Cardiac involvement and lupus nephritis (LN) were developed in 20% of patients. About 13.75% of patients had neurological manifestations, 26.5% articular complications, 16% vascular involvement and 10% of them developed infectious complications. Eight percent of these infections were diagnosed concomitantly with the diagnosis of SLE and 92% of them after the diagnosis of lupus with an average of 25 months. About 52.9% of the patients developed more than 2 episodes of infection. The spectrum of infectious complications was: pulmonary in 33.3%, urinary in 22.2% and cutaneous in 13.9%. Tuberculosis was the most frequent infection 12.5%. Lupus flare complicated the infection in 28.6% of patients with mean SLEDAI score at 10. Comparative study between group 1 and group 2 revealed that LN, corticosteroids and immunosuppressors were associated with a high risk of infection ($p=0.002$, $p=0.017$ and $p=0.034$ respectively). In multivariate analysis only LN was an independent predictive factor (OR=3.5, 95% CI=1.06-12.87, $p=0.049$).

Conclusions: Infections may complicate the course of SLE with flares presenting in 1/3 of cases. Half of the patients had more than 2 episodes of infection during their follow up. The presence of LN represents a predictive factor of such complication.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2017-eular.3009

AB0518 NEW 2016 ACR/EULAR CLASSIFICATION CRITERIA FOR SJÖGREN SYNDROME: USEFULNESS AND APPLICABILITY IN CLINICAL PRACTICE

M. Retuerto Guerrero¹, L. Sierra Herranz², C. Moriano Morales¹, C. Iñiguez Ubiaga¹, M. Garijo Bufort¹, A. Crespo Golmar¹, C. Álvarez Castro¹, E. Díez Álvarez¹, A. López Robles¹, M. Martín Martínez¹, T. Perez Sandoval¹.
¹Rheumatology Department; ²Pneumology Department, University Health Care Complex of León, Leon, Spain

Background: The Sjögren syndrome (SS) is an autoimmune disease where the cellular and humoral mechanisms affect the exocrine glands. In 2016, new classification criteria validated by ACR and EULAR were established.

Objectives: To compare the new criteria with those used so far in our hospital, as well as to assess the need for changes in the current diagnostic strategy.

Methods: Retrospective observational study in which 65 patients diagnosed with SS at the Hospital of León were randomly included. We reviewed the diagnostic tests performed and the fulfilment of the different classification criteria developed since 1993. Other variables studied were: sex; age at the time of diagnosis and the months from the onset of symptoms; xerostomia and xerophthalmia; extraglandular involvement, ESSDAI; immunosuppression; Raynaud; lymphoma development; and analytical alterations.

Results: The mean age at the time of diagnosis was 54.9 years ± 14 [23–82], with an average of months from the onset of symptoms to the diagnosis of 10.2 $\pm$ 9.5 [0–36]. 90.8% were women. 87.7% presented xerostomia; and 91% showed xerophthalmia, being severe in 43.1%. 64.6% had extraglandular manifestations; being the most prevalent the joint manifestation (60%) and the cutaneous one (18.4%). Over the past year, 37% developed haematological alterations in the form of cytopenias, and 73% biological alterations. At the time of the study, 32.8% presented low activity, 38.5% moderate activity and 9.2% high activity, measured by ESSDAI; being higher in anti-Ro positive patients ($p=0.011$). There was no association between ESSDAI and other antibodies, Raynaud or severe ocular involvement. 10.8% required systemic immunosuppression (RTX 5, AZA 2) and 18.5% needed ocular immunosuppression (topical cyclosporine). Only one patient developed lymphoma.

A Schirmer's test (ST) was performed in 92.3% (positive in 89.2%), saving the Van Bijsterveld test for patients with severe ocular involvement. The Ocular Staining Score (OSS) was not performed in any patient.

The scintigraphy of the salivary glands was positive in 70.8% of the patients and was not performed in 21.5%. The parotid sialography was only performed in two patients and the study of the salivary flow was not stimulated in none of them. Regarding the autoimmunity, 80% presented positive antiRo; 61.5% antiLa; 89% ANA; 61.5% RF; 43% quadruple positivity.

Labial gland biopsy was performed only in 18.4%, with a positive result in 75%.

All patients met the 1993 European Criteria; 86.2% met the European-American criteria of 2002; and only 10.8% met the SICCA-ACR Criteria. The new criteria validated by ACR and EULAR were verified in 80%. Four patients who fulfilled the European criteria did not meet the new criteria, coinciding with those patients with negative ST, but positive scintigraphy.

Conclusions: In our hospital, the method for electing the xerostomia study was the salivary scintigraphy; therefore, we cannot establish direct comparisons with the new criteria.

The incorporation of non-stimulated salivary flow in our diagnostic strategy is necessary.

We should consider conducting a lip biopsy more systematically for histological confirmation since there are no validated diagnostic criteria.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2017-eular.5255

AB0519 LABORATORY ABNORMALITIES IN PATIENTS WITH PRIMARY SJÖGREN'S SYNDROME

M. Fernandez Castro¹, J.L. Andreu², C. Sanchez-Piedra³, V. Martínez Taboada⁴, A. Olive⁵, J. Rosas⁶ on behalf of SJOGRENSER group, part of the Spanish Society of Rheumatology Systemic Autoimmune Diseases Study Group (EASSER). ¹Rheumatology, Hospital Infanta Sofía; ²Rheumatology, Hospital Puerta de Hierro Majadahonda; ³Research unit, Spanish Society of Rheumatology; ⁴Rheumatology, Hospital Marqués de Valdecilla, Madrid; ⁵Rheumatology, Hospital Hospital Germans Trias i Pujol, Barcelona; ⁶Rheumatology, Hospital Hospital Marina Baixa, Alicante, Spain

Background: Primary Sjögren's syndrome (pSS) is a chronic systemic autoimmune disease often accompanied by analytical abnormalities. Altered levels in serum protein concentration, blood cell count and autoantibodies contribute to the broad spectrum of biological manifestations that characterize pSS.

Objectives: The objective of this study is to evaluate the presence of laboratory abnormalities in patients with pSS from the SjogrenSER registry.

Methods: We conducted a multicentre transversal study of a cohort of pSS patients fulfilling 2002 European/American criteria, from 33 Spanish rheumatology departments. Every patient was interviewed for data collection and signed an informed consent. Data were also collected by reviewing medical records. Local ethics committees approved the study. Variables were analysed by descriptive statistical methods, using means, medians and rates. Chi-square was used to establish the statistical associations. A $p<0.05$ was considered significant.

Results: Four hundred and thirty-seven patients were included. Ninety-five percent of them were women. The median age of the cohort was 58 years. AntiRo antibodies were present in 93.6% of patients and AntiLa antibodies in 67.3% of patients. All patients were ANA+. Rheumatoid factor (RF) was positive in 64.8% of patients. Low levels of C3 and C4 were observed in 14.87% and 14.19% of patients, respectively. Polyclonal hypergammaglobulinemia (HGG) was present in 53% of patients. Thirteen patients had cryoglobulins (2.97%). An increase in β 2microglobulin was observed in 22.2% of patients. Fifty-six percent of the patients had hematological involvement: 29% of the patients had anemia, 38% had leukopenia (38% lymphopenia, 10.5% had neutropenia), and 9% had thrombocytopenia. The median ESR was 25 mm. Age at diagnosis and age at onset of symptoms were significantly lower in patients presenting RF+ vs RF- (48.71 vs 53.73, $p<0.001$ and 44.76 vs 49.53, $p=0.001$, respectively), decreased C3 vs normal C3 (45.66 vs 51.18, $p=0.004$ and 42.2 vs 46.99, $p=0.018$, respectively), decreased C4 vs normal C4 (47.02 vs. 50.89, $P=0.042$, for age at diagnosis) and HGG (47.59 vs 54, $p<0.001$, and 43.44 vs 50.16, $p<0.001$, respectively). ESR was significantly higher in patients with hematological involvement (35.94 vs 26.24, $p<0.001$), RF+ (36.39 vs 22.91, $p<0.001$), decreased C3 (37.8 vs 30.53, $p=0.026$) and C4 (38.71 vs. 30.42, $p=0.04$), HGG (36.21 vs 26.03, $p<0.001$) and increased β 2microglobulin (38.80 vs 29.71, $p=0.009$). ESSDAI (Eular Sjögren Syndrome Disease Activity Index) was significantly higher in patients with haematological involvement (5.58 vs 3.69, $p<0.001$), RF+ (5.40 vs 3.53, $p<0.001$) and HGG (5.31 vs 3.93, $P=0.011$). The median ESSDAI score was 2 (P25-P75, 0–4).

Conclusions: In SjogrenSER registry all patients were serologically positive. More than half of the patients presented abnormalities in serum proteins and 14% had hypocomplementemia. More than half of the patients had abnormalities, mostly leucopenia and lymphopenia. Patients with analytical alterations were younger at the time of diagnosis and had more often elevated ESR and higher ESSDAI score.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2017-eular.2603

AB0520 THE ROLE OF LEPTIN IN SJÖGREN'S DISEASE

M. Erdoğan¹, M.E. Tezcan², K. Başak³. ¹Internal medicine division of Rheumatology, Cerrahpaşa Medical Faculty; ²Internal medicine division of Rheumatology; ³Pathology, Dr. Lütfi Kırdar Kartal Training and Research Hospital, Istanbul, Turkey

Background: Sjögren's disease (SD) is a multisystemic disease mostly manifest with sicca symptoms. Lymphocytic infiltration of the glandular and extra glandular organs is the dominant pathologic feature of the disease. Multiple stimulators were accused in the pathogenesis of SD. Leptin, an endogenous peptide, involves in various metabolic processes as well as influence immune system (1). Increased serum leptin level is observed in patients with autoimmune diseases such as SD, systemic lupus erythematosus and rheumatoid arthritis when compared to healthy controls (2).

Objectives: Even if serum leptin level increases in the patient with SD, there is no data about its effect on exocrine glands. We aimed to compare density of leptin in the salivary gland of SD patients with control group. Furthermore we evaluated the relation between intensity of lymphocytic infiltration and density of leptin in the salivary glands of SD patients

Methods: We applied leptin immunostain to minor salivary glands samples of 24 SD patients, who were fulfilled American College of Rheumatology Sjögren's Disease Classification Criteria (ACR-SDCC) and 19 patients who undergo minor salivary gland biopsy due to clinically on suspicion of SD but not fulfilling the ACR-SDCC and had no lymphocytic focus on biopsy. Herein, leptin density in acinar and ductal structures of the salivary glands were evaluated in both groups. Moreover,