Diffuse scleroderma	6/11
Limited scleroderma	5/11
Pulmonary hypertension	4/11
History of digital ulcers	5/11
Time of endurance in minutes (ergospirometry)	5:32" (± 65 ")
Diffusion capacity (%pred)	82 $\pm$ 15
Pulmonary fibrosis	7/11
VE/VCO ₂ (mean values $\pm$ sd)	32 $\pm$ 5
Patterns of capillaroscopy	1/11 normal 1/11 early 4/11 active 5/11 late

References:

- [1] Ghizzoni C, Sebastiani M, Manfredi A, et al. Prevalence and evolution of scleroderma pattern at nailfold videocapillaroscopy in systemic sclerosis patients: Clinical and prognostic implications. *Microvasc Res.* 2015 May; 99: 92–5.
- [2] Castellví I, Simeón-Aznar CP, Sarmiento M, et al. Association Between Nailfold Capillaroscopy Findings and Pulmonary Function Tests in Patients with Systemic Sclerosis. *J Rheumatol.* 2015 Feb; 42(2): 222–7.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2017-eular.6599

AB0602 CAROTID ATHEROSCLEROTIC PLAQUES DETECTED BY ULTRASOUND IN SYSTEMIC SCLEROSIS PATIENTS UNDER 55 YEARS OLD

I. Sanz Pérez¹, F. Martínez Valle¹, A. Guillén del Castillo¹, N. Pizzi², R. Solans Laque¹, E. Callejas Moragas¹, I. González Nieto¹, V. Fonollosa Pla¹, C.P. Simeón Aznar¹. ¹Internal Medicine; ²Nuclear cardiology, Hospital Universitari Vall d'Hebron, Barcelona, Spain

Background: In Systemic sclerosis (SSc), as in other autoimmune diseases such as rheumatoid arthritis or systemic lupus erythematosus, cardiovascular events are one of the most frequent causes of mortality not attributed to the disease itself.

Objectives: The aim of this study was to assess the presence of subclinical atherosclerosis by carotid Doppler ultrasound in SSc patients under 55 years.

Methods: We conducted a cross-sectional study that included 78 SSc patients without cardiovascular events from H. Vall d'Hebron cohort (Barcelona). Carotid Doppler ultrasound was performed to measure the Carotid Intima Media Thickness (CMT) of common carotid artery (CCA) and detection of cholesterol plaques in CCA, bulb and internal and external carotid arteries, according to Mannheim consensus criteria. The results were compared to a healthy cohort from Barcelona, adjusted to age and sex¹. We used SCORE for populations with low risk and REGICOR as cardiovascular risk assessment charts.

Results: Risk factors and SSc related features are described in table 1. Twenty three patients (29.5%) had carotid plaques (CP) being the presence of CP statistically significant compared to the healthy cohort (29.5% vs 15.6%; $p < 0.05$; IC 0.04–0.24). None of the patients were catalogued as high risk according to SCORE chart. According to REGICOR chart, 4 patients (5.1%) were catalogued as intermediate risk and none as high risk. The four intermediate risk patients had CP. Carotid Intima Media Thickness mean (CMTm) and maximum (CMTmax), were statistically significant increased compared to the healthy cohort (CMTm 0.57 vs 0.53; $p < 0.05$ and CMTmax 0.74 vs 0.61; $p < 0.05$).

We performed multivariate regression analysis. Age, CMTmax, low High Density Lipid (HDL), the presence of pulmonary hypertension, and Diffuse cutaneous SSc patients were independent factors for the presence of CP.

Female	62 (79.5%)
Age	44 (20–55)
Disease duration	14 (0–38)
Limited cutaneous SSc	41 (57.7%)
Diffuse cutaneous SSc	17 (23.9%)
Sine Scleroderma SSc	8 (11.3%)
Initial Scleroderma	5 (7%)
Arterial Hypertension	18 (23.1%)
Diabetes mellitus	2 (2.8%)
Statin treatment	14 (17.9%)
Smokers	15 (19.2%)
Digital ulcers	30 (42.3%)
Pulmonary Hypertension*	4 (5.1%)
Interstitial lung disease	32 (41%)

*Mean pulmonary artery pressure > 25 mm Hg at right heart catheterization.

Conclusions: SSc patients under 55 years old often have subclinical atherosclerotic disease which could be detected by carotid ultrasound. Risk charts do not correlate with the presence of subclinical atherosclerotic disease. The presence of carotid plaque is more frequent in SSc patients with pulmonary hypertension, diffuse cutaneous subtype and patients with low HDL. CMT could be useful although it is currently not recommended in cardiovascular risk guidelines.

References:

- [1] Junyent M, Gilabert R, Núñez I, Corbella E, Vela M, Zambón D, et al. Carotid ultrasound in the assessment of preclinical atherosclerosis. Distribution of intima-media thickness values and plaque frequency in a Spanish community cohort. *Med Clin (Barc).* 2005;125(20):770–4.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2017-eular.6471

AB0603 QUANTITATIVE VIDEOCAPILAROSCOPY STUDY IN SYSTEMIC SCLEROSIS

A. Guillén-Del Castillo, C.P. Simeón-Aznar, E.L. Callejas-Moraga, S. Alonso-Vila, V. Fonollosa-Pla, A. Selva-O'Callaghan. *Department of Systemic Autoimmune Diseases, Hospital Universitari Vall d'Hebron, Universitat Autònoma de Barcelona, Barcelona, Spain*

Background: Systemic sclerosis (SSc) is an autoimmune disease characterized by immunological features, microvascular disturbances and fibrosis as a consequence of a massive production of extracellular matrix.

Objectives: The aim of this study was to analyse the alterations of the capillary nailfold bed and describe their correlation with clinical variables.

Methods: 134 patients were selected with SSc according to LeRoy and Medsger diagnostic criteria from the Hospital Universitari Vall d'Hebron cohort. Patients underwent an echocardiogram, pulmonary function test and nailfold videocapillaroscopy (NVC) within a 3 months period. NVC was performed from 2nd to 5th finger of both hands. NVC features were quantitatively measured. Informed consent was obtained by all the participants. SPSS 20.0 were used for statistical analysis. A P-value < 0.05 was considered as significant.

Results: Female was the predominant gender (113, 84.3%). The most common cutaneous subtypes were limited SSc, diffused SSc (88, 65.7% and 28, 20.9%, respectively). Almost 80% of patients met the ACR/EULAR 2013 classification criteria. The age ($\pm$ standard deviation) at NVC was 38.5 ($\pm$ 15.7) years, and the interval from the first SSc symptom and NVC was 16 ($\pm$ 12.6) years. Anti-centromere antibodies (ACA) were the most frequent in 36.6% of patients, followed by anti-topoisomerase I in 23.1%. Regarding organ involvement, 62 (46.3%) patients suffered from digital ulcers, 58 (43.3%) interstitial lung disease (ILD), 11 (8.2%) pulmonary hypertension, 110 (82.1%) gastrointestinal disturbances and 103 (76.9%) some cardiac alterations. The forced vital capacity was 80.8 ($\pm$ 20.1) % and diffusion capacity for carbon monoxide (DLCO) was 66.2 ($\pm$ 23.7) %. Regarding echocardiography results, the tricuspid regurgitation velocity (TRV) was 2.8 ($\pm$ 0.3) m/s and the right ventricular systolic pressure 30.7 ($\pm$ 12.0) mmHg. NVC findings were described as follows: the median of number of capillaries was 5.4 /mm, enlarged capillaries 0.8/mm, giant capillaries 0.2 /mm, microhemorrhages 0.1 /mm, ramified capillaries 0.3 /mm, tortuous capillaries 0.6 /mm and disorganized capillaries 0.0 /mm. SSc-ILD patients presented lower capillary density 4.8 /mm compared with 5.8 /mm in the other group ($P = 0.005$), and also more frequent ramified capillaries 0.3 /mm compared with 0.2 /mm ($P = 0.013$). With respect to the correlations, the number of capillaries was related with the DLCO ($\rho = 0.26$, $P = 0.003$) but negatively with the RVSP ($\rho = -0.21$, $P = 0.03$). Giant capillaries were correlated with LVEF ($\rho = 0.27$, $P = 0.001$) and greater tricuspid annular plane systolic excursion (TAPSE) ($\rho = 0.21$, $P = 0.01$). Microhemorrhages were associated with LVEF ($\rho = 0.29$, $P = 0.001$) although negatively with age at NVC ($\rho = -0.25$, $P = 0.003$) and time from first symptom ($\rho = -0.25$, $P = 0.003$). Disorganized capillaries were related with age at NVC ($\rho = 0.17$, $P = 0.04$) but negatively with DLCO ($\rho = -0.18$, $P = 0.035$).

Conclusions: The existence of a correlation between NVC features and clinical variables suggests that microvascular alterations may play a role as a pathogenic link with the cardiopulmonary SSc manifestations.

Acknowledgements: Guillén-Del Castillo A. is a researcher supported by the Contratos Predoctorales de Formación en Investigación (PFIS) grant from Instituto de Salud Carlos III [FI14/00643].

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2017-eular.4612

AB0604 RITUXIMAB EXPERIENCE IN PATIENTS WITH LONGSTANDING SYSTEMIC SCLEROSIS-ASSOCIATED INTERSTITIAL LUNG DISEASE: A SERIES OF 14 PATIENTS

A. Sari¹, D. Guven², B. Armagan¹, L. Kilic¹, A. Erden¹, A. Akdogan¹.

¹Rheumatology; ²Internal Medicine, Hacettepe University School of Medicine, Ankara, Turkey

Background: Interstitial lung disease (ILD) is the leading cause of mortality in systemic sclerosis (SSc) patients. Treatment options are rather limited in SSc associated ILD (SSc-ILD).

Objectives: Objective of this study was to report the experience of RTX treatment in a series of patients with longstanding SSc-ILD in whom unsatisfactory response in lung functions was noted under conventional treatments.

Methods: We retrospectively reviewed charts of 197 SSc patients evaluated between April 2015 and November 2016. 14 patients who received rituximab (RTX) for SSc-ILD participated in this analysis. The severity of ILD based on PFTs was defined as follows; mild (FVC between 71% and 80% of predicted), moderate (FVC between 51% and 70% of predicted) and severe (FVC $\leq 50\%$). The extent of skin disease was clinically measured by using Modified Rodnan Skin Score (mRSS) tool. End of follow-up was considered as six months after the last RTX dose.

Results: Median (IQR, interquartile range) age was 53.2 (46.8–55.5) and median disease duration was 9.1 (5.1–13.6) years. Median FVC was 52.5 (41.5–64.0)