(median 12 (IQR 7.25–17) vs. 7.5 (5–9), $P < 0.001$), while the original MASEI did not differ significantly (Table 1). Confluent PD over a larger area was only seen in 8% of the established PsA patients. Structural damage on ultrasound was more pronounced in the patients compared to the healthy volunteers. Number of PD locations and PD score did not distinguish the two groups.

Table 1. Participant characteristics and sonographic enthesitis scores

	PsA patients (n=84)	Healthy Volunteers (n=25)
Male, n (%)	45 (54)	12 (48)
Age, mean (SD)	55 (11)	47 (6)*
Disease duration, median years (IQR)	8.0 (4.9–12.3)	
LEI, median (IQR)	0.5 (0–2)	0 (0–0)**
Ultrasound		
MASEI, median (IQR)	15.5 (11–22)	13 (9–18)
Modified MASEI, median (IQR)	12 (7.25–17)	7.5 (5–9)**
structural components	7 (3–10)	3 (1–6)**
inflammatory components	6 (3.5–8.5)	3.5 (2–5.5)
Power Doppler		
in any entheses, n (%)	74 (88)	22 (88)
score 3, n (%)	7 (8%)	0 (0%)
locations, median (IQR)	2 (1–3)	2 (1–3)
if positive, median (IQR)	1.5 (1.25–1.75)	1.5 (1.25–1.5)

PsA: Psoriatic Arthritis; SD: standard deviation, IQR: interquartile range; LEI: Leeds Enthesitis Index; MASEI: Madrid Sonographic Enthesitis Score; modified MASEI: MASEI with new PD scoring method (1: one spot of PD, 1.5: some spots of PD, 2: confluent signal, 3: severe signal) and without knee entheses thickness. Structural components: erosions, calcifications, structure. Inflammatory components: bursitis, thickness and PD signal. PD: Power Doppler. * $P < 0.01$, ** $P < 0.001$ (Wilcoxon rank sum test).

Conclusions: Inflammatory and structural changes of the entheses measured with ultrasound are common in both unselected PsA patients and healthy volunteers, but more pronounced in established PsA patients.

References:

- [1] Wervers K, Rasappu N, Vis M, Tchetverikov I, Kok MR, Gerards AH, et al. AB0733 Masei Shows Substantial Changes in The Entheses of Young Healthy Volunteers – Amending Its PD Score and Excluding Knee Entheses Thickness Provides Better Discrimination of Enthesitis in Psoriatic Arthritis Patients. *Ann Rheum Dis* 2016;75:1155.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2017-eular.3431

AB0770 CHANGE OF PSORIATIC ARTHRITIS IMPACT OF DISEASE (PSAID12) QUESTIONNAIRE RELATED TO CHANGE IN DISEASE ACTIVITY IN EARLY PSORIATIC ARTHRITIS

K. Wervers¹, J.J. Luime¹, I. Tchetverikov², A.H. Gerards³, M.R. Kok⁴, C.W.Y. Appels⁵, W. van der Graaff⁶, J.H.L.M. van Groenendaal⁷, L.-A. Korswagen⁸, J.J. Veris-van Dieren⁹, J.M.W. Hazes¹, M. Vis¹ on behalf of Cicero. ¹Rheumatology, Erasmus MC, Rotterdam; ²Rheumatology, Albert Schweitzer Hospital, Dordrecht; ³Rheumatology, Vlietland Hospital, Schiedam; ⁴Rheumatology, Maasstad Hospital, Rotterdam; ⁵Rheumatology, Amphia Hospital, Breda; ⁶Rheumatology, Rivas Hospital, Gorinchem; ⁷Rheumatology, Reumazorg Zuid West Nederland, Roosendaal; ⁸Rheumatology, Sint Franciscus Gasthuis, Rotterdam; ⁹Rheumatology, Reumazorg Zuid West Nederland, Goes, Netherlands

Background: The Psoriatic Arthritis Impact of Disease 12-item questionnaire (PSAID12) has been developed to measure impact of Psoriatic Arthritis (PsA) for purposes of monitoring and clinical management. Although validated in patients with longstanding disease, data on validity and sensitivity to change in early PsA is lacking.

Objectives: We aim to relate change in disease activity to change in PSAID12 score in early PsA and evaluate which PSAID domains are more likely to change.

Methods: Patients with a new diagnosis of PsA were included in the Dutch southwest Early Psoriatic Arthritis cohort (DEPAR). For this analysis, patients that have PSAID12 (range 0–10) and Composite Psoriatic Disease Activity Index (CPDAI, range 0–15) data at two consecutive visits (i.e. 3 months apart) within the first year were included. In case multiple periods per patients were available, the first time period was chosen. The change in PSAID is compared to the change in disease activity over this period, measured with the CPDAI using Spearman's correlation coefficient. Change in score on individual domains of the PSAID was analysed in subgroups of patients that perceived improvement in health and those that perceived worsening. The SF-36 question on self-perceived change in health was used to determine these subgroups.

Results: 143 unique patients had at least one period with two PSAID and CPDAI measurements (67 from baseline-3 months, 26 3–6 months, 25 6–9 months and 25 9–12 months). Mean age was 51 (SD 13.7) and 70 (49%) were male. The initial median PSAID was 3.35 (IQR 1.4–5.1) and the subsequent score was 2.25 (0.95–4.8 with a mean delta of 0.52 ($P < 0.01$)). Median first CPDAI score was 4 (2–7) and 3 (1–5) for the second with a mean delta of 0.45 ($P < 0.05$). The difference in PSAID score was significantly but moderately correlated with the difference in CPDAI (Spearman's rho 0.267, $P = 0.0013$). 58 patients (41%) report a better health status compared to 3 months ago. Figure 1 shows that patients with self-perceived improvement of health have the highest improvement in pain and only domains of skin problems and embarrassment/shame did not improve

significantly. Patients reporting worsening of health (n=29) only have significantly lower scores in fatigue, discomfort and social domains.

Figure 1. Mean absolute differences in PSAID12 domains in patients with self-perceived improvement of health (n=58, black) and patients reporting worsening of health (n=29, grey). Shown as mean value and 95%-Confidence Interval

Conclusions: Improvement in CPDAI disease activity is significantly but moderately associated with improvement in PSAID score, with the biggest improvement in the pain domain in patients with a self-reported improvement of health.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2017-eular.3817

AB0771 SAFETY AND EFFICACY OF APREMILAST THROUGH 104 WEEKS IN PATIENTS WITH MODERATE TO SEVERE PSORIASIS WHO CONTINUED ON APREMILAST OR SWITCHED FROM ETANERCEPT TREATMENT IN THE LIBERATE STUDY

K. Reich¹, M. Goodfield², L. Green³, K. Nogales⁴, R. Chen⁴, E. Levi⁴, R.G.B. Langley⁵. ¹Dermatologikum Hamburg, Hamburg, Germany; ²Leeds General Infirmary, Leeds, United Kingdom; ³George Washington University School of Medicine, Washington, DC; ⁴Celgene Corporation, Summit, United States; ⁵Dalhousie University, Halifax, Canada

Background: Many patients (pts) with chronic plaque psoriasis exhibit nail and scalp involvement that can markedly affect quality of life and be difficult to treat.

Objectives: The phase 3b LIBERATE (Evaluation in a Placebo-Controlled Study of Oral Apremilast and Etanercept in Plaque Psoriasis) study (NCT01690299) evaluated the efficacy and safety of apremilast or etanercept vs. placebo in biologic-naïve pts with moderate to severe plaque psoriasis. Efficacy assessments included effects on preexisting nail and scalp disease and skin lesions.

Methods: In this double-blind, double-dummy study, pts were randomized (1:1:1) to placebo (PBO), apremilast 30 mg BID (APR), or etanercept 50 mg QW (ETN) through Week 16; thereafter, all pts switched to or continued APR (PBO/APR, ETN/APR, APR/APR) through Week 104. The primary end point was achievement of a $\geq 75\%$ reduction from baseline in Psoriasis Area and Severity Index (PASI) score (PASI-75) at Week 16 with APR vs. PBO; the secondary end point was PASI-75 achievement at Week 16 with ETN vs. PBO. Physician assessments were also conducted for overall disease activity (static Physician's Global Assessment [sPGA]); scalp disease activity (Scalp Physician Global Assessment [ScPGA], limited to patients with score ≥ 3 at baseline, indicating moderate to very severe scalp disease); and nail disease (Nail Psoriasis Severity Index [NAPSI], limited to patients with active disease [NAPSI ≥ 1] in the target nail at baseline). Responses were assessed at Week 104 using the last-observation-carried-forward (LOCF) methodology.

Results: The APR extension phase (Weeks 16 to 104) included 226 pts (PBO/APR n=73; APR/APR n=74; ETN/APR n=79). At Week 16, PASI-75 scores vs. PBO were significant for both APR and ETN; long-term treatment with APR maintained both PASI-75 and sPGA 0 or 1 response levels (Table). Improvements were seen in nail and scalp disease at Week 16, and responses continued to improve with APR treatment over 104 weeks and in pts who switched from ETN to APR (Table). ScPGA 0 or 1 was achieved by 50.0% to 59.2% of pts across treatment arms, and mean percent improvement from baseline NAPSI score ranged from -48.1% to -51.1% (Table); the proportion of pts achieving NAPSI-50 response ranged from 48.6% to 65.2%. Adverse events (AEs) occurring in $\geq 5\%$ of pts during Weeks 0 to 16 were diarrhea, nausea, nasopharyngitis, upper respiratory tract infection, and headache; long-term assessment by exposure-adjusted incidence rates (EAIR)/100 pt-yrs showed no increase with longer-term APR exposure. No increase in EAIR/100 pt-yrs of serious AEs occurred during the APR extension phase (3.45 to 5.49, across groups) vs. Weeks 0 to 16 (PBO 0.0; ETN 7.91; APR 12.57). Changes in laboratory parameters were infrequent and transient; EAIR/100 pt-yrs remained low across groups through 104 weeks.

Conclusions: APR demonstrated efficacy through Week 104 in pts who continued APR and pts who switched from PBO or ETN to APR at Week 16. The AE profile remained consistent with prolonged APR exposure, and no new safety or tolerability issues were observed through Week 104 in pts with moderate to severe plaque psoriasis.

Table. LIBERATE Efficacy Outcomes

Pts Achieving Response, % (LOCF)	PBO n=84	APR n=83	ETN n=83
Week 16			
PASI-75	11.9	39.8*	48.2*
sPGA 0 or 1 [§]	3.6	21.7*	28.9*
ScPGA 0 or 1 [§]	25.9	44.4**	50.0†
NAPSI-50 ^{§§}	10.9	25.0***	48.0*
Percent change from BL in NAPSI ^{¶¶}	-10.1	-18.7	-37.7
Week 104			
PASI-75	50.7	45.9	51.9
sPGA 0 or 1 [§]	27.4	18.9	26.6
ScPGA 0 or 1 [§]	50.0	59.2	56.6
NAPSI-50 ^{§§}	48.6	60.4	65.2
Percent change from BL in NAPSI ^{¶¶}	-48.1	-48.2	-51.1

APR=apremilast 30 mg BID; ETN=etanercept 50 mg QW; LOCF=last observation carried forward.
NAPSI=Nail Psoriasis Severity Index; NAPSI-50=50% or greater reduction from baseline in NAPSI score; PASI=Psoriasis Area and Severity Index; PASI-75=75% or greater reduction from baseline in PASI score; PBO=placebo; ScPGA=Scalp Physician Global Assessment; sPGA=static Physician's Global Assessment.
*P<0.0001; †P=0.0005; **P<0.0021; ***P=0.0458; ††P=0.0083; †††P=0.0701 vs. PBO (LOCF). Italics indicate values are nominally significant due to hierarchical testing of study end points.
§With ≥2-point reduction from baseline.
§In pts with ScPGA ≥3 (moderate to very severe) at baseline (exploratory): PBO n=58; APR n=54; ETN n=54 at Week 16 and PBO/APR n=50; APR/APR n=49; ETN/APR n=53 at Week 104.
§§In pts with NAPSI ≥1 at baseline: PBO n=46; APR n=52; ETN n=50 at Week 16 and PBO/APR n=37; APR/APR n=48; ETN/APR n=46 at Week 104.
¶¶PBO n=42; APR n=50; ETN n=50 at Week 16; PBO/APR n=33; APR/APR n=48; ETN/APR n=45 at Week 104.

Disclosure of Interest: K. Reich Consultant for: AbbVie, Amgen, Biogen, Boehringer Ingelheim, Celgene Corporation, Centocor, Covagen, Eli Lilly, Forward Pharma, GlaxoSmithKline, Janssen-Cilag, LEO Pharma, Medac, Merck Sharp & Dohme, Novartis, Ocean Pharma, Pfizer (Wyeth), Regeneron, Takeda, UCB Pharma, and Xenoport, Speakers bureau: AbbVie, Amgen, Biogen, Boehringer Ingelheim, Celgene Corporation, Centocor, Covagen, Eli Lilly, Forward Pharma, GlaxoSmithKline, Janssen-Cilag, LEO Pharma, Medac, Merck Sharp & Dohme, Novartis, Ocean Pharma, Pfizer (Wyeth), Regeneron, Takeda, UCB Pharma, and Xenoport, M. Goodfield: None declared, L. Green Consultant for: AbbVie, Amgen, Celgene Corporation, LEO Pharma, Novartis, Pfizer, and Valeant, Speakers bureau: AbbVie, Amgen, Celgene Corporation, LEO Pharma, Novartis, Pfizer, and Valeant, K. Nograles Employee of: Celgene Corporation, R. Chen Employee of: Celgene Corporation, E. Levi Employee of: Celgene Corporation, R. Langley Speakers bureau: AbbVie, Amgen, Celgene Corporation, Eli Lilly, LEO Pharma, Merck, Novartis, and Pfizer

DOI: 10.1136/annrheumdis-2017-eular.5163

AB0772 SECUKINUMAB PROVIDES SUSTAINED IMPROVEMENT IN FUNCTION, QUALITY OF LIFE AND FATIGUE OVER 2 YEARS IN PATIENTS WHO ACHIEVED LOW DISEASE ACTIVITY RELATED TO PSORIATIC ARTHRITIS DISEASE ACTIVITY SCORE (PASDAS)

L.C. Coates^{1,2}, T.K. Kvien³, P. Nash⁴, L. Gossec⁵, V. Strand⁶, L. Pricop⁷, L. Rasouliyan⁸, K. Ding⁹, S. Jugl⁹, C. Gaillez⁹ on behalf of the FUTURE 2 study group. ¹University of Leeds; ²Leeds Teaching Hospitals NHS Trust, Leeds, United Kingdom; ³Diakonhjemmet Hospital, Oslo, Norway; ⁴University of Queensland, Brisbane, Australia; ⁵UPMC Université Paris 06, Paris, France; ⁶Stanford University School of Medicine, Palo Alto; ⁷Novartis Pharmaceuticals Corporation, East Hanover, United States; ⁸RTI Health Solutions, Barcelona, Spain; ⁹Novartis Pharma AG, Basel, Switzerland

Background: PASDAS is a composite index measuring disease activity in psoriatic arthritis (PsA), well correlated with HAQ and health related quality of life¹.

Objectives: To report the impact of secukinumab on individual core components of PASDAS and its relationship with function (HAQ-DI), health related quality of life (SF36-PCS and MCS, DLQI, PsAQoL) and fatigue in patients (pts) who achieved PASDAS low disease activity (LDA) vs. high disease activity (HDA) through Week (Wk) 104 using *post-hoc* analysis from FUTURE 2 trial.

Methods: 397 pts with active PsA were randomised to s.c. secukinumab (300, 150, or 75mg) or placebo in FUTURE 2 study.² PASDAS is derived from physician's global VAS, pts global VAS, SF-36 PCS, tender and swollen joints (TJC68 and SJC66), Leeds enthesitis count, dactylitis count and CRP level and has cut-points for HDA (≥5.4), LDA (<3.2) and remission (REM≤1.9).³ PASDAS was assessed at Wks 16, 52 and 104 and reported as observed using non-mutually exclusive categories at group level. Additionally, the SF-36 PCS, SF-36 MCS, HAQ-DI, FACIT-Fatigue, PsAQoL, and DLQI were assessed by PASDAS LDA and HDA at Wks 16, 52 and 104 using MMRM analyses.

Results: Pts presented similar baseline characteristics. Mean±SD score of each PASDAS component among pts reaching LDA and HDA at Wk 16 for each treatment group are shown in table and were similar at Wk 104. Secukinumab treated pts achieving PASDAS LDA had significantly greater improvements in function, physical and mental health quality of life and fatigue compared to HDA through Wk 104 (Figure).

Conclusions: In pts treated with secukinumab, the most improved individual components with PASDAS LDA were related to dactylitis, enthesitis, SF36-PCS, Physician global VAS and SJC at Wk 16 and Wk 104. PASDAS LDA was associated with better improvement in function, quality of life and fatigue than

Table 1. Scores of PASDAS components at Wk 16

Core components, Mean ± SD	Secukinumab/PBO (s.c.), mg	LDA 300mg: n=37; 150mg: n=34; PBO: n=14	HDA 300mg: n=12; 150mg: n=23; PBO: N=34
Physician/Pt Global VAS	300 150 PBO	7.3±7.2/14.6±10.3 8.3±8.7/13.2±10.3 11.4±9.8/16.7±13.4	43.2±10.2/65.7±21.6 55.7±16.2/67.8±14.9 56.7±12.8/64.8±15.6
SF-36 PCS	300 150 PBO	51.5±5.8 51.8±5.5 49.4±6.6	34.9±6.0 34.2±6.5 33.5±8.2
SJC 66/TJC 68	300 150 PBO	2.2±3.0/2.3±3.4 1.9±3.6/2.6±5.0 1.7±2.0/3.4±4.7	12.3±10.0/27.9±16.1 15.9±11.1/32.3±21.7 10.1±6.1/25.4±14.0
Leeds Enthesitis Score	300 150 PBO	0.3±0.8 0.1±0.3 0.2±0.6	3.0±2.5 3.0±2.5 2.4±1.8
Tender Dactylitis count	300 150 PBO	0.03±0.2 0 0.1±0.5	0.9±2.6 1.4±3.1 1.3±2.4
CRP (mg/L)	300 150 PBO	3.3±3.5 3.3±3.5 5.8±15.2	5.0±4.5 8.1±11.6 14.7±19.5

Figure: Mean Change in PRO scores by PASDAS states (LDA vs. HDA) at Wks 16, 52 and 104

*P < 0.0001; †P < 0.001; ††P < 0.01 versus HDA.
DLQI, Dermatology Life Quality Index; FACIT-Fatigue, Functional Assessment of Chronic Illness Therapy-Fatigue; HAQ-DI, Health Assessment Questionnaire Disability Index; HDA, high disease activity; LDA, low disease activity; LS, least squares; n, number of pts with measurements; N, number of pts in each group of the specified analysis set; PsAQoL, PsA-specific quality of life; SF-36 MCS, Short Form-36 Mental Component Summary; SF-36 PCS, Short Form-36 Physical Component Summary
Wk 16: n=16 (LDA) and n=19 (HDA); Wk 52: n=125 (LDA) and n=21 (HDA); Wk 104: n=126 (LDA) and n=13 (HDA).
Analysis data were pooled across treatment arms. Placebo group not reported at Wk 16.

HDA confirming the importance to reach stringent target in a PsA pts with multifaceted disease.

References:

- [1] Helliwell PS and Kavanaugh A. Arth Care and Res. 2014;66:749–56.
- [2] McInnes IB et al. Arthritis Rheumatol.2016;68 (suppl 10).
- [3] Coates LC & Helliwell PS. J Rheumatol. 2016;43:371–5.

Disclosure of Interest: L. Coates Grant/research support from: Abbvie, Janssen, Consultant for: Abbvie, BMS, Celgene, Pfizer, UCB, MSD, Sun Pharma, Novartis, Lilly, Janssen, T. Kvien Consultant for: bbVie, Biogen, BMS, Boehringer Ingelheim, Celltrion, Eli Lilly, Epirus, Janssen, Merck-Serono, MSD, Mundipharma, Novartis, Oktal, Orion Pharma, Hospira/Pfizer, Roche, Sandoz and UCB, Speakers bureau: bbVie, Biogen, BMS, Boehringer Ingelheim, Celltrion, Eli Lilly, Epirus, Janssen, Merck-Serono, MSD, Mundipharma, Novartis, Oktal, Orion Pharma, Hospira/Pfizer, Roche, Sandoz and UCB, P. Nash Grant/research support from: Novartis, Abbvie, Roche, Pfizer, BMS, Janssen, and Celgene, Consultant for: Novartis, Abbvie, Roche, Pfizer, BMS, Janssen, and Celgene, Speakers bureau: Novartis, Abbvie, Roche, Pfizer, BMS, Janssen, and Celgene, L. Gossec Grant/research support from: AbbVie, BMS, Celgene, Janssen, Novartis, MSD, Roche and UCB, V. Strand Consultant for: AbbVie, Amgen, BMS, Celgene, Celltrion, CORRONA, Genentech/Roche, GSK, Janssen, Lilly, Merck, Novartis, Pfizer, Regeneron, Sanofi, and UCB, L. Pricop Shareholder of: Novartis, Employee of: Novartis, L. Rasouliyan Consultant for: Novartis through employment at RTI Health Solutions, Employee of: RTI Health Solutions, K. Ding Shareholder of: Novartis, Employee of: Novartis, S. Jugl Employee of: Novartis, C. Gaillez Shareholder of: Novartis, BMS, Employee of: Novartis

DOI: 10.1136/annrheumdis-2017-eular.1482