AB0891 HIGH FREQUENCIES OF CVD RISK FACTORS, OBESITY, METABOLIC SYNDROME, AND VITAMIN DEFICIENCIES IN A DANISH COHORT OF GOUT PATIENTS

O. Slot. Copenhagen Center for Arthritis Research, Rheumatology, Copenhagen University Rigshospitalet Glostrup, Glostrup, Denmark

Background: Gout is associated with increased risk of Cardio-Vascular Disease and death.

Objectives: To measure the frequencies of a number of potentially modifiable CVD risk factors in gout patients.

Methods: Comorbidities and CVD risk factors were measured and registered in a cohort of 100 consecutive new crystal proven gout patients.

Results: 88 males (62.1±13.6 years) and 12 females (74.1±6.9 years) were included. See table for results.

Comorbidities	
Previous CVD (n=36/100)	36%
Hypertension (BT>140/90 mmHg) (n=80/100)	80%
Hypertension without antihypertensive treatment (n=22/100)	22%
Diabetes (n=28/100)	28%
Nephropathy (eGFR<60 ml/min) (n=29/99)	29%
One or more comorbidities	86%
Comorbidity risk factors	
BMI, mean±SD (n=95)	29.6±5.3
BMI ≥30 (n=44/95)	46%
Metabolic Syndrome (Int Diab Fed 2006) (n=68/95)	72%
P-High Density Lipoprotein <1.00 mmol/l (n=30/96)	32%
P-Low Density Lipoprotein >3.0 mmol/l (n=39/91)	43%
P-Triglyceride >2.00 mmol/l (n=48/96)	50%
Homocysteine >15 μmol/l (n=28/71)	40%
Homocysteine ≥20 μmol/l (n=17/71)	24%
Cobalamine <200 pmol/l (n=14/90)	16%
Folate ≤9 nmol/l (n=22/86)	26%
25 (OH)-vitamin D3 <50 nmol/l (n=45/94)	48%
25 (OH)-vitamin D3 <20 nmol/l (n=18/94)	19%

Conclusions: Nephropathy, obesity, and dyslipidemia are known to correlate to hyperuricemia and gout. Reciprocally hyperuricemia and gout may lead to development of comorbidities, i.e. hypertension, CVD, and nephropathy.

Almost half the patients in this study were obese emphasizing the correlation between hyperuricemia and gout and obesity. Weight reduction is beneficial for control of hyperuricemia and gout, and is an important lifestyle factor that may be addressed in the treatment of hyperuricemia.

Metabolic Syndrome is characterized by abdominal obesity and at least two of the following criteria: hypertension, diabetes, hypertriglyceridemia, and low blood levels of high-density-lipoprotein. Metabolic Syndrome is associated with a two-fold risk of CVD and was present in 72% of the gout patients.

High homocysteine levels are correlated to endothelial dysfunction and increased risk of CVD. In this study the majority of patients had levels of homocysteine in the upper normal range, and a quarter had hyper-homocysteinemia. Elevated homocysteine has previously been reported in gout correlated to impaired renal function, but hyper-homocysteinemia can also be induced by low blood levels of cobalamin, which was present in 16%, and low levels of folate, which was found in 26%. In addition, low folate has been associated with endothelial dysfunction and CVD independently of homocysteine.

Low levels of 25(OH)-vitamin D3 were observed in nearly half of the gout patients and deficiency were found in 19%. Obesity is known to be associated with low levels of vitamin D. Negative correlation between uric acid and Vitamin D has though been reported after correction for weight. Uric acid suppresses 1-alpha hydroxylase leading to low levels of 1,25(OH)2-vitamin D3 which is correlated to dyslipidemia and Metabolic Syndrome. Urate Lowering Therapy has been shown to counteract suppression of 1-alpha hydroxylase.

Due to the increased risk of CVD and death associated with gout and the high prevalence of comorbidities and CVD risk factors in gout patients - comprehensive evaluation, information, and treatment of reversible co-morbidities and risk factors is recommended to be a part of the clinical management of gout.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2017-eular.3701

AB0892 PROGRESSIVE CLINICAL BENEFIT IN CHRONIC REFRACTORY GOUT PATIENTS ACHIEVING A PERSISTENT URATE LOWERING EFFECT FROM PEGLOTICASE TREATMENT

P. Khanna¹, B. Mandell², R. Fleischmann³, J. Kay⁴, A. Yeo⁵, P. Lipsky⁶.
¹Rheumatology, University of Michigan, Ann Arbor, MI; ²Dept. of Medicine, Cleveland Clinic, Cleveland, OH; ³Metroplex Clinical Research Center, Dallas, TX; ⁴U Mass Memorial Medical Center, Worcester, MA; ⁵Horizon Pharma, Lake Forest, IL; ⁶AMPEL BioSolutions, LLC, Charlottesville, VA, United States

Background: Pegloticase is a recombinant uricase conjugated to polyethylene glycol approved for treatment of patients with chronic refractory gout. It profoundly decreases serum uric acid (UA) and resolves tophi, but the quality of the clinical benefit in patients achieving long-term UA lowering has not been assessed.

Objectives: To assess the clinical benefit of long-term UA-lowering with pegloticase in patients with gout.

Methods: This analysis used results from two randomized controlled trials (RCTs) of 6- months duration and the 2-year open-label extension (OLE) of these studies.^{1,2} Efficacy was assessed in responders to the approved treatment regimen (8 mg pegloticase every 2 weeks [q2w]) in the RCTs (i.e., patients with plasma UA <6.0 mg/dL for ≥80% of the assessments around the 3 and 6 month time periods.) Clinical assessments included serum UA, frequency of gout flares, Patient Global Assessment (PGA), tender and swollen joints (TJC and SJC), pain measured with a 100-mm visual analog scale (VAS), Health Assessment Questionnaire Disability Index, bodily pain and the Arthritis-Specific Health Index from the Medical Outcomes Study Short Form 36 item, and reduction of target tophi.

Results: 33 patients who responded to pegloticase in the RCTs were followed throughout the OLE. Of these, 20 received 8 mg pegloticase q2w and 13 q4w. Both groups maintained markedly decreased serum UA levels during the OLE. Results for patients who received pegloticase q2w indicated significant improvements between RCT baseline and the final OLE evaluation for serum UA (P<0.0001), PGA (P=0.02), TJC (P=0.04), SJC (P=0.01), and pain VAS (P=0.01); 61.5% of patients in this group had complete target tophus resolution (P<0.0001). Results for those treated q4w indicated significant improvements between RCT baseline and the final OLE evaluation for serum UA (P<0.001), PGA (P=0.004), TJC (P=0.004), SJC (P=0.012), and pain VAS (P=0.01); 100% of patients in this group had complete tophus resolution (P<0.001). After an initial increase in gout flares during the initial 3 months of the RCT, there was a persistent decrease throughout the OLE. Maximal target tophus reduction was observed after 13–25 weeks of the OLE. Two of 20 (10%) patients receiving pegloticase q2w and 4 of 13 (31%) treated q4w lost the persistent UA-lowering effect during the OLE.

Conclusions: There were significant sustained clinical benefits with long-term pegloticase treatment in patients with chronic refractory gout achieving a UA lowering effect during the first 6 months of therapy. Significant decreases in TJC, SJC, and pain were noted along with significant improvements in PGA, suppression of gout flares, and resolution of tophi. In most patients, maximal benefit was noted after 6–12 months of pegloticase therapy, with many patients meeting newly proposed criteria for gout remission.³

References:

[1] Sundy JS, et al. JAMA. 2011;306:711–720.

[2] Becker MA, et al. Ann Rheum Dis. 2013;72:1469–1474.

[3] de Lautour H, et al. Arthritis Care Res. 2016;68:667–672.

Disclosure of Interest: P. Khanna Grant/research support from: AstraZeneca, Consultant for: Horizon, Ironwood, Selecta Bio, B. Mandell Grant/research support from: Horizon, Consultant for: Horizon, Ironwood, R. Fleischmann Consultant for: Horizon, J. Kay: None declared, A. Yeo Consultant for: Horizon Pharma, P. Lipsky Consultant for: AstraZeneca, Celgene, EMD Serono, GSK, Horizon Pharma, Janssen, Medimmune, Pfizer, Roche, Sanofi, UCB

DOI: 10.1136/annrheumdis-2017-eular.4119

AB0893 DISEASE RELATED KNOWLEDGE IN GOUT PATIENTS AND THE RELATIONSHIP WITH ADHERENCE TO URATE-LOWERING THERAPY IN EAST CHINA

R. Yin¹, T. Fu¹, Q. Zhang¹, L. Zhang¹, L. Li², Z. Gu¹. ¹Department of Rheumatology, Affiliated Hospital of Nantong University; ²School of Nursing, Nantong University, Nantong, China

Background: Gout is a chronic rheumatic disease caused by deposition of monosodium urate crystals in and around the joint, with a reported prevalence of 1.1% in mainland China and 6.24% in Taiwan, making it the most common form of inflammatory arthritis. With the disease progression, gout can cause permanent joint destruction, bone erosion, and organ damage. Urate-lowering therapy (ULT) is necessary to lower and maintain serum urate (sUA) levels at a therapeutic target of <360 μmol/L as this is associated with fewer gout flares, reduction of tophus size, and depletion of urate crystal stores in synovial tissues, making gout the only chronic arthritis that can be "cured". Disease related knowledge of gout patients should be assessed before attempting to improve health education. To date, except one study from south China, all other published papers about gout knowledge and medication adherence are from other countries. To our knowledge, there is no survey from east China.

Objectives: The current study aimed to investigate knowledge related to gout and its risk factors, and the relationship with adherence to urate-lowering therapy in patients with gout in east China.

Methods: A cross-sectional study of 229 gout patients recruited from the Affiliated Hospital of Nantong University between August 2015 and November 2016 was conducted with two questionnaire, Gout Knowledge Questionnaire (GKQ) and Compliance Questionnaire on Rheumatology (CQR). Chi-square analysis, t-test, rank sum test, as well as logistic regression analysis were used to analyze data.

Results: 21.5% (49/228) of gout patients in east China had knowledge of gout, and 9.1% (13/143) adhered to ULT. Age, employment, income, alcohol use, family history, acute flares in preceding 1 year, and colchicine use were associated with awareness of gout-related knowledge, and age, income, alcohol use, and colchicine use were the predictors. Among patients with ULT, patients adherent to ULT tended to have gout-related knowledge, compared with non-adherent patients. Awareness of the cause of gout attack, flare prevention induced by ULT and comorbidity were correlated with medication adherence, and the cause of gout attack as well as flare prevention induced by ULT were predictors of adherence to ULT in gout patients in east China.

Conclusions: In this study, 78.5% of patients in east China didn't have gout-related knowledge. Patients' knowledge on gout is a significant independent determinant of adherence to ULT.

Acknowledgements: This study was supported by grants from the Cultivate Distinguished Young Scholars Project of Nantong University (2nd); the 2015 Graduate Innovation Project of Nantong University (YKC15075); and College graduate research and innovation of Jiangsu Province (KYZZ15-0353).

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2017-eular.3071

AB0894 TREATMENT ADHERENCE TO URATE-LOWERING THERAPY IN CHINESE GOUT PATIENTS

R. Yin¹, H. Cao¹, T. Fu¹, Q. Zhang¹, L. Zhang¹, L. Li², Z. Gu¹. ¹Department of Rheumatology, Affiliated Hospital of Nantong University; ²School of Nursing, Nantong University, Nantong, China

Background: Gout, which is characterised by deposition of monosodium urate monohydrate (MSU) in synovial fluid and other tissues, is the most common form of inflammatory arthritis in man and older women. In addition to recurrent acute arthritis, subcutaneous tophi and chronic painful arthritis, gout also affects morbidity and premature mortality. Previous studies have reported that effective ULT can decrease sUA levels enough to prevent further crystal formation, dissolve existing urate crystals and eliminate the causative agent, thus reducing the frequency of acute gout attacks and preventing urate nephropathy, uric acid nephrolithiasis, and the deposition of tophi: a common cause of progressive joint damage, deformity and functional impairment, making gout the only chronic arthritis that can be "cured". However, gout patients' adherence rate to ULT is low, ranged from 10 to 46%. Emerging data suggest that poor adherence to long-term ULT use may be an important contributor to the suboptimal outcomes seen in gout and nonadherence may be worse in gout than in any other chronic disease. This suggests that there is a strong need to make a further study on drug adherence and its risk factors.

Objectives: Non-adherence in gout patients using urate-lowering therapy (ULT) may lead to joint destruction and permanent disability. Purpose of this cross-section survey was to explore Chinese gout patients' adherence rates and investigate potential risk factors for medication non-adherence.

Methods: A total of 129 gout patients were recruited from the Affiliated Hospital of Nantong University from August 2015 to September 2016. Patients were asked to complete a standardized self-report questionnaires (Compliance Questionnaire on Rheumatology, Treatment Satisfaction Questionnaire for Medication version II, Health Assessment Questionnaire, Confidence in gout treatment questionnaire, Gout Knowledge Questionnaire, Patient Health Questionnaire-9, Generalized Anxiety Disorder-7, and 36-Item Short Form Health Survey). Data was analyzed by independent sample t-test, rank sum test, chi-square analysis as well as logistic regression modeling.

Results: Based on CQR, 9.6% of gout patients were adherent to ULT. Adherence was associated with HAQ, GKQ, treatment satisfaction for medication, confidence in gout treatment and MCS. Other demographic, clinical and psychological characteristics were not related to adherence. Logistic regression models identified HAQ, GKQ and MCS as predictors of medication non-adherence.

Conclusions: In the current study, 90.4% of gout patients didn't adhere to their ULT prescription. HAQ, GKQ, treatment satisfaction for medication, confidence in gout treatment and MCS were relevant to medication adherence, and HAQ, GKQ and MCS were independent predictors of medication non-adherence in patients with gout. These findings could help medical personnel develop useful interventions to improve gout patients' medication adherence and quality of life.

Acknowledgements: This study was supported by grants from the Cultivate Distinguished Young Scholars Project of Nantong University (2nd); the 2015 Graduate Innovation Project of Nantong University (YKC15075); and College graduate research and innovation of Jiangsu Province (KYZZ15-0353).

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2017-eular.3075

AB0895 OSTEOPOROSIS RISK FACTORS IN PATIENTS WITH CALCIUM PYROPHOSPHATE CRYSTAL DEPOSITION DISEASE (CPPD)

S. Vladimirov, M. Eliseev, O. Zhelyabina, A. Smirnov, S. Glukhova. V. A. Nasonova Research Institute of Rheumatology, Moscow, Russian Federation

Background: Osteoporosis (OP) risk factors (RF) in CPPD patients are not sufficiently studied, although some of these RFs may be more prevalent in CPPD pts than in general population [1]

Objectives: To identify the input of individual RFs into OP development in pts with CPPD

Methods: 64 patients with CPPD (35 males and 29 females), but without OP were included into open prospective study. CPPD was confirmed based on McCarty criteria (detection of calcium pyrophosphate crystals in the synovial fluid using polarized light microscopy and detection of chondrocalcinosis based on joint radiography or US examination). 40 pts of those had acute and/or chronic arthritis, and 24 pts had osteoarthritis with CPP crystal depositions. Mean age was 57.6±10.2 years, mean follow up – 4.85±0.96 years. Bone mass density (BMD) was measured by dual energy X-ray absorptiometry (DXA) at the

forearm, lumbar spine and total hip was performed in all pts at baseline. OP was diagnosed by BMD (T-criterion ≤-2.5). The following OP risk factors were evaluated in this study: sex, age >55 years for females and >65 years for males, smoking, alcohol, fractures in past medical history, fractures in parents, BMI ≤20 kg/m², BMI ≤25 kg/m², serum levels of calcium, magnesium, vitamin D, hypoparathyroidism (HPT), chronic kidney disease (GFR ≤60 mL/min), intake of diuretics and glucocorticosteroids (GCs), Erythrocyte sedimentation rate (ESR) >20 mm/h, C-reactive protein (CRP) >5.0 mg/L. Odds ratio (OR) (95% confidence interval, CI) was estimated for each risk factor and logistic regression analysis was performed. Statistical analysis was made using SPSS v. 11 package, p values of <0.05 were considered statistically significant.

Results: OP was identified in 22 (34%) out of 64 pts with CPPD (13 males and 9 females) by the end of the study. The following factors were associated with OP: age >55 years in females (odds ratio (OR) 5.0, 95% CI: 1.021–24.49; p=0.047), age >65 years in males (OR 3.9, 95% CI 1.3–11.5; p=0.014), HPT (OR 15.38, 95% CI 1.7–137.9; p=0.015), elevated ESR (OR 3.38, 95% CI 1.14–10.0; p=0.028) and CRP (OR 6.42, 95% CI 2.05–20.05; p=.001) (Fig.1).

Only hypoparathyroidism was identified by logistic regression analysis (sensitivity-71%, specificity -82%) as OP-associated risk factor (OR 14.24, 95% CI 1.05–194.05; p=0.046) (Table 1).

Table 1. Data from multiple logistic regression analysis

Study	OR	Lower	Upper	p
sex	1,215	0,259	5,709	0,805
BMI<25kg/m ²	1,483	0,325	6,774	0,611
Age>65	1,293	0,274	6,093	0,745
Diuretics intake	2,771	0,189	40,617	0,457
HPT	14,245	1,046	194,055	0,046
CKD	1,025	0,082	12,876	0,985
lower vit.D3	0,503	0,116	2,173	0,357
elevated ESR	3,830	0,764	19,214	0,101
elevated CRP	3,855	0,608	24,44	0,152

Fig 1. Risk factors for osteoporosis (forest plot).

Conclusions: Hypoparathyroidism is the key risk factor for OP in CPPD pts. Among other risk factors chronic inflammation (ESR and CRP levels) is of highest importance.

References:

[1] Ross P.D. Osteoporosis: frequency, consequences, and risk factors. Archives Internal Med.1996; 156:1399–141.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2017-eular.3755

AB0896 THE EFFECT OF SERUM URIC ACID LEVELS ON TOPHUS STATUS AND FLARES IN PATIENTS WITH GOUT: A SYSTEMATIC REVIEW

S. Mitchell¹, L. Manounah¹, H. Liedgens², E. Johannes³. ¹Decision Resources Group, Bicester, United Kingdom; ²Head Health Economics and Outcomes Research, Market Access, SBU Europe?; ³Head Market Access Europe, Grünenthal GmbH, Aachen, Germany

Background: Gout is a chronic, progressive, inflammatory disease characterised by elevated serum uric acid (sUA) levels. sUA levels above its saturation point result in the deposition of monosodium urate crystals, which lead to gout flares and tophi (1). Multiple studies support the use of sUA levels as a marker for clinical improvements; hence the therapeutic goal is to lower sUA levels (<6.0 mg/dL) to improve the symptoms of gout and reduce the risk of associated comorbidities (2,3).

Objectives: To conduct a systematic review to identify studies reporting the effect of sUA levels on the incidence of gout flares and tophus status in adult patients with gout/hyperuricaemia, with a focus on publications reporting a correlation between the parameters.