categorically. Furthermore, potentially severe lupus manifestations, such as hemolytic anemia are not scored in SLEDAI.

Objectives: To evaluate the performance of SLEDAI-2K to detect a clinically meaningful change in SLE disease activity.

Methods: Prospective cohort study of SLE patients followed at a tertiary care lupus clinic from January 2014 to December 2016. Consecutive patients fulfilling the ACR'97 and/or the SLICC'12 classification criteria were included. At each outpatient visit, disease activity from the last 30 days was scored in the Physician Global Assessment (PGA) (0–3 cm scale) and in SLEDAI-2k. The association between PGA and SLEDAI-2K at each visit was tested with Spearman's Correlation. A clinically meaningful change in SLE disease activity was defined as difference in PGA ≥ 0.3 cm at follow-up compared to the baseline visit. Performance of change in SLEDAI-2K was tested in two models: against worsening and improvement in PGA ≥ 0.3 cm from baseline using Receiver Operating Characteristic (ROC) curve analysis. Sensitivity, specificity, positive and negative predictive values (PPV, NPV) of SLEDAI-2K to change in PGA was calculated. Statistical significance was set at 0.05.

Results: We included 334 patients (87.1% female, mean age at baseline - 44.8 $\pm$ 14.5 years). At baseline, median PGA and SLEDAI-2k score was 0.2 points (range 0–2.5) and 2 points (range 0–19), respectively. Eighty-three patients (24.8%) had a PGA ≥ 0.4 points at baseline. During follow-up of 36 months, 2129 visits were performed. PGA and SLEDAI-2K scores presented a high correlation ($\rho=0.82$, $p<0.0001$) (fig. 1). Reductions in SLEDAI-2K presented in ROC analysis an area under curve (AUC) of 0.697 [95% CI (0.628–0.766), $p<0.0001$] for an improvement in PGA ≥ 0.3 . For a worsening of PGA ≥ 0.3 points, increase in SLEDAI-2K presented an AUC of 0.877 [95% CI (0.822–0.932), $p<0.0001$]. Estimated sensitivities, specificities, PPV and NPV are presented in table 1.

Table 1. Performance of Sledai-2K to detect a clinically meaningful change in PGA, using cut-offs of decrease and increase (for a clinical improvement and worsening, respectively) in SLEDAI-2K ≥ 1 and ≥ 4 points

	Δ SLEDAI-2K ≥ 1				Δ SLEDAI-2K ≥ 4			
	Sens.	Spec.	PPV	NPV	Sens.	Spec.	PPV	NPV
Improvement PGA ≥ 0.3	0.7	0.571	0.397	0.825	0.288	0.929	0.622	0.763
Worsening PGA ≥ 0.3	0.725	0.903	0.627	0.936	0.353	0.996	0.947	0.873

Sens: Sensitivity; Spec: Specificity; PPV: Positive predictive value; NPV: Non predictive value.

Fig. 1 Scatter diagram showing linear positive correlation between PGA and SLEDAI-2K scores (Spearman's correlation coefficient (ρ) = 0.82, $p<0.0001$).

Conclusions: SLEDAI-2K presents a limited performance in detecting a clinically meaningful change in SLE disease activity, failing to identify more than a quarter of cases with clinically meaningful improvement or worsening. There is a need to optimize SLE disease activity measures.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2017-eular.3895

THURSDAY, 15 JUNE 2017

Joint EULAR - EFIS session: tilting the balance: from disease to tolerance induction

OP0207 B CELL DEPLETION INCREASES REGULATORY T CELLS AND AMELIORATES SKIN AND LUNG FIBROSIS IN A BLEOMYCIN-INDUCED SYSTEMIC SCLEROSIS MODEL MOUSE

H. Numajiri, A. Yoshizaki, T. Fukasawa, S. Ebata, Y. Asano, S. Sato. *Department of Dermatology, The University of Tokyo Graduate School of Medicine, Hongo, Bunkyo-ku, Tokyo, Japan*

Background: B cells play a critical role in systemic autoimmunity and disease expression through various functions such as cytokine production and induction of other immune cell activation. Recently, some clinical studies have shown that the efficacy of B cell depletion therapy with rituximab, a chimeric monoclonal antibody against human CD20, in systemic sclerosis (SSc) patients. However, it still remains unclear why B cell depletion can be an effective treatment for SSc.

Objectives: The purpose of this study is to assess the role of B cell depletion in

SSc. We evaluated the skin and lung fibrosis of bleomycin (BLM)-induced SSc model mice treated with B cell depletion. Furthermore, we investigated the effect of B cell depletion on T cell cytokine profile.

Methods: To generate BLM-induced SSc model mice, 300 μ g of BLM was injected subcutaneously into the shaved backs of the C57BL/6 mice every other day for 4 weeks. Anti-mouse CD20 monoclonal antibodies, which can deplete mouse B cells, were also injected every 2 weeks from 2 weeks later starting BLM treatment. After 4 weeks of BLM treatment, skin and lung fibrosis were assessed histopathologically. T cells and B cells were isolated from spleen using magnetic cell sorting system. Purified T cells (5×10^5 cells) were cultured with B cells (5×10^5 cells) in the presence of phorbol 12-myristate 13-acetate, ionomycin, and anti-CD3/CD28 antibodies. Cytokine expressions in the fibrotic skin and lung were quantified by real-time polymerase chain reaction. Cytokine production of T cells and B cells were analyzed by flow cytometric analysis.

Results: Dermal thickness and lung fibrosis score increased in BLM-treated mice compared with phosphate buffer saline (PBS)-treated control mice. Frequencies of interleukin (IL)-10 producing splenic B cells significantly decreased in BLM-treated mice compared with PBS-treated mice, while IL-6 producing B cell frequencies increased. Moreover, interferon (IFN)- γ , IL-4, or IL-17 producing T cell frequencies increased in BLM-treated mice. There were no significant differences in regulatory T cell frequencies between BLM-treated and PBS-treated mice. B cell depletion increased IL-10 producing regulatory T cell frequencies in BLM-treated mice. By contrast, frequencies of IFN- γ , IL-4, or IL-17 producing T cells were significantly decreased by B cell depletion in BLM-treated mice. In addition, fibrogenic cytokine mRNA expression levels of skin and lung decreased in BLM-treated mice with B cell depletion. To assess the role of B cells on T cell cytokine production, purified splenic B cells from BLM- or PBS-treated mice were cultured with naive T cells. T cells which were cultured with B cells from BLM-treated mice produced greater amounts of INF- γ , IL-4, and IL-17 than those cultured with PBS-treated mouse B cells. By contrast, B cells from PBS-treated mice induced a higher amount of IL-10 production from T cells than those from BLM-treated mice.

Conclusions: B cell depletion inhibited skin and lung fibrosis in BLM-treated mice. Furthermore, B cell depletion increased regulatory T cell frequencies in BLM-treated mice, though INF- γ , IL-4, and IL-17 producing T cell frequencies were decreased by B cell depletion. These results suggest that B cell depletion alters T cell cytokine profile, which results in inhibition of fibrosis in this model.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2017-eular.4189

OP0208 SYNOVIAL TISSUE OF RA PATIENTS IN REMISSION CONTAINS A UNIQUE POPULATION OF REGULATORY MACROPHAGES

A. Elmesmar¹, S. Alivernini², B. Tolusso², D. Vaughan¹, G. Di Sante², L. Petrica², E. Gremese², G. Ferraccioli², I.B. McInnes¹, M. Kurowska-Stolarska¹. ¹*Institute of Infection, Immunity and Inflammation, University of Glasgow, Glasgow, United Kingdom;* ²*Division of Rheumatology, Fondazione Policlinico Universitario A. Gemelli, Catholic University of the Sacred Heart, Rome, Italy*

Background: The majority of RA treatments target inflammation or the adaptive immune response. Partial- or non-response is common and only a minority have sustained remission. There is a knowledge gap in understanding the mechanisms that could reinstate synovial homeostasis in RA. Tissue macrophages may have a role in this process; they are present in healthy synovium and aid resolution of the inflammation in experimental models of RA. However, little is known about the regulatory properties of human synovial tissue macrophages.

Objectives: Our hypothesis is that healthy and RA synovium in remission contain macrophages with anti-inflammatory/repair properties and identifying the effector pathways that drive their function could facilitate therapeutic restoration of synovial homeostasis in RA.

Methods: We developed a flow cytometry sorting strategy for harvesting tissue-resident macrophages obtained from digested synovial biopsies of RA patients (n=21, including in remission n=5; and active RA n=16). Cells were labelled with cell lineage-specific antibodies; then macrophages were gated based on their expression of CD64^{pos}CD11b^{pos}MHCII^{pos}Lineage^{neg}. The potential homeostatic/repair macrophage was preliminary identified by the presence of CD206 marker. CD206^{pos} and CD206^{neg} macrophages were sorted using a FACS Aria III and RNAseq performed to characterise their functional signature. In some experiment, macrophages were seeded on collagen-coated plates and production of TNF α evaluated.

Results: All synovial tissue macrophages from RA in remission were CD206^{pos} whereas a substantial number of synovial macrophages from active RA tissue were CD206^{neg}. Gene expression analyses and functional assays suggest that these populations represent distinct phenotypes in the activation spectrum. CD206^{neg} macrophages have high expression of microRNA-155, which drives production of inflammatory mediators e.g. TNF α . In contrast, CD206^{pos} macrophages showed regulatory properties characterised by increased expression of soluble (e.g. IL10, TGFB), surface (e.g. IL4/14R, TGFB1/2) and cellular (e.g. SHIP1, TAM, SMAD2, STAT6) inhibitors of inflammatory activation, and increased expression of repair markers (e.g. ARG2 and CCL18).

Conclusions: We propose therefore that anti-inflammatory/repair macrophages may be present in human synovial tissues in remission representing a hitherto unnoticed regulatory tissue mechanism.

References:

- [1] Smolen JS, Aletaha D, McInnes IB. Rheumatoid Arthritis. *Lancet* Volume 388, Issue 10055, 22–28 October 2016, Pages 2023–2038.
- [2] Misharin AV et al. Non-classical Ly6C(-) monocytes drive the development of inflammatory arthritis in mice. *Cell Rep*. 2014;9:591.
- [3] Kurowska-Stolarska M, Alivernini S et al. MicroRNA-155 as a pro-inflammatory regulator in clinical and experimental arthritis. *PNAS* 2011;108:11193.
- [4] Elmesari A, McInnes IB, Kurowska-Stolarska M. MicroRNA-155 regulates monocyte chemokine and chemokine receptor expression in Rheumatoid Arthritis. *Rheumatology* (Oxford). 2016 Nov;55(11):2056–2065. Epub 2016 Jul. 13.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2017-eular.5910

THURSDAY, 15 JUNE 2017

What to do about comorbidity?

OP0209-HPR

INCIDENCE OF FIRST CARDIOVASCULAR EVENT IN SPANISH PATIENTS WITH CHRONIC INFLAMMATORY RHEUMATIC DISEASES: PROSPECTIVE DATA FROM THE CARMA PROJECT

M.A. Martin Martinez¹, S. Castañeda², C. Gonzalez-Juanatey³, F. Sanchez-Alonso¹, S. Marsal Barril⁴, M. Romero-Baures⁵, J. Santos-Rey⁶, J.A. Pinto-Tasende⁷, J. Tornero-Molina⁸, R. Lopez-Gonzalez⁹, M.D. Sanchez-Gonzalez¹⁰, J. Llorca¹¹, M.A. Gonzalez-Gay¹². ¹Research of Unit of Spanish Society of Rheumatology; ²Division of Rheumatology. Hospital Universitario la Princesa, IIS-Princesa, Madrid; ³Division of Cardiology. Hospital Lucus Augusti, LUGO; ⁴Division of Rheumatology. Hospital Universitario Vall d Hebron; ⁵Division of Rheumatology. Hospital Universitario Bellvitge, Barcelona; ⁶Division of Rheumatology. Hospital Virgen de la Salud, TOLEDO; ⁷Division of Rheumatology. Complejo Hospitalario a Coruña, la Coruña; ⁸Division of Rheumatology. Hospital Universitario de Guadalajara, Guadalajara; ⁹Division of Rheumatology. Hospital Virgen de la Concha, ZAMORA; ¹⁰Division of Rheumatology. Hospital Universitario de Salamanca, Salamanca; ¹¹Division of Epidemiology and Computational Biology, School of Medicine, University of Cantabria, Ciber Epidemiología y Salud Pública (Ciberesp); ¹²Division of Rheumatology, Hospital Universitario Marques de Valdecilla, and Epidemiology, Genetics and Atherosclerosis Research Group on Systemic Inflammatory Diseases, Rheumatology Division, IDIVAL, Santander, Spain

Objectives: To determine the incidence and risk factors implicated in the development of first cardiovascular event (CVE) in patients with chronic inflammatory rheumatic diseases (CIRD) attending rheumatology clinics after 2.5 years of follow-up.

Methods: Analysis of data after 2.5 years of follow-up in an observational prospective study [CARDiovascular in rheUMAtology (CARMA) project] that includes a cohort of patients with CIRD [rheumatoid arthritis (RA), ankylosing spondylitis (AS), and psoriatic arthritis (PsA)] and another cohort of matched individuals without CIRD attending outpatient rheumatology clinics from 67 hospitals in Spain. The cumulative incidence per 1000 patients and the incidence density per 1000 patient-months of non-fatal CVE were estimated in both cohorts at 2.5 years from the start of the project. Weibull proportional hazard model was used to calculate the Hazard Ratio (HR) and 95% confidence interval (95% CI) of the risk factors involved in the development of CVD events. Losses to follow-up and their causes were also analyzed.

Results: The total number patient who completed the follow-up visit at 2.5 years was 2,598 (89.2% of those who started the study). Seven patients had died due to CVE and 23 because of non-CVE. The higher number of losses to follow-up was found in the control group (15.81%), because many of them were not periodically follow-up at the outpatient clinics. Cardiovascular cumulative incidence in patients with CIRD 15.30 cases per 1000 patients (95% CI: 12.93–17.67), being higher in AS patients 22.03 (95% CI: 11.01–33.04). The higher risk of developing a first CVE during the 2.5 years of follow-up was in patients with AS (HR: 4.11, 95% CI: 1.07–15.79; p: 0.04), those with older age (HR:1.09; 95% CI: 1.05–1.13, p<0.001), higher systolic blood pressure (HR: 1.02; 95% CI: 1.00–1.04, p=0.01) and longer duration of the rheumatic disease (HR: 1.07; 95% CI: 1.03–1.12), p<0.01). In contrast, woman gender was a protective factor (HR: 0.43; 95% CI: 0.18–1.00, p=0.05).

Conclusions: Patients with AS prospectively followed-up at rheumatology outpatient clinics show higher risk of developing a first CVE than those with RA or PsA. Besides traditional CVD risk factors a longer time course of the disease is a risk factor for the development of CVD in patients with CIRD.

Acknowledgements: This project has been supported by an unrestricted grant from Abbvie, Spain. The design, analysis, interpretation of results and preparation of the manuscript has been done independently of Abbvie.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2017-eular.3838

THURSDAY, 15 JUNE 2017

Difficult to reach patient groups

OP0210-PARE

THE DEVELOPMENT AND EVALUATION OF AN INTERACTIVE HEALTH COMMUNICATION APPLICATION TO EDUCATE AND EMPOWER YOUNG PEOPLE LIVING WITH JUVENILE IDIOPATHIC ARTHRITIS: A PILOT STUDY

S.R. Stones^{1,2}, S.M. Cruickshank³, S.M.M. Verstappen⁴ on behalf of MyJIA Expert Advisory Group. ¹School of Healthcare, University of Leeds, Leeds; ²Faculty of Life Sciences; ³School of Biological Sciences; ⁴Arthritis Research UK Centre for Epidemiology, The University of Manchester, Manchester, United Kingdom

Background: In parallel with clinical care, it is equally important to empower young people with juvenile idiopathic arthritis (JIA) to develop self-efficacy skills, so that they can competently self-manage their health, particularly as they transition into adult health services.¹ Capitalising on the popularity of technology, web-enabled tools represent a novel and effective way of engaging young people with JIA. A Cochrane review of Internet Health Communication Applications (IHCA) found that IHCA have a positive effect on self-efficacy, empowering individuals to become more knowledgeable.²

Objectives: The aim of this study was to: i) develop and evaluate an IHCA in young people with JIA, aged 16 to 25 years; and ii) investigate whether an IHCA enhanced young people's condition-specific knowledge, and their self-efficacy to manage their health.

Methods: Young people aged 16 to 25 with a self-reported diagnosis of JIA were recruited to take part via social media. The IHCA was built using Microsoft PowerPoint and iSpring Suite (Figure 1). A virtual advisory group of patients, parents and healthcare professionals were involved throughout the design and development of the IHCA. The contents of the IHCA was obtained for existing published sources. Matching anonymous pre- and post-participation questionnaires were used to assess differences in knowledge of JIA and self-efficacy skills.

Results: In total, 23 (79%) young people with JIA completed both pre- and post-participation questionnaires. Most young people reported that their preferred source of condition-related information was obtained via an internet search engine (83%). Prior to completing the IHCA, only 22% of young people reported that they had an excellent understanding of JIA. After completing the IHCA, there was a significant improvement in young people's condition-specific understanding. Similarly, after completing the IHCA, there was a significant improvement in young people's confidence to better self-manage their health. Interestingly, only 30% of young people reported that the information which health professionals provided to them about JIA was adequate, in terms of comprehension and usefulness. Qualitative findings identified three core themes: resource content, practical support and accessibility/functionality. Young people liked the way that information was presented, as well as practical steps they could take to improve their health.

Conclusions: Quantitative results correspond with qualitative findings, indicating that the IHCA was well-received by young people with JIA. Condition-specific understanding was enhanced after completing the IHCA, as was participants' confidence in their self-efficacy and disease self-management capabilities. These results indicate that with ongoing development and larger scale evaluation, implementation of this IHCA is feasible.

References:

- [1] Lindsay, S., Kingsnorth, S. et al. (2014). A systematic review of self-management interventions for children and youth with physical disabilities. *Disability and Rehabilitation* 36(4): 276–288.
- [2] Murray, E., Burns, J. et al. (2005). Interactive health communication applications for people with chronic disease. *Cochrane Database of Systematic Reviews* (4).

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2017-eular.1025