levels or due to a better response to anti-TNF of those patient with higher plasma levels of 25 hydroxy-vitamin D.

References:

- [1] Nisar MK, Masood F, Cookson P, Sansome A, Östör AJ. What do we know about juvenile idiopathic arthritis and vitamin D? A systematic literature review and meta-analysis of current evidence. *Clin Rheumatol* 2013; 32:729–734.
- [2] Santos-Antunes J, Corte-Real Nunes A, Lopes S, Macedo G. The Relevance of Vitamin D and Antinuclear Antibodies in Patients with Inflammatory Bowel Disease Under Anti-TNF Treatment: A Prospective Study. *Inflamm Bowel Dis* 2016;22:1101–1106.
- [3] Ham M, Longhi MS, Lahiff C, Cheifetz A, Robson S, Moss AC. Vitamin D levels in adults with Crohn's disease are responsive to disease activity and treatment. *Inflamm Bowel Dis*. 2014; 20(5):856–60.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2017-eular.6232

AB0969 IMPACT OF JUVENILE IDIOPATHIC ARTHRITIS ON SCHOOLING

S. Rahmouni, A. Fazaa, K. Ouenniche, S. Kassab, S. Chekili, K. Ben Abdelghani, A. Laater. *Rheumatology Department, Mongi Slim hospital, marsa, Tunisia*

Background: Juvenile idiopathic arthritis (JIA) is the most common chronic rheumatic disease in children and is one of the major causes of morbidity and physical disability. Due to frequent absences, children with chronic health impairments are also often confronted with educational difficulties.

Objectives: The aims of this study were to assess the impact of JIA on children's schooling and to determine the factors that influence their school level.

Methods: This is a cross-sectional study including patients with JIA (ILAR criteria). A detailed questionnaire was completed for each participant by interviewing them or their parents as well as by information obtained from their medical records. Collected data included age, sex, subtype of JIA, disease duration, level of disability according to the Childhood Health Assessment Questionnaire (CHAQ), visual analog scale for patient's overall assessment of disease activity, duration of morning stiffness, joint counts, erythrocyte sedimentation rate, C-Reactive Protein, Disease Activity Score (DAS28). Medications used for JIA treatment were also documented.

Data on the school performance of patients and their siblings were obtained using telephone interviews (educational level, absenteeism, school delay by repetition, drop-out).

The comparison of quantitative variables was performed with the Mann-Whitney test and the comparison of qualitative ones was performed with the Chi square test. The significance level was set at 0.05.

Results: A total of 38 patients with JIA were included, 23 female and 15 male, with a mean age of 26 years [12–51] and a mean disease duration of 237 months [5–496]. The average age of the onset of the disease was 7.4 years [1.5–16].

The most common subtype was rheumatoid factor-positive polyarthritis (n=16) followed by systematic (n=7), oligoarticular (n=6), rheumatoid factor-negative polyarthritis (n=5) and Entesitis-related arthritis (n=4). The mean DAS28 was 2.63 [0.76 - 5.55] and the median CHAQ was 0.528 [0–3]. Twenty-seven of the children were receiving corticosteroid. Disease-modifying anti-rheumatic drugs were used by 34 of the 38 patients: methotrexate (n=23), sulfasalazine (n=8), leflunomide (n=7), biotherapies (n=14). Twenty patients had complications: Hip arthritis (n=15), growth stunting (n=12), uveitis (n=4). Joint replacement was required in 9 cases. Four patients were illiterate, 12 had dropped out of school, 21 reported repeated absences due to illness. A year of schooling was repeated by 61.7% of patients. Ten out of 32 patients over the age of 20 had an university level. Almost 80% of patients were exempted of physical education.

There were no significant associations between the school-related problems, the socio-demographic characteristics and the various parameters of clinical and biological activity studied. School failure was similar among patients and their siblings (p=0.05).

Conclusions: Our study suggested that JIA negatively affects schooling of children. More studies with a larger sample are needed to identify the variables associated with school failure in order to ensure the proper management of these patients and to increase their academic performance.

References:

- [1] Bouaddi et al. Impact of juvenile idiopathic arthritis on schooling. *BMC Pediatrics* 2013;13:2.
- [2] Kamrul Laila et al. Impact of Juvenile Idiopathic Arthritis on School Attendance and Performance. *American Journal of Clinical and Experimental Medicine* 2016;4(6):185–190.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2017-eular.6037

AB0970 CLINICAL AND IMMUNOLOGICAL CHARACTERISTICS IN CHILDHOOD-ONSET SYSTEMIC LUPUS ERYTHEMATOSUS PATIENTS TREATED WITH RITUXIMAB

S. Garrote, A. Alia Jimenez, A. Boteanu, M.A. Blazquez, M.L. Gamir. *Rheumatology, Ramon y Cajal Hospital, Madrid, Spain*

Background: Systemic lupus erythematosus (SLE) is an autoimmune disease that is more severe in pediatric population than in adults. Biological therapy with

anti-CD20 (rituximab) is an option in patient that do not respond to conventional therapy.

Objectives: The aim of this study is to determine the clinical and immunological response in 9 patients with childhood-onset systemic lupus erythematosus (cSLE) that received treatment with rituximab in a third level hospital

Methods: This is a retrospective observational study. 9 patients treated with Rituximab between November 2007 and October 2016 were included and their medical records were reviewed. The response to treatment at 6 months and one year after the first infusion of Rituximab were assessed. Patients with overlap syndromes were excluded. All patients fulfilled four or more of the 1982 revised American College of Rheumatology criteria for the diagnosis of SLE (<16 years).

Results: Nine pediatric patients with SLE treated with rituximab were included, all of them were female. The age at diagnosis of SLE was a mean of 15,22 years. The mean time duration of disease was 87,55 months (5–255m). 7 patients were caucasians. Rituximab was indicated in 6 patients with class IV of lupus nephritis (LN) 1/9 with class III LN, 1/9 with severe cutaneous lupus, and with severe hematological manifestations in 1 case (haemolytic anemia). In addition, 6/9 patients had mucocutaneous and articular manifestations. The disease activity of all patients was assessed using SELENA-SLEDAI index pre rituximab infusion, the mean was

17,11 (8–33). All patients had low level of complement C3 and C4 and 8/9 increased anti-DNA. In 8/9 patients Rituximab was used as a rescue treatment and in a single case as a first line.

3/6 patients with renal involvement were previously treated with cyclophosphamide (CF) iv and mycophenolate, 2/6 CF. In case of cutaneous involvement the previous treatment was methotrexate, azathioprine (AZA) and dapsone and in case of hemolytic anemia was AZA.

The treatment protocol was 1 gram x 2 (1 cycle) in 7/9 patients, 375mg/m² x 4 in 1/9 cases and 600mg monthly for 5 months in the case of hemolytic anemia. Five patients received more than 1 cycle. After the administration of Rituximab, the SELENA-SLEDAI activity index was 4.5 points. At 6 months a complete response was obtained in the case of hematological and cutaneous manifestations, in 2 cases of lupus nephritis (proteinuria <0.5 g/day) and partial response was obtained in 2 cases. Data were not analyzed in 2 patients (death and less than 6 months of the first dose of rituximab). Patients with partial response and lack of response achieved complete response at 12 months. 2/9 patients had side effects (Rituximab pneumonitis in 1 case and infections in 2 cases). Mortality was 11.11% (1/9 patients, per infection and lupus activity, SLEDAI pre rituximab =33).

Conclusions: In our study, although it consisted of few patients, it was objected that Rituximab therapy in patients with cSLE is effective, reduces lupus activity index, especially in cases of renal, cutaneous and hematologic involvement, that don't respond to conventional therapy. It may be consider in the future as an effective alternative treatment at first line treatment.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2017-eular.6846

AB0971 CHARACTERISATION OF THE PATIENTS AT THE TIME OF THE TRANSITION INTO THE ADULT RHEUMATOLOGY. 63% OF THE PATIENTS ARE IN REMISSION AT THE TIME OF THE TRANSITION

S. Tatsis¹, I. Foeldvari². ¹Department of Rheumatology, Marienkrankenhaus; ²Hamburg Center for Pediatric and Adolescent Rheumatology, Am Schön Klinik Eilbek, Hamburg, Germany

Background: The issue of transition from pediatric to adult rheumatology service is an emerging important topic. In 2012 a transition clinic was established according to the "Berliner Transitionsprogramm" [1] in cooperation of the Hamburger Zentrum für Kinder- und Jugendrheumatologie and the Rheumatology Unit of the Marien Hospital in Hamburg. The Berliner programm suggests three visits of the patient in transition process, in the presence of the pediatric and an adult rheumatologist together, and the fourth visit conducted by the adult rheumatologist alone. We present the characteristics of the patients at the time of the 4th visit.

Objectives: To characterize the patient population at the time of enrollment into the adult service in the frame of our transition programm.

Methods: We collected patient data starting 8/2012 to 11/2016. We summarized the patient population, who successfully transitioned from pediatric to adult rheumatologic service, concerning diagnosis, sex, age at the time of diagnosis, disease duration at the time of transition, JADAS, HAQ, VAS globular assessment, VAS pain, medication and disease activity.

Results: 73 patients were transitioned. 65% of them female. Mean age at diagnosis of the patients was 12.5 years. Mean disease duration at time of transition was 10.8 years. The mean JADAS Score was 3.18 and the mean HAQ Score was 0.136. The patients global activity score was, on a VAS of 0 to 100, 14.03 and the global pain score, on a VAS of 0 to 100, 12.33. 39.7% of the patients received synthetic DMARDs and 34% biologic DMARDs: Only 1 patient received steroids. 24.6% of the patients were off medication. 63% of the patients were in remission, 61% of them on medication and 39% off medication in remission.

Conclusions: In this monocenter cohort 63% of patients were in remission, and with the mean JADAS Score of 3.18 most of them have low disease activity under the current treatment. The mean HAQ Score with 0.136 reflects a score, which is expected in healthy controls. But 75.4% of the patients needed medication to