positive; 45.7% (N=21) were RF-ELISA positive and RF-nephelometry negative and 4.3% (N=2) the opposite, without a statistical significance.

Conclusions: ELISA is superior to nephelometry detecting RF in patients with RA, as also in quantifying high-positive values.

References:

[1] Routine measurement of IgM, IgG, and IgA rheumatoid factors: high sensitivity, specificity, and predictive value for rheumatoid arthritis. Swedler W, Wallman J, Froelich CJ, Teodorescu M. *J Rheumatol*, 1997 Jun;24(6):1037-44.

[2] Is rheumatoid factor still a superior test for the diagnosis of rheumatoid arthritis? Singh U, Vishwanath A, Verma PK et al. *Rheumatol Int*, 2010 June; 30(8):1115-9.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2017-eular.2606

AB1044 EVALUATION OF A FLUOROENZYME IMMUNOASSAY (ELIA-CTD) IN THE SCREENING OF PATIENTS SUSPECTED FOR AUTOIMMUNE CONNECTIVE TISSUE DISEASES

M. Elkhalfi¹, R. Abdulhadi¹, H. Ramadan¹, S. Barrientos¹, S. Saleh¹, O. Alsaed², A.-W. Al-Ailaf². ¹Laboratory Medicine and Pathology; ²Medicine, Hamad Medical Corporation, Doha, Qatar

Background: Detection of auto-antibodies directed against nuclear antigens (antinuclear antibodies or ANA) have important diagnostic and prognostic implications in connective tissue diseases (CTD). The conventional indirect immunofluorescence assay on Hep-2 cell line (ANA-IIF) is the most commonly used method to detect ANA. The ANA-IIF can be labor intensive and suffers from lack of specificity.

Objectives: To evaluate the utility of a new fluoroenzyme Immunoassay “EliA-CTD” as an alternative for screening patients suspected for autoimmune connective tissue diseases.

Methods: Sixteen Hundred (1600) consecutive patients' sera submitted for anti-nuclear antibodies were tested using the ANA-IIF (Diasorin S.P.A, Saluggia, Italy) and the new EliA-CTD screen (Phadia GmbH, Freiburg, Germany). ANA testing was ordered by both primary and secondary care physicians. The EliA-CTD screening assay is a fluoroenzyme immunoassay which is performed on the Phadia-250 automated platform. The EliA-CTD assay contains ANA-targeted recombinant antigens including dsDNA, Sm-D, Rib-P, PCNA, U1-RNP (70, A, C), SS-A/Ro, SS-B/La, Centromere B, Scl-70, Fibrillarin, RNA Polymerase III, Jo-1, Mi-2, and PM-scl. The test results are expressed as ratio, with >1.0 considered positive. For ANA-IIF, the cut off for positive results was 1:40 or greater. Additionally, further testing for dsDNA and other extractable nuclear antigens (ENA) was undertaken on a subset of sera that were ANA-IIF+ or whenever there was discrepancy between the two methods.

Results: The overall agreement between the two methods was 84.2%. Three hundred and eight (308) out of 1600 (19.3%) samples tested positive by ANA-IIF positive as compared to 101/1600 (6.6%) for the EliA-CTD assay. Additional testing showed that 105 samples were positive for ENA including dsDNA. Of those, 101 were EliA-CTD positive and 81 were ANA-IIF positive. By incorporating the ENA results, the calculated sensitivity and specificity for the EliA-CTD were 97.1% and 99.7% respectively with positive and negative predictive values for the EliA-CTD assay of 96.1% and 99.8%, respectively. The corresponding sensitivity, specificity, positive predictive value (PPV) and negative predictive value (NPV) for the ANA-IIF assay at different dilutions is shown below:

Titer	Sensitivity (%)	Specificity (%)	PPV (%)	NPV (%)
≥1:40	77.7	84.8	26.0	98.2
≥1:80	60.3	95.3	32.4	98.4
≥1:160	57.4	97.4	41.3	98.4
≥1:320	46.5	98.7	48.8	98.5

Conclusions: The new automated EliA-CTD assay shows superior sensitivity and specificity compared to the conventional labor intensive ANA-IIF. The EliA-CTD can be used as an upfront screening tool for connective tissue diseases. Depending on the clinical details, any EliA-CTD positive results could be reflexly followed by additional testing including ANA-IIF testing to elucidate the titer and pattern.

References:

[1] Hayashi N, Kawamoto T, Mukai M, Morinobu A, Koshiba M, Kondo S, Maekawa S, Kumagai S. Detection of Antinuclear Antibodies by Use of an Enzyme Immunoassay with Nuclear Hep-2 Cell Extract and Recombinant Antigens: Comparison with Immunofluorescence Assay in 307 Patients. *Clinical Chemistry* 2001; 47(9):1649-1959.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2017-eular.2679

AB1045 MACROPHAGE TARGETED POSITRON EMISSION TOMOGRAPHY (PET) FOR THE IMAGING OF INFLAMMATORY ARTHRITIS; AN IN VIVO AND IN VITRO INVESTIGATION OF TRANSLOCATOR PROTEIN (TSPO) TRACER UPTAKE

N. Narayan¹, H. Mandhair¹, C. Coello², A. Saleem², A. Sabokbar¹, P. Taylor¹.

¹Nuffield Dept of Orthopaedics, Rheumatology, Musculoskeletal Sciences, University of Oxford, Oxford; ²Imanova Centre for Imaging Sciences, London, United Kingdom

Background: TSPO targeted PET tracers are increasingly recognised as cellular imaging markers of macrophage infiltration, due to the high expression of TSPO on activated macrophages. Previous work demonstrated the ability of [¹¹C]PK11195 TSPO PET to detect subclinical inflammation in RA, and predict flare in both those with established RA and ACPA positive arthralgia. However, high background uptake of [¹¹C]PK11195 in bone, and inability of [¹¹C]PK11195 to detect lesions in AS has driven the investigation of newer TSPO tracers for the detection of inflammatory arthritis.

Objectives: Here, we present data confirming the ability of the TSPO tracer [¹¹C]PBR28 to detect and quantify synovitis in both RA and PsA, and in vitro work that assesses more fully what TSPO tracer accumulation in inflamed synovium actually reflects at a cellular level, especially considering TSPO is ubiquitously expressed.

Methods: 10 patients (5 with RA, 5 with PsA) with evidence of inflammation in one or both knees (as confirmed by clinical examination and US) and 4 healthy volunteers underwent PET/CT both knees using the TSPO tracer [¹¹C]PBR28. Arthritis patients underwent synovial biopsy of one knee within 7 days of scan. Healthy synovium was obtained from patients undergoing knee arthroscopy for ligamentous injury. Synovial tissue was stained for CD68, CD163 and TSPO. For in vitro work, human monocytes, lymphocytes and synovial FLS from RA patients were harvested, and macrophages differentiated from monocytes. RNA was extracted for PCR. Other cells underwent density centrifugation to extract the cytoplasmic cell fraction, and a radioligand binding assay with [³H]PBR28 was undertaken, to assess tracer binding to TSPO in each cell type

Results: Tracer uptake correlated significantly with severity of inflammation on clinical examination and ultrasound, as did synovial sublining staining for CD68, TSPO and CD163 (see table 1). There was negligible staining for all stains in healthy control synovium. qPCR demonstrated highest TSPO mRNA in stimulated FLS (fold change 62.75±10.03) and M2 macrophages (60.69±2.38), with lymphocytes having the least TSPO expression. PBR28 saturation binding confirmed these findings at protein level (see graph 1).

Tissue marker	Clinical synovitis severity		
	1 (mild)	2 (moderate)	3 (severe)
TSPO	1.50±0.08	2.19±0.19	3.06±0.12
CD68	2.17±0.3	2.42±0.21	2.96±0.05
CD163	1.61±0.03	2.19±0.1	2.81±0.04

Conclusions: Our data demonstrates that the TSPO tracer PBR28 is capable of detecting and quantifying synovitis in RA and PsA. PBR28 tracer uptake correlates with macrophage marker staining, but not with fibroblast marker staining in our patient cohort. mRNA and protein data demonstrate, however, that there is a similar expression of TSPO in activated macrophages and activated FLS, hence TSPO tracer accumulation is as likely to represent FLS activation as it is macrophage activation.

References:

[1] Gent YY et al. Macrophage positron emission tomography imaging as a biomarker for preclinical rheumatoid arthritis: findings of a prospective pilot study. *Arthritis Rheum* 2012; 64:62-6.

[2] Gent YYJ et al. Subclinical synovitis detected by macrophage PET, but not MRI, is related to short-term flare of clinical disease activity in early RA patients: an exploratory study. *Arthritis Res Ther* 2015; 17:266.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2017-eular.1838

AB1046 COULD SERUM TWEAK LEVEL BE AN INDICATOR OF SUBCLINICAL ATHEROSCLEROSIS IN RHEUMATOID ARTHRITIS?

M. Ucer¹, N. Yilmaz², B. Cengiz Elcioglu³, T. Sahin³, P. Gun Atak⁴, S. Yavuz². ¹Internal Medicine; ²Rheumatology; ³Cardiology; ⁴Biochemistry, Istanbul Bilim University, Istanbul, Turkey

Background: TWEAK is a type 2 transmembrane glycoprotein of TNF family that has multiple functions such as angiogenesis, regulation of tissue production-