of foot pain and foot care among rheumatoid arthritis patients taking and not taking anti-TNFalpha therapy: an epidemiological study. *Rheumatol Int.* 2011;31:1515–9.

[3] Hennessy K, Woodburn J, Steultjens MPM. Custom foot orthoses for rheumatoid arthritis: A systematic review. *Arthritis Care Res (Hoboken)*. 2012;64:311–20.

[4] Conrad KJ, Budiman-Mak E, Roach KE, Hedeker D, Caraballada R, Burks D, Moore H. Impacts of foot orthoses on pain and disability in rheumatoid arthritis. *J Clin Epidemiol.* 1996 Jan;49(1):1–7.

[5] Novak P, Burger H, Tomsic M, Marincek C, Vidmar G. Influence of foot orthoses on plantar pressures, foot pain and walking ability of rheumatoid arthritis patients—a randomised controlled study. *Disabil Rehabil.* 2009;31(8):638–45.

Disclosure of Interest: None declared
DOI: 10.1136/annrheumdis-2017-eular.2739

THURSDAY, 15 JUNE 2017

HPR measuring health (development and measurement properties of PROs, tests, devices) —

THU0752-HPR

DEVELOPMENT OF THE “TREATMENT BELIEFS IN KNEE AND HIP OSTEOARTHRITIS (TOA)” QUESTIONNAIRE

E.M. Seltén¹, J.E. Vriezcekolk¹, H.J. Schers², M.W. Nijhof³, W.H. van der Laan⁴, R.G. van der Meulen-Dilling⁵, R. Geenen⁶, C.H. van den Ende¹.

¹Rheumatology, Sint Maartenskliniek; ²Primary and Community Care, Radboud University Nijmegen Medical Centre; ³Orthopaedics, Sint Maartenskliniek, Nijmegen; ⁴Rheumatology, Sint Maartenskliniek, Woerden; ⁵Physical Therapy and Manual Therapy, Partnerschip Velperweg, Arnhem; ⁶Psychology, Utrecht University, Utrecht, Netherlands

Background: Use of non-surgical treatment modalities in osteoarthritis (OA) is suboptimal¹, which might be influenced by patients’ beliefs about treatments. An instrument for measuring treatment beliefs in OA is not available yet.

Objectives: To develop a questionnaire assessing patients’ beliefs about treatment modalities of hip and knee OA: the “Treatment beliefs in OA (TOA) questionnaire” and to evaluate its clinimetric properties.

Methods: The item pool, drawn from two previous qualitative studies, comprised beliefs regarding 5 treatment modalities: physical activity, pain medication, physiotherapy, injections and arthroplasty. A draft questionnaire comprising beliefs on these 5 treatment modalities was developed (200 items, Table 1). Two samples of patients with knee or hip OA (N=840, N=700) were recruited from our hospital to test the clinimetric properties of the TOA questionnaire. Descriptive analyses, explorative factor analyses (EFA; sample 1) and confirmatory factor analyses (CFA; sample 2) were conducted for each treatment module separately. Internal consistency was assessed with Cronbach’s Alpha (both samples). In order to examine test-retest reliability a subsample of sample 2 (N=67) was asked to fill out the final TOA questionnaire again after two weeks.

Results: 351 patients filled out the draft TOA questionnaire (sample 1), 289 patients filled out the final TOA questionnaire (sample 2), with a subsample (N=50) who filled out the final questionnaire twice. EFA yielded a two factor solution for each treatment modality. The factors were labeled “positive treatment beliefs” and “negative treatment beliefs”. The final TOA questionnaire comprised 60 items; items per treatment modality ranged from 9 to 14. CFA showed adequate fit indices for physical activities and physical therapy, while fit indices for the treatment modalities pain medication, injections and arthroplasty just failed to reach adequate cut-off values. Internal consistency was good to excellent for the subscale positive treatment beliefs (Cronbach’s α between 0.84 and 0.90), and mediocre to acceptable for the subscale negative treatment beliefs (Cronbach’s α between 0.66 and -0.79). Test-retest reliability was satisfactorily to good with ICCs from 0.66 to 0.88 and SEMs from 0.17 to 0.52.

Table 1: Example items in the TOA questionnaire

Items TOA-questionnaire	Disagree				Agree
I think [doing physical activities] leads to risks	1	2	3	4	5
My quality of life improves through [using pain medication]	1	2	3	4	5
I think [physiotherapy] causes pain	1	2	3	4	5
I can do household chores more easily through [an injection]	1	2	3	4	5
I think [a joint replacement] is pervasive	1	2	3	4	5

Conclusions: The TOA questionnaire is the first questionnaire assessing positive and negative beliefs regarding five treatment modalities for knee and hip OA. The TOA questionnaire showed moderate structural validity and good internal consistency and test-retest reliability. The TOA questionnaire is useful for a better understanding of patients’ treatment beliefs. Future research will examine how treatment beliefs, in interaction with other variables, influence treatment choices.

References:
[1] Churchill 2016. The development and validation of a multivariable model

to predict whether patients referred for total knee replacement are suitable surgical candidates at the time of initial consultation. *Can J Surg.* Vol. 59, No. 6, pp. 407–414.

Disclosure of Interest: None declared
DOI: 10.1136/annrheumdis-2017-eular.4815

THURSDAY, 15 JUNE 2017

HPR epidemiology and public health (including prevention) —

THU0753-HPR

PREDICTING THE FUTURE DEVELOPMENT OF SPONDYLOARTHRITIS AMONG PATIENTS WITH IDIOPATHIC ACUTE ANTERIOR UVEITIS USING REAL-WORLD DATA

M. Haroon¹, K.A. Betts², F. Mu², M. Skup³, J.K. Anderson³, A.D. Joshi³.

¹Division of Rheumatology, University Hospital Kerry, Tralee, Ireland; ²Analysis Group, Boston; ³AbbVie Inc., North Chicago, United States

Background: Previous studies have suggested an association between idiopathic acute anterior uveitis (AAU) and spondyloarthritis (SpA), including showing that at least 40% of AAU patients have undiagnosed SpA. However, the clinical factors to predict the incident diagnosis or future development of SpA remain poorly recognized.

Objectives: The objectives of this study were: 1) to describe the patient characteristics of AAU patients with and without SpA diagnosis, and 2) to identify the predictive factors of SpA diagnosis among AAU patients using real-world data.

Methods: Adult patients with at least one diagnosis of AAU (ICD-9-CM: 364.01 or 364.02) were selected from a large US insurance claims database (01/01/2008–06/30/2015). The first AAU diagnosis was defined as the index date. Patients were required to have at least 6 months of continuous data availability before the index date. Patients with intraocular surgery, penetrating or blunt eye trauma, or a diagnosis of rheumatoid arthritis or SpA on or prior to the index date were excluded. Potential predictive factors for subsequent diagnosis of SpA included demographic characteristics, type of AAU (primary [first diagnosis] vs. recurrent), SpA-related comorbidities, and healthcare resource utilization prior to the index date. Factors predictive of SpA diagnosis were selected into a multivariable Cox proportional hazards model based on statistical significance and clinical relevance. Hazards ratios (HR) and p-values were estimated for each factor.

Results: A total of 48,822 patients with AAU were included, and among them, 1,032 patients were newly diagnosed with SpA during the follow-up period which was 24 months on average. Patients with SpA were younger (45.7 vs. 50.4 years), more likely to be male (52% vs. 42%), more likely to have recurrent AAU (44% vs. 29%), back pain (21% vs. 13%), SpA-related comorbidities including inflammatory bowel disease (IBD) (4% vs. 1%) and psoriasis (3% vs. 1%), and more likely to use corticosteroids (45% vs 40%). Predictive factors in the final Cox model were: male vs. female (HR=1.55; p-value<0.01), age<45 vs. ≥45 years old (1.65; <0.01), recurrent vs. primary AAU (1.94; <0.01), back pain under age 45 vs. no back pain (1.90; <0.01), back pain above age 45 vs. no back pain (1.46; <0.01), psoriasis (5.16; <0.01), IBD (2.50; <0.01), chiropractor/physical therapist visits (1.29; 0.01), conjunctivitis (1.23; 0.02), nonsteroidal anti-inflammatory drugs use (1.44; <0.01), use of conventional synthetic disease-modifying anti-rheumatic drugs (1.32; 0.05), joint pain (1.14; 0.19), imaging test use (1.03; 0.74), corticosteroids use (1.07; 0.33), and enthesitis/heel pain (1.13; 0.36). When sacroiliitis was included in the model, results remained largely similar.

Conclusions: There are significant differences among isolated AAU patients and AAU patients that developed SpA later. The most predictive factors of SpA diagnosis were male, age<45 years, recurrent AAU, back pain, and other extra-articular manifestations of SpA such as IBD and psoriasis. Since delayed diagnosis is common among SpA patients, identifying such predictive factors can help inform risk stratification.

Disclosure of Interest: M. Haroon Grant/research support from: AbbVie and Pfizer, K. Betts Consultant for: AbbVie, F. Mu Consultant for: AbbVie, M. Skup Employee of: AbbVie, J. Anderson Employee of: AbbVie, A. Joshi Employee of: AbbVie
DOI: 10.1136/annrheumdis-2017-eular.3028

THU0754-HPR

WORK STATUS IN WOMEN WITH FIBROMYALGIA – A 12-YEAR FOLLOW UP STUDY

S. Johannesson¹, L. Nordeman^{2,3}, M. Joelsson¹, K. Mannerkorpi², A. Bergenheim^{1,2,3}. ¹Närhälsan rehabilitation centre, Region Västra Götaland; ²Institute of Neuroscience and Physiology/Physiotherapy, Sahlgrenska Academy, University of Gothenburg, Gothenburg; ³Närhälsan Research and Development Primary Health Care, Region Västra Götaland, Sweden

Background: Fibromyalgia (FM) affects approximately 1–3% of the general population in the Western world and about 10% have been found to report Chronic Widespread Pain (CWP). The work ability is often affected in persons with FM and CWP. Previous research has shown a small improvement of symptoms over time in patient with FM and CWP. Sustain in work has been