of people with rheumatic diseases and fatigue. The current intervention was developed as a supplement to medical treatment to strengthen coping and quality of life, and reducing fatigue, depression and pain.

Methods: People with inflammatory rheumatic diseases and fatigue were recruited for 6–9 sessions of cognitive therapy sessions, in addition to treatment as usual at a rheumatology outpatient clinic in Norway. The intervention aimed at reducing fatigue, depression and pain, consisted of four main elements; understanding fatigue, assessment and activity planning, mental and cognitive self-help skills. The project was developed, and data collected at the rheumatology outpatient clinic at Diakonhjemmet Hospital in 2014–2016.

Results: This pilot project recruited 40 people with inflammatory rheumatic disease from a rheumatologic outpatient clinic in Norway. Participants had a disease duration of mean 14 years, they were mainly women (n=36) with a mean age of 45 (Standard Deviation =10) years (Table 1). Repeated ANOVA analyses and Paired t-tests showed promising statistically significant changes on a group level for fatigue and depression, not for pain (Table 2).

Table 1. Demographic data of included patients (n=40)

N=40	Mean values
Age (SD)	45 (10)
Women (men)	36 (4)
Disease duration (years)	14 (11)

Conclusions: This pilot project supports the idea of conducting a trial on the effectiveness of a brief cognitive therapy intervention for people with inflammatory rheumatic diseases and fatigue. ANOVA analyses and Paired t-tests showed promising statistically significant changes on a group level for fatigue and depression, but not for pain, whether these changes are clinically meaningful, and if there is a difference, compared to treatment as usual remains to be explored.

Acknowledgements: This project would not have been possible without the support from the national patient organization, through a grant from the Norwegian ExtraFoundation for Health and Rehabilitation. We acknowledge patients who contributed to this project and the staff at the rheumatological outpatient clinic at Diakonhjemmet Hospital.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2017-eular.6096

AB1245-HPR NEW HEALTH TECHNOLOGIES AND LIFESTYLE MANAGEMENT FOR PATIENTS WITH OSTEOARTHRITIS

M. Loiskandl¹, H. Ritschl², T. Stamm¹, G. Steiner¹, J. Smolen¹, D. Aletaha¹.

¹Rheumatology, Medical University of Vienna, Vienna; ²Joanneum, University of Applied Sciences, Graz, Austria

Background: Osteoarthritis (OA) is one of the most common chronic diseases of the elderly worldwide. It represents significant impairments in terms of quality of life [1]. It notably affects self-care-tasks, body-image, self-esteem, well-being, social-activities and relationships. A wide-ranged field of therapy concepts exist. Medication is often one part, but it cannot solve other existing problems due to OA [2]. The demographic trends combined with the growth of mobile devices among the older population suggest that using digital devices, as a platform for interventions in health, may be a viable way forward and open new opportunities in healthcare [2].

Objectives: To (A) analyse the literature to assistive devices and applications for OA patients and (B) to evaluate existing health applications.

Methods: We performed a systematic literature review on this topic and explored existing mobile and web-based apps. The app findings were classified into those categories of the Activities of Daily Living Model, which were related to the impairments of OA.

Results: We identified 6 studies, which contained relevant information. The main results showed, that there is interest and willingness to utilize mobile Health solutions [3]; a positive impact of web-based interventions; the benefit of online social support regarding the level of patient empowerment [4], self-management, lifestyle, physical activity [5] and self-efficacy [6].

The app exploration results in 16 web-based, 15 iOS and 9 Android applications. The areas most commonly found were disease- and treatment information (n=23),

forum, report, self-help groups (n=16), self- and disease-management (n=11), as well as nutrition tips (n=8).

Conclusions: The recent years have seen a dramatic growth of digital applications. Also, the scientific interest has grown and numerous studies emphasise the power of these solutions to support OA patient in all areas of lifestyle management. Further research is necessary, to investigate the benefit of web-based or mobile applications for persons with OA.

References:

- [1] T. Aigner, S. Söder, Typisierung, Graduierung und Stadieneinteilung der Osteoarthritis: Histopathologische Begutachtung der Gelenkdegeneration, *Z. Für Rheumatol.*, 2008;67(1):32–40.
- [2] J. Joe, G. Demiris, Older adults and mobile phones for health: A review, *J. Biomed. Inform.* 2013;46(5):947–954.
- [3] S.J. Parker, S. Jessel, J.E. Richardson, M.C. Reid, Older adults are mobile tool/identifying the barriers and facilitators to older adults' use of mHealth for pain management, *BMC Geriatr.* 2013;13(1):43.
- [4] A. Allam, Z. Kostova, K. Nakamoto, P. J. Schulz, The Effect of Social Support Features and Gamification on a Web-Based Intervention for Rheumatoid Arthritis Patients: Randomized Controlled Trial, *J. Med. Internet Res.* 2015;17(1):e14.
- [5] H. Umapathy, K. Bennell, C. Dickson, F. Dobson, M. Fransen, G. Jones, D.J. Hunter, The Web-Based Osteoarthritis Management Resource My Joint Pain Improves Quality of Care: A Quasi-Experimental Study, *J. Med. Internet Res.* 2015;17(7):e167.
- [6] K.R. Lorig, P.L. Ritter, D.D. Laurent, K. Plant, The internet-based arthritis self-management program: A one-year randomized trial for patients with arthritis or fibromyalgia, *Arthritis Rheum.* 2008;59(7):1009–1017.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2017-eular.6652

AB1246-HPR INTRA ARTICULAR OZONE OR HYALURONIC ACID INJECTION: WHICH ONE IS SUPERIOR IN PATIENTS WITH KNEE OSTEOARTHRITIS? A 6-MONTH RANDOMIZED CLINICAL TRIAL

S.A. Raeissadat¹, S.M. Rayegani², M. Moridnia³, S. Rahimi Dehgolan⁴.

¹Physical Medicine and Rehabilitation Department, Shahid Modarres Hospital;

²Physical Medicine and Rehabilitation Department, Shohadaye Tajrish Hospital;

³Deputy of Education; ⁴Physical Medicine and Rehabilitation Department,

Shahid Beheshti University of Medical Sciences, Islamic Republic Of

Background: Knee osteoarthritis (OA) is a common disease with great burden through pain and decreased function (1). Except for knee replacement surgery as a definite treatment (2), there are many other therapeutic modalities including non-pharmacologic choices and some heterogeneous drugs like oral, topical and intra articular ones (3). Recent studies have showed promising results of ozone application in OA (2,4–5).

Objectives: The aim of this study was to compare the effects of ozone therapy versus hyaluronic acid (HA) intra articular injection in patients with knee OA.

Methods: This prospective RCT studied 174 patients in two random groups: ozone and HA. We performed three weekly injections of Hyalgan® versus 10 ml of ozone solution in moderate knee OA patients. Participants were primarily evaluated and 6 months after injection using pain, stiffness and function questionnaires.

Results: No major adverse events were noted in the study. Total WOMAC score decreased in the ozone group from 40.8±9.8 to 20.4±4.9 (p<0.05) and in the HA group from 38.5±7.9 to 17.1±4.2 (p<0.05). A similar trend was observed in pain improvement according to VAS (table 1). Pain, stiffness and function significantly improved in both groups but no differences were found between groups.

Conclusions: Although both ozone and HA can be effectively used for improving function and reducing pain in selected patients with knee OA, neither of the two showed no superiority at 6 months follow up.

References:

- [1] Richmond J, Hunter D, Irrgang J, et al. American Academy of Orthopaedic Surgeons clinical practice guideline on the treatment of osteoarthritis (OA) of the knee. *J Bone Joint Surg Am* 2010; 92:990–993.
- [2] Kr Mishra S, Pramanik R, Das P, Pratim Das Pl. Role of intra-articular ozone

Abstract AB1244-HPR – Table 2. Self-reported fatigue and depression at baseline, after treatment and at 6 months follow-up

Scale	Mean (SD) t1	Mean (SD) t2	Mean (SD) t3	P-value diff ANOVA	Paired t-test T1–3
BRAF fatigue ^a	13.89 (5.4)	10.85 (4.9)	9.46 (5.8)	p>0.001	Diff 4.57 (CI: 2.2/7.0) (p>0.001)**
PHQ ^a	25.13 (5.9)	20.67 (6.8)	18.25 (7.1)	p>0.001	Diff 6.44 (CI: 4.0/8.9) (p>0.001)**
VAS pain [†]	45.54 (24.4)	39.08 (21.1)	36.9 (23.0)	p>0.241	Diff 7.64 (CI: -1.4/16.7) (p>0.096)

^aBRAF (Bristol Rheumatoid Arthritis Fatigue Multi-Dimensional questionnaire) fatigue global score (0–30), ^aPHQ-9 (Patient Health Questionnaire) depression score (0–70), [†]VAS pain (0–100).

Abstract AB1246-HPR – Table 1

Measure	Ozone Therapy (Mean ± SD)		Raw Mean Difference (SE) [95% CI]	HA Therapy (Mean ± SD)		Raw Mean Difference (SE) [95% CI]
	Before	After		Before	After	
VAS	7.6±2.8	2.6±2.0	-5.0 (0.2) [4.6–5.4]	7.1±3.2	3.0±2.4	-4.1 (0.15) [3.8–4.4]
WOMAC Pain	9.3±4.4	3.2±1.6	-6.1 (0.2) [5.7–6.5]	8.8±4.0	2.9±1.6	-5.9 (0.2) [5.5–6.3]
WOMAC Stiffness	2.3±2.4	1.1±1.6	-1.2 (0.05) [1.1–1.3]	2.1±1.6	1.1±0.8	-1.0 (0.05) [0.9–1.1]
WOMAC Function	29.2±7.0	16.1±4.2	-13.1 (0.15) [12.8–13.4]	27.6±6.6	13.1±3.2	-14.5 (0.3) [13.9–15.1]
WOMAC Total	40.8±9.0	20.4±5.0	-20.4 (0.2) [20.0–20.8]	38.5±8.0	17.1±4.2	-21.4 (0.2) [21.0–21.8]

in osteoarthritic knee for functional and symptomatic improvement. *IJPMR* 2011;22:65–69.

[3] Bellamy N, Campbell J, Robinson V, Gee T, Bourne R, Wells G. Viscosupplementation for the treatment of osteoarthritis of the knee. *Cochrane Database Syst Rev* 2006;(2):CD005321.

[4] Sello Lebohag Manoto, Makwese Johanness Maepa, Shirley Keolebogile Motaung. Medical ozone therapy as a potential treatment modality for regeneration of damaged articular cartilage in osteoarthritis [Article in Press]. *Saudi Journal of Biological Sciences* (2016).

[5] Velio Alvaro Bocci. Scientific and Medical Aspects of Ozone Therapy: State of the Art. *Archives of Medical Research* 37 (2006) 425–435.

Disclosure of Interest: S. A. Raeissadat Shareholder of: None, Grant/research support from: None, Consultant for: None, Employee of: None, Paid instructor for: None, Speakers bureau: None, S. M. Rayegani Shareholder of: None, Grant/research support from: None, Consultant for: None, Employee of: None, Paid instructor for: None, Speakers bureau: None, M. Moridnia Shareholder of: None, Grant/research support from: None, Consultant for: None, Employee of: None, Paid instructor for: None, Speakers bureau: None, S. Rahimi Dehghan Shareholder of: None, Grant/research support from: None, Consultant for: None, Employee of: None, Paid instructor for: None, Speakers bureau: None

DOI: 10.1136/annrheumdis-2017-eular.5484

HPR professional education, training and competencies

AB1247-HPR AN INTEGRATED APPROACH IN CUSTOMIZED MEDICAL FEMORAL COMPONENT PRODUCTION TO MINIMIZE THE CONDYLE CUT OFF DURING THE TKA

M. Moayedfar, A.M. Abdul Rani. *Mechanical Engineering, Universiti Teknologi Petronas, Bandar Seri Iskandar, Malaysia*

Background: To prepare the femoral component during the Total knee arthroplasty (TKA), it is needed to cut off the damaged condyle to a dimension matched with one of the available implant sizes in the market. Since there is a variety of humans' knee size and shape, therefore, some parts of the distal femoral are over cut. The solution to this problem is customisation of the implants via a cohesive procedure specifically designed for medical orthopaedic implant production.

Objectives: This research aims to provide a fully integrated system of digital image processing of Magnetic Resonance Imaging (MRI) data and Finite Element Method (FEM) for testing and analysing prior to manufacturing of customized medical femoral component.

Methods: An automatic algorithm to segment MR images from end-of-femur condyle is developed and employed for 3D reconstruction followed by computer-aided design (CAD) system, FEM and incremental sheet forming (ISF) process. In femur segmentation, multi-resolution edge detection is applied that extracts all the edges at bone surfaces. The edge detection process followed by the selection and extraction of strong edges with enhanced active contour technique that results in separation between femur and tibia components. Morphological processing is applied to extract the femur region as the segmented outcome. Consequential of segmentation, a surfaces rendering method is applied for the 3D structure that automatically converts the model into standard tessellation language (STL) format for later CAD system and rapid prototyping preparation. Generated STL file is applied for creating a solid part and solid body. The Proper femoral component is designed using region growing technique then exported to FEM system to calculate the best metal thickness and angles, regarding the normal cartilage, to bear the maximum load with the lighter material. The process of ISF that involves machining and sheet forming parameters is using to manufacture customized medical metal femoral component but in low surface quality which increases the risk of ion release at *in vivo* condition. Therefore, during this technique, optimum parameters using Design of Experiment method are applied to modify and enhance the sheet stretching of final parts.

Results: Applying this integrated system demonstrates a cohesive user-friendly system for creation of patients specific implants while it showed a substantial reduction in implant production time compares to the current manufacturing method. In ISF process optimum parameters are used to increase the rate of stretching together with the best sheet thickness ~2.5 mm based on validated sheet stretching simulation. The results of this study also show that the high surface quality for femoral component up to ~Ra 2.527 is achievable using this newly developed method.

Conclusions: An integrated system that combines computational method, CAD, FEM and ISF procedure is developed that shows its potential for manufacturing of customized medical femoral component. Flexibility in size and shape make this system prominent. The best sheet thickness with respect to the cartilage thickness is calculated and obtained with the optimum process parameters for manufacturing section.

Acknowledgements: The authors are thankful to the Ministry of Higher Education, Malaysia for supporting this research via FRGS/1/2014/TK01/UTP/02/8 research grant.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2017-eular.1351

AB1248-HPR THE ROLE OF PATIENT SUPPORT PROGRAMMES IN HEALTH LITERACY OF PATIENTS. RESULTS OF A 1.5-YEAR-LONG HUNGARIAN STUDY

A. Domján¹, Z. Szekanez¹, J. Pilling², G. Purebl², Z. Szabo¹, P. Kakuk³.
¹Department of Rheumatology, University of Debrecen Faculty of Medicine, Debrecen; ²Institute of Behavioural Sciences, Semmelweis University, Budapest; ³Institute of Behavioural Sciences, University of Debrecen, Faculty of Public Health, Debrecen, Hungary

Background: Health literacy means our ability to find our way in health issues, to understand health information and how to utilize them, that is how we can understand and use health information. Unfortunately, every second person has serious deficit in this area in Hungary.

Objectives: We wished to determine the effectiveness of a specialized patient support program in Hungary.

Methods: The present study was conducted between August, 2015 and December, 2016 in Hungary. Altogether 24 health professionals in 15 investigational sites including rheumatology, gastroenterology and dermatology centres were involved. All participating health professionals was qualified and skilled in patient education. We tried to enrol mainly new patients with relatively little knowledge of their chronic illness. The participation was completely voluntary. The programme was based on 4 presentations on various but closely related topics, presented in small-group sessions by health professionals, and then discussed with the patients. The topics of the lectures were as follows: How to manage stress, associated with the disease? How to deal with the emotional burden of the disease? How to use social support when fighting a chronic disease? How should I tell? (physician-patient communication). The theoretical framework of the lectures was composed by 2 psychiatrists, proficient in the subject. The health professionals had also received extensive training before the programme. Altogether 1950 participants were present at least one of the lectures: Only those were carried forward and allowed to complete the questionnaire who participated in all 4 lectures of different topics. Thus, 480 patients completed the 8 questions of the questionnaire that evaluated the effectiveness of the programme.

Results: Among the 480 patients, who completed the study, 66% had joint and musculoskeletal complaints, 32% had inflammatory skin or gastrointestinal disease and 2% of them suffered from other chronic diseases. Altogether 24% was a member of a patient organization. Two-third of the patients felt better prepared for their consecutive doctor-patient or nurse-patient appointment after participating in the programme. In total, 49% of the responders found the doctor-patient communication lecture most helpful. In addition, 73% of them were able to successfully manage disease-related events and stress. Two-third of the patients wished to contact and/or keep in contact with a patient organization in order to take advantage of the opportunities of social support and 76% understood the place and role of patient organizations in the society. We wish to continue the programme in 2017 and we are aiming to target communication channels appropriate for the needs and opportunities of our age (e.g. Facebook).

Conclusions: The educational programme described above was successful among our patients. We think that appropriate health literacy can lead to the better understanding of the disease and maybe to the improvement of unfavourable health indicators.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2017-eular.1514

AB1249-HPR THE EFFECT OF TASK-ORIENTED TRAINING ON FUNCTIONALITY, DEXTERITY AND ADL PERFORMANCE IN RHEUMATOID ARTHRITIS

Ö.F. Özçelep¹, E.I. Üstün², Z.C. Algun³. ¹Prosthetics Orthotics Research Center, Istanbul Medipol University; ²Physical Therapy and Rehabilitation, Başçılar Education and Research Hospital; ³Orthotics and Prosthetics, Istanbul Medipol University, Istanbul, Turkey

Background: Rheumatoid arthritis is the most common and most serious of the inflammatory arthritides, and it dominates clinical rheumatological practice (1). This inflammation leads gradually to a destruction of bone and cartilage, responsible for loss of function (2). The main goals of treatment for RA are to prevent or control joint damage, prevent loss of function, and decrease pain (3).

Objectives: The purpose of the study is to investigate the effect of task oriented training on functionality, dexterity and ADL performance in rheumatic hands. Also the other goal of this study is to bring a new perspective to RA rehabilitation.

Methods: Thirty seven women (16 patients in control group and 17 patients in study group) performed therapy programme. Control group received hand exercise therapy (MCP, DIP, PIP mobilization, range of motion exercise, isometric exercise) and study group received both hand exercise therapy and task oriented training (fork use, drinking water with a glass, face washing, sit up-stand up and t-shirt wearing) twice a week for 5 weeks. Every therapy sessions took 40–45 minutes for study group and 20–25 minutes for control group. According to the patient's condition, the rest interval was given and it was said that the exercises should not be done too fast. The results were evaluated before and after the therapy program with Jamar Hand Dynamometer, Nine Hole Peg Test, Health Assessment Questionnaire and Duruöz Hand Index.

Results: Age distribution of participants was 49, 57 ($p > 0.005$) and were found to be homogeneously dispersed ($p = 0.15$). Hand grip strength was not statistically