years in axSpA patients. Radiographic findings of cam abnormality (figure) were significantly more frequent in axSpA patients in comparison with control subjects (30/150 [20%] vs 17/193 [9%] and $P=0.004$). Cam-type radiographic abnormality was only present in 2 female control subjects and none of female axSpA patients. FAI was significantly correlated with the presence of HLA-B27 ($r=0.213$ and $P=0.048$), smoking ($r=0.194$ and $P=0.018$), height ($r=0.283$ and $P=0.001$) and gender ($r=0.443$ and $P<0.001$).

Conclusions: Our results showed that radiographic findings compatible with PGD were frequent in axSpA patients. In addition to repetitive injury to the proximal femoral physis, new bone formation may be responsible for increased FAI in axSpA. In axSpA patients with hip or trochanteric pain, FAI may be kept in mind as an alternative explanation of the symptoms.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2017-eular.3678

FRI0463 OPTIMIZING THE ASSESSMENT OF SPONDYLOARTHROPATHY AMONG OPHTHALMOLOGISTS – A REGIONAL SURVEY

S. Goel¹, P. Kumar², A. Moorthy². ¹Medical student, University of Leicester; ²University Hospitals of Leicester NHS Trust, UK, Leicester, United Kingdom

Background: Spondyloarthropathy (SpA), is an under diagnosed clinical condition with an average delay in diagnosis of up to 8 years. The delay occurs due to non-specific symptoms which occur in young patients who are unlikely to present to healthcare services as well as presentations to different specialties as a result of extra-articular manifestations.

30% of Ankylosing Spondylitis (AS) patients have acute anterior uveitis (AAU) at some point due to the common link with HLA B-27. Uveitis may precede the diagnosis of Ankylosing Spondylitis (AS) by several years. Previous studies have shown that up to 40% (1) people with AAU have undiagnosed SpA, and are not routinely seen in rheumatology services due to lack of symptom and risk recognition. Early diagnosis in this group is essential to reduce the delay in diagnosis and improve the outcome of SpA.

Objectives: 1. To identify the current assessment of inflammatory back pain, in patients with recurrent AAU, by local ophthalmologists. 2. To identify ways to improve cross specialty referral and reduce delay in the diagnosis of SpA.

Methods: This is a prospective, questionnaire based semi qualitative study. The target population includes specialist ophthalmology trainees, consultants and GPs with a special interest in ophthalmology across the East Midlands. The questionnaire was piloted locally at a regional uveitis meeting. The improved questionnaire was then electronically communicated to ophthalmologists across East Midlands. The data was collated and analyzed using smart survey software.

Results: We received 61 responses (response rate = 51%), of which 57 (n=57) were analysed. The respondents largely comprised doctors working at university hospitals (74%) and consultants (45%).

17% of the respondents were not confident with taking an inflammatory back pain history. Of the remaining 83%, only 67% respondents would normally think to ask about the main symptoms of back pain related to AS. Thus 44% respondents were not competent in taking an inflammatory back pain history.

With regards to their clinical practice, only 54% of respondents would routinely ask about symptoms of SpA in patients with recurrent AAU, even though 79% would test for HLA-B27. Furthermore, according to 42% of respondents they would never refer a recurrent AAU patient to a rheumatologist.

Most respondents felt that work pressure was the most likely reason for deficit of proper SpA assessment and referrals, followed by lack of support by guidelines.

Conclusions: Our survey demonstrates a clear knowledge deficit among ophthalmologists regarding inflammatory back pain and the importance of assessing it in recurrent AAU patients. This can be improved by educating ophthalmologists about SpA in conferences and developing local guidelines about patients with HLA B-27 recurrent AAU related to assessment and referral.

References:

- [1] A novel evidence-based detection of undiagnosed spondyloarthritis in patients presenting with acute anterior uveitis: the DUET (Dublin Uveitis Evaluation Tool).
- [1] Muhammad Haroon, Michael O'Rourke, Pathma Ramasamy, Conor C Murphy, Oliver FitzGerald.

Acknowledgements: Respondents.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2017-eular.5434

FRI0464 ARTERIAL WALL INFLAMMATION IS NOT AFFECTED BY ANTI-IL17 TREATMENT IN PATIENTS WITH PERIPHERAL SPONDYLOARTHRITIS

L.J.J. Van Mens¹, S.L. Verweij², A.W.R. van Kuijk³, E.S.G. Stroes², D.L. Baeten^{1,4}. ¹AMC, Amsterdam Immunology and Rheumatology Center; ²Vascular medicine, Academisch Medisch Centrum; ³Reade, Amsterdam Immunology and Rheumatology Center, Amsterdam, Netherlands; ⁴UCB, Brussels, Belgium

Background: Patients with spondyloarthritis (SpA), a chronic inflammatory disease, have an increased cardiovascular risk, which is partly due to increased inflammatory activity in the arterial wall.

IL-17A blockade with secukinumab is an effective treatment for SpA. The role of IL-17A in atherogenesis is controversial, some studies suggest that IL-17A is pro-atherogenic, while others indicate that IL-17A is athero-protective. So, it is not known what the effect is of treatment with IL-17A blockade on inflammatory activity in the arterial wall.

Objectives: To assess the effect of 3 months treatment with secukinumab on arterial wall inflammation in SpA patients with peripheral disease (pSpA).

Methods: We included 20 patients with clinical pSpA in a 12 week open-label trial. Treatment consisted of 300 mg secukinumab once a week during the first 4 weeks and then every 4 weeks thereafter. EULAR DAS response was used to define a responder/non responder state. To measure arterial wall inflammation we performed a 18-fluorodeoxyglucose positron emission tomography with computed tomography (¹⁸F-FDG PET/CT) imaging in 18 patients at baseline and week 12, which is a validated method to quantify arterial wall inflammation. Arterial wall inflammation is measured in both the ascending aorta and carotids, maximal FDG uptake is shown as the maximal target-to-background ratio (TBR_{max}).

Results: 18 patients with pSpA (age 44±12, 72% male) and without a previous cardiovascular event underwent imaging. Overall, three months treatment with secukinumab resulted in a significant improvement of disease activity with 17/18 patients achieving a EULAR DAS response (9 good and 8 moderate responders). Correspondingly, CRP levels decreased significantly (baseline: 3.2 [1.2–12.40] mg/dl vs. wk 12: 2.0 [1.1–5.8] mg/dl, $p=0.011$). Importantly, treatment with secukinumab did not affect cholesterol levels (total cholesterol baseline: 5.1±1 mmol/l vs wk 12: 5.5±1 mmol/l, $p=0.167$; LDL-c baseline: 3.2±0.8 mmol/l vs wk 12: 3.5±0.9 mmol/l, $p=0.219$). Additionally, arterial wall inflammation as measured by PET-CT did not change over the course of the 12 weeks treatment with secukinumab (aorta TBR_{max} baseline: 3.3±0.9 vs. wk 12: 3.3±0.7, $p=0.861$; carotid TBR_{max} baseline: 1.88±0.6 vs. wk 12: 1.76±0.4, $p=0.067$).

Conclusions: This pilot study in 18 patients with pSpA without any preexisting CV events showed that treatment with secukinumab for 3 months has no effect on arterial wall inflammation as measured by PET-CT. Further research in larger patient groups, over a longer period of treatment, and with different measurements remains warranted to fully elucidate the effect of IL-17A blockade on vascular inflammation.

Acknowledgements: This study was funded by an unrestricted grant from Novartis.

Disclosure of Interest: L. Van Mens: None declared, S. Verweij: None declared, A. van Kuijk Grant/research support from: UCB, Pfizer, MSD, Janssen, Consultant for: Novartis, Celgene, E. Stroes Speakers bureau: Amgen, Sanofi, Merck, D. Baeten Grant/research support from: Pfizer, MSD, AbbVie, Novartis, UCB, Janssen, Boehringer Ingelheim, Consultant for: Pfizer, MSD, AbbVie, Novartis, UCB, Janssen, Boehringer Ingelheim, Eli Lilly, Roche, BMS, Glenmark, Employee of: UCB

DOI: 10.1136/annrheumdis-2017-eular.4709

FRI0465 ANGLES OF SACRUM INCLINATION EFFECT ON RADIOLOGIC IMAGING READING IN SPONDYLOARTHRITIS (THE ANTELOPE-DESIR STUDY)

M. Herbert¹, L. Deloire², F. Garrigues², L. Gossec³, A. Simon⁴, A. Feydi⁵, F. de Bruin⁶, M. Reijnders⁶, D. van der Heijde⁷, D. Loeuille⁸, P. Claudepierre⁹, T. Marhadour¹, A. Saraux¹ on behalf of DESIR cohort. ¹Rheumatology; ²Radiology, CHU Brest and Université Bretagne Occidentale, Brest; ³Rheumatology, Pitié-Salpêtrière, Paris; ⁴Neurosurgery, CHU Brest and Université Bretagne Occidentale, Brest; ⁵Radiology, CHU Cochin, Paris, France; ⁶Radiology; ⁷Rheumatology, Leiden, Leiden, Netherlands; ⁸Rheumatology, CHU, Nancy; ⁹Rheumatology, CHU Creteil, Paris, France

Objectives: To assess the impact of spinal angles on clinical and imaging features of suspicion of axial spondyloarthritis (axSpA).

Methods: The DESIR cohort is a prospective longitudinal cohort study of adults aged 18–50 with inflammatory back pain (IBP) ≥3 months, ≤3 years. Baseline lateral lumbar radiography of patients included in DESIR cohort were read by two central blinded fellow readers (and a rheumatologist spine specialist in case of discrepancy) for Sacral Horizontal Angle (SHA), Lumbosacral angle (LSA) and total Lordotic Angle (TLA) measures. On the basis of literature, patients were classified depending on whether they had TLA more or less than 50°, SHA more or less than 40° or LSA more or less than 15°. Associations between angles and baseline clinical variables, presence of X-Rays (New York) and MRI (ASAS and MORPHEO proposal definition) sacroiliitis, presence of spinal signs of spondyloarthritis (mSASSS, BASRI-total, SPARCC scores), presence

of spinal degenerative MRI signs on X-rays (yes or no) and MRI (presence of Modic abnormalities, Pfirrmann score, Canal stenosis, Extrusion, High intensity zone Facet osteoarthritis) according to central reading (two readers) and axSpA diagnostic confidence (according to local clinician's confidence on a 0–10 visual analog scale) were assessed by univariate analysis using the chi-square test (or Fisher's exact test where appropriate) and the Mann-Whitney test. Adjustment for multiple testing was performed according to Bonferroni method.

Results: Of 708 patients, data were available for 677, 675 and 672 for SHA, LSA and TLA, measures with a mean value of 39.2°, 14.5° and 51.5° respectively. Clinical features and diagnostic confidence did not differ between the SHA, LSA and TLA groups. More sacroiliitis imaging, according to ASAS (41.4% versus 32.0%) and MORPHO definition (48.6% versus 39.3%), were reported in TLA <50° group but the differences did not reach statistical significance. Radiological scores were low with a mean value of 0.49 (±1.83), 0.30 (±0.78) and 4.9 (±9.0) for mSASSS, BASRI-total and SPARCC score, respectively, and no inter-group difference was found. In L5S1, more grade 3 and 4 Pfirrmann class and MODIC discopathy (types 1 and 2) were observed for SHA <40°, and TLA <50° ($p < 0.001$) whereas the difference did not reach the significance level for LSA <15° ($p = 0.05$) (table).

Level	L5/S1					
	SHA		LSA		TLA	
Angles	<40	>40	<15	>15	<50	>50
Number of patients	328	300	313	313	270	353
Pfirrmann class =3	99 (30.2)	53 (17.7) **	92 (29.3)	58 (18.5)*	83 (30.7)	68 (19.3) **
High Intensity Zone	61 (18.6)	35 (11.7)*	60 (19.2)	36 (11.5)*	40 (14.8)	56 (15.9)
Disc protrusion	37 (11.3)	22 (7.3)	27 (8.6)	30 (9.6)	34 (12.6)	24 (6.8)*
Disc extrusion	29 (8.8)	17 (5.7)	28 (8.9)	18 (5.8)	24 (8.9)	22 (6.2)
Canal stenosis	8 (2.4)	2 (0.7)	2 (0.7)	7 (2.2)	7 (2.6)	3 (0.8)
MODIC 1 signal	12 (3.7)	5 (1.7)*	15 (4.8)	2 (0.6)*	13 (4.8)	4 (1.1)*
MODIC 2 signal	7 (2.1)	1 (0.3)*	2 (0.6)	1 (0.3)*	5 (1.8)	3 (0.4)*
Facet joint osteoarthritis	1 (0.3)	2 (0.7)	1 (0.3)	2 (0.6)	1 (0.4)	2 (0.6)

Conclusions: Lumbar spine morphology is not associated with any clinical variable, presence on X-Rays or MRI of spinal signs of spondyloarthritis or sacroiliitis. At the L5S1 level, a more horizontal SHA and a reduction of TLA is associated with more degenerative radiological lumbar spine manifestations.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2017-eular.5128

FRI0466 REAL-LIFE PERFORMANCE OF THE ASAS HEALTH INDEX IN ROUTINE CARE OF PATIENTS WITH SPONDYLOARTHRITIS

U. Kiltz, T. Wiatr, X. Baraliakos, K. Fedorov, J. Braun. *Rheumatology, Rheumazentrum Ruhrgebiet, Herne, Germany*

Background: The measurement properties of the ASAS Health Index (ASAS HI) which has been developed to measure health and functioning of patients with spondyloarthritis (SpA) have been evaluated in a cohort study. Its association with structural changes has not been examined to date.

Objectives: To investigate the relationship between clinical data, function and health as assessed by the ASAS HI in patients with axial SpA (axSpA) and to study the influence of structural spinal changes on this instrument.

Methods: Patients fulfilling ASAS classification criteria for axSpA were recruited prospectively. Information was collected on clinical assessments (ASAS HI, NRS pain, BASDAI, ASDAS, BASFI, BASMI), laboratory parameters and spinal x-rays. Images were scored using mSASSS by two independent readers. The relationship between ASAS HI and other health outcomes was evaluated by Spearman correlation.

Results: A total of 150 patients (57 nr-axSpA and 93 AS patients) were included: 68.7% male, mean (SD) age 46.4 (14.1), symptom duration 18.7 (13.5) and diagnosis since 11.4 (11.8) years, and HLA-B27 positive in 82.0%. Values of clinical assessments were ASAS HI 7.4 (4.1), BASDAI 4.7 (2.3), ASDAS 2.7 (1.1), BASMI 3.3 (1.8), pain 5.7 (2.7), and BASFI 4.7 (2.6). Elevated CRP levels were found in 34.8% of the patients. Radiographs of the sacroiliac (SI) joint and the spine were available in 138 patients, 38.0% of which had syndesmophytes and 7.5% had a bamboo spine. The median (IQR) mSASSS value was 4.3 (IQR 1.0–22.1) in AS and 0.2 (IQR 0.0–1.4) in nr-axSpA patients. Patients received a treatment with NSAIDs (62.7%), DMARDs (20.9%) and/or biologics (49.4%). Significant correlation of ASAS HI was found for BASFI ($r = 0.8$), BASDAI

($r = 0.7$), ASDAS ($r = 0.5$) and BASMI ($r = 0.5$, and, all $p < 0.05$). No correlations were found for ASAS HI and radiographic damage (mSASSS $r = 0.2$, occurrence of syndesmophytes $r = 0.01$, occurrence of bamboo spine $r = 0.2$) and CRP levels ($r = 0.07$). Stratifying patients by symptom duration (cut-off 3 years) did not affect these results.

Conclusions: Established measures of function and spinal mobility correlate well with the ASAS HI in patients with axSpA. The influence of structural changes as measured by mSASSS on the ASAS HI was limited in this study, probably due to relatively low mSASSS scores at baseline. Further study in more and more severe patients is needed to study the association of physical activity, spinal mobility, function and radiographic damage in axSpA.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2017-eular.2202

FRI0467 DEVELOPMENT AND PRELIMINARY VALIDATION OF THE COMPUTED TOMOGRAPHY SACROILIAC STRUCTURAL SCORE (CT-SSS) FOR ASSESSMENT OF STRUCTURAL LESIONS IN AXIAL SPONDYLOARTHRITIS

W.P. Maksymowich^{1,2}, M. Raynal³, D. Loeuille³, M.-A. D'Agostino⁴, J. Paschke², R.G. Lambert⁵. ¹Medicine, University of Alberta; ²CaRe Arthritis, Edmonton, Canada; ³Rheumatology, CHRU de Nancy, Vandœuvre-lès-Nancy; ⁴Rheumatology, Université Paris Ouest, Paris, France; ⁵Radiology, University of Alberta, Edmonton, Canada

Background: Computed tomography (CT) is considered the imaging benchmark for the assessment of certain structural lesions in the sacroiliac joints (SIJ) of patients with axial spondyloarthritis (axSpA). Availability of low dose radiation techniques may lead to more widespread use, potentially as a structural endpoint in clinical trials research.

Objectives: We aimed to validate a new CT-based scoring method, the CT Sacroiliac Structural Score (CT-SSS), for assessing structural lesions in the SIJ.

Methods: CT scans of the SIJ from 44 patients (26 females, mean age 49.4 years, mean symptom duration 9.1 years) were reconstructed in the semicoronal plane parallel to the superior border of the sacrum and scoring of lesions was confined to this plane. Structural lesions were scored in consecutive slices in SIJ quadrants (erosion, sclerosis) or SIJ halves (ankylosis) on a dichotomous basis (present/absent) using the same anatomical principles as developed for the SPARCC MRI SIJ inflammation and structural scores. The most anterior slice is defined as visible joint ≥ 1 cm vertical height and when < 3 cm is defined as having only upper iliac and sacral quadrants. A visible joint ≥ 3 cm vertical height is defined as having 4 quadrants. At the posterior aspect of the SIJ, there is a natural separation of iliac and sacral cortical bone by structures in the ligamentary portion. Scoring is terminated when < 1 cm of iliac and sacral bone is appositional. Two readers independently scored CT scans without a prior calibration exercise and using direct online data entry onto a schematic of the SIJ. Reliability was assessed by kappa statistics, intra-class correlation coefficient (ICC), and Bland-Altman limits of agreement.

Results: Scoring was feasible (5–10 minutes per scan) and both ankylosis (ICC=0.95) and erosion (ICC=0.81) were reliably scored (Table). Sclerosis was less reliably scored (ICC=0.39). Presence/absence of ankylosis was reliably detected irrespective of whether this was based on a single slice ($\kappa = 0.77$) or 3 consecutive slices ($\kappa = 0.81$). Reliable detection was lower for erosion ($\kappa = 0.50$ for 1 or 3 slices) and sclerosis ($\kappa = 0.44$ and 0.48 for 1 and 3 slices, respectively). Bland-Altman graphs illustrate reliability across the range of scores for ankylosis and erosion.

Table 1. Descriptive and Reliability data for CT-SSS score

	Mean (SD) score		Median (IQR) score		ICC	95% L of A
	R1	R2	R1	R2		
Erosion	4.4 (9.7)	6.5 (11.5)	0 (25.5)	0 (21)	0.95	-10.7, 13.5
Ankylosis	7.5 (14.1)	6 (13.6)	0 (12)	0 (12)	0.81	-16.2, 12.2
Sclerosis	6.6 (14.4)	7.4 (18.2)	0 (20)	0 (24.5)	0.39	-34.5, 33.0

IQR interquartile range L of A Limits of Agreement.

Conclusions: The CT-SSS method is feasible and reliable for scoring ankylosis and erosion with minimal calibration. Sclerosis requires further standardization and calibration.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2017-eular.5254

FRI0468 RADIOGRAPHIC HIP INVOLVEMENT IN PATIENTS WITH ANKYLOSING SPONDYLITIS. A STUDY OF ITS PREVALENCE AND DETERMINING FACTORS

X. Michelena, M. López de Recalde, H. Borrell, J. Lluch, P. Juárez, I. Morales, M. Pascual, J.-M. Nolla, X. Juanola. *Department of Rheumatology, Hospital Universitari de Bellvitge, L'Hospitalet de Llobregat, Barcelona, Spain*

Background: Hip involvement in Ankylosing Spondylitis (AS) is the most frequent extraspinal manifestation and a common cause of disability and limited mobility, often ending with hip replacement. There are few studies analysing the radiographic changes of the hip and their relationship with other disease variables.