secondary osteoporosis was detected. Serum vitamin D concentration was <30 ng/ml in 5 patients, including 2 with concentration <10 ng/ml.

Conclusions: Our study reveals that odontoid fractures mainly occur in elderly osteoporotic patients after a low energy impact. Although WHO osteoporosis definition excludes cervical fractures, odontoid fracture may be considered as an osteoporotic fracture. Further studies are required to confirm these results.

References:

- [1] Watanabe M, et al. Analysis of predisposing factors in elderly people with type II odontoid fracture. *Spine J.* 2014 Jun 1; 14(6):861–6.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2017-eular.3503

FRI0579 INFLUENCE OF ORAL PREDNISOLONE ON EFFECT OF DENOSUMAB ON OSTEOPOROSIS IN PATIENTS WITH JAPANESE RHEUMATOID ARTHRITIS; 24 MONTHS OF FOLLOW-UP ~A MULTICENTER REGISTRY STUDY~

Y. Kanayama¹, Y. Hirano², N. Takahashi³, S. Asai³, N. Ishiguro³, T. Kojima³ on behalf of TBCR-BONE study group. ¹Orthopedic Surgery and Rheumatology, Toyota Kosei Hospital, Toyota; ²Rheumatology, Toyohashi Municipal Hospital, Toyohashi; ³Orthopedic Surgery, Nagoya University Graduate School of Medicine, Nagoya, Japan

Background: Denosumab (DMB) is a fully human monoclonal antibody to the RANKL that blocks its binding RANK, inhibiting the development and activity of osteoclasts, decreasing bone resorption, increasing bone density and reducing fracture risk. Osteoporosis (OP) is more frequent in patients with rheumatoid arthritis (RA) than in the general population due to active systemic inflammation as well as the use of glucocorticoid and immobility. However efficacy of DMB is not clear in patients with glucocorticoid-induced OP in RA. Therefore we investigated the influence of oral prednisolone on effect of DMB in patients with Japanese RA from initiation to 24 months at this time.

Objectives: This prospective study investigated the efficacy of DMB for 24 months on glucocorticoid-induced OP in RA patients.

Methods: Patients with a diagnosis of RA according to the 2010 ACR/EULAR criteria who had been prescribed DMB from Tsurumi Biologics Communication Registry (TBCR)-BONE between October 2013 and October 2015 were enrolled. The final study cohort of 63 patients received continuous DMB therapy more than 24 months. The DMB dose was 60mg at once every 6 months. In all cases native or activated vitamin D has been used. We reviewed the results for 24 months about the increase and decrease of bone mineral density (BMD) of lumbar spine (LS) and total hip (TH) by DEXA and bone turnover markers, intact n-terminal propeptide type I procollagen (PINP) and tartrate-resistant acid phosphatase form 5b (TRACP-5b).

Results: In the patients receiving oral prednisolone group (n=23, GC+) and not receiving group (n=40, GC-), the number of female was each 21 (91%) and 39 (98%) cases (p=0.548). The mean age was 69.8±7.0 and 71.0±7.3 years old (p=0.622); disease duration was 16.0±8.9 and 15.7±12.6 years (p=0.592); the body mass index was 20.7±3.5 and 19.7±3.0 (p=0.335) and the FRAX was 34.2±19.1 and 24.6±13.9 (p=0.040). Clinical findings related to RA and OP at baseline were as follows; CRP 1.2±1.4 and 0.5±1.0 mg/dl (p=0.009); DAS-CRP 3.16±1.17 and 2.50±1.27 (p=0.025); m-HAQ 1.24±0.90 and 0.82±0.82 (p=0.065); PINP 60.1±38.6 and 56.9±33.5 µg/l (p=0.711); TRACP-5b 513±257 and 505±202 mU/dL (p=0.689); LS-BMD 0.87±0.18 and 0.80±0.14 g/cm² (p=0.074) and TH-BMD 0.60±0.11 and 0.59±0.08 g/cm² (p=0.457). The rate of decreased PINP from baseline to 6, 12, 18 and 24 months were each -25.6% vs -41.6% (p=0.129) at 6 month, -8.0% vs -42.6% (p=0.031) at 12 month, -19.5% vs -27.5% (p=0.235) at 18 month and -13.8% vs -33.8% (p=0.134) at 24 month and TRACP-5b were -26.5% vs -38.7% (p=0.710) at 6 month, -22.0% vs -35.1% (p=0.229) at 12 month, -25.2% vs -28.1% (p=0.792) at 18 month and -20.6% vs -27.3% (p=0.663) at 24 month in the GS+ vs GS- group. The rate of increased LS-BMD from baseline to 6, 12, 18 and 24 months were each 4.1% vs 4.6% (p=0.671) at 6 month, 5.3% vs 7.0% (p=0.361) at 12 month, 8.1% vs 8.1% (p=0.438) at 18 month and 10.0% vs 7.7% (p=0.104) at 24 month and TH-BMD were 3.3% vs 2.7% (p=0.690) at 6 month, 1.4% vs 3.8% (p=0.804) at 12 month, 6.4% vs 4.0% (p=0.259) at 18 month and 5.7% vs 4.3% (p=0.945) at 24 month in the GS+ vs GS- group (Fig.1,2).

Conclusions: DMB was effective in OP of RA patients. Oral prednisolone use did not influence the efficacy of DMB for 24 months.

Disclosure of Interest: Y. Kanayama: None declared, Y. Hirano Speakers bureau:

Abbvie Japan, Eisai, Mitsubishi Tanabe Pharma, Pfizer, Chugai Pharmaceutical, and Bristol-Myers Squibb, N. Takahashi Speakers bureau: bbvie Japan Co. Ltd, Eisai Co. Ltd, UCB Japan Co. Ltd, Mitsubishi Tanabe Pharma Corporation, Takeda Pharmaceutical Company Ltd, Pfizer Co. Ltd, Chugai Pharmaceutical Co. Ltd, Janssen Pharmaceutical K.K., and Bristol-Myers Squibb Co. Ltd, S. Asai: None declared, N. Ishiguro Grant/research support from: Daiichi Sankyo, Takeda Pharmaceutical, Hisamitsu Pharmaceutical, Otsuka Pharmaceutical, Taisho Toyama Pharmaceutical, Kaken Pharmaceutical, Eisai, Janssen Pharmaceutical, Bristol-Myers Squibb, AbbVie, ChugaiPharmaceutical, Mitsubishi Tanabe Pharmaceutical, Astellas Pharma and Pfizer Japan, T. Kojima Speakers bureau: Mitsubishi Tanabe Pharma, Takeda Pharma, Eisai Pharma, AbbVie, Bristol-Myers Squibb, and Pfizer, Janssen Pharmaceutical Companies, Astellas Pharma and Chugai Pharma

DOI: 10.1136/annrheumdis-2017-eular.4677

FRI0580 PREDICT SARCOPENIA BY SONOELASTOGRAPHY OF QUADRICEPS MUSCLE IN OSTEOPOROTIC PATIENTS

Y.-C. Chen, C.-H. Ko. Rheumatology, Kaohsiung Chang Gung Memorial Hospital, Kaohsiung, Taiwan, Province of China

Background: Reduced muscle mass had associated with high mortality. So it is urgent for simple techniques to early detection sarcopenia. Our objective was to examine the validity of sonoelastography to predict sarcopenia in osteoporotic patients.

Objectives: To evaluate the association of sonoelastography and dual-energy X-ray absorptiometry in patients with sarcopenia and osteoporosis.

Methods: We conducted an observational study in Kaohsiung Chang Gung Memorial Hospital. Sarcopenia was determined using a dual-energy X-ray absorptiometry. Osteoporosis was defined through estimated bone mass (BM). Sonoelastography was performed over mid thigh over quadriceps muscle. We measure hardness and elastography ratio of quadriceps over subcutaneous fat tissue. ROC analysis was used to find best cut-off point.

Results: A total 47 (23 sarcopenia, 24 non-sarcopenia) osteoporotic patients were enrolled. The mean age was 71.04±9.64 years, and most patients (88.9%) were women. Sonoelastography showed sarcopenia patients had more soft than non-sarcopenia patients, furthermore the elastography ratio of quadriceps over subcutaneous tissue was lower than non-sarcopenia patients. When the cut points determined by receiver operating characteristic (ROC) curve analysis were applied, The best cut-point of hardness was 42.5 (sensitivity, 0.969; 1-specificity, 0.066), while the best cut-point of quadriceps over subcutaneous tissue was 70.5% (sensitivity, 1.00; 1-specificity, 0.079).

Characteristics of study patients

Variables	Sarcopenia (n=23)	Non-sarcopenia (n=24)	P-value
Age (years)	71.04 ±9.64	70.82±8.624	0.316
Body mass index (kg/m ²)	23.80±3.45	23.83±4.56	0.977
Gender (Female %)	22 (75.9%)	78 (84.8%)	0.624
EX2/1	0.37±0.15	1.36±0.57	<0.001
Hard%	15.44±13.15	76.70±20.12	<0.001

Conclusions: Sonoelastography was easily applicable in patients with sarcopenia and osteoporosis. Using hardness content and elastography ratio of quadriceps over subcutaneous ratio render more information of muscle character. Early detection of sarcopenia with sonoelastography in patients with osteoporosis afford future trend of preventive medicine in geriatric patients.

References:

- [1] Teber MA, Ogur T, Bozkurt A, Er B, Turan A, Gulbay M, et al. Real-time sonoelastography of the quadriceps tendon in patients undergoing chronic hemodialysis. *J Ultrasound Med.* 2015;34(4):671–7.

Acknowledgements: This study was supported by Kaohsiung Chang Gung Memorial Hospital and Chang Gung University College of Medicine, Kaohsiung (grant CMRPG8F1731).

Disclosure of Interest: None declared
DOI: 10.1136/annrheumdis-2017-eular.2709

FRI0581 HEPATITIS B INCREASE MORTALITY IN PATIENTS WITH OSTEOPOROTIC VERTEBRAL FRACTURES

Y.-C. Chen, C.-H. Ko. *Rheumatology, Kaohsiung Chang Gung Memorial Hospital, Kaohsiung, Taiwan, Province of China*

Background: Hepatitis B virus (HBV) is a major cause of chronic liver diseases worldwide, particularly cirrhosis and hepatocellular carcinoma, and the most important liver disease in Taiwan. However, little is known the impact of hepatitis b on osteoporotic patients.

Objectives: This study aimed to determine if hepatitis b can increase the risk of mortality in patients with osteoporotic vertebral fracture.

Methods: This retrospective study reviewed of cases of osteoporosis patients with acute vertebral fractures between 2001 and 2008. All associated co-morbidities were recorded. Kaplan-Meier and cox regression analysis were performed.

Results: There were 432 patients with acute vertebral fractures. The mean age of 72.85±9.28. 31 (7.2%) patients had chronic hepatitis b. Using the Kaplan-Meier curve, hepatitis b had a significant effect on mortality ($p<0.001$, by log rank test). After adjusting for potential confounders, patients with hepatitis b still had a high mortality rate ($p=0.019$; HR: 2.436~5.136) 2.137, 95% CI: 1.156~5.136). The mortality rate also increased among smokers ($p=0.026$; HR: 3.6043.891; 95% CI: 1.056~12.301).

Table 1. Multivariable Cox regression analysis of the hazard ratios for adjacent fracture

Variables	Regression coefficient	P value	HR (95CI)
Gender	-0.127	0.764	0.88 (0.383–0.022)
Age	0.004	0.785	1.004 (0.976–1.032)
Smoking	1.282	0.041	3.604 (1.056–12.301)
Alcohol consumption	-0.701	0.396	0.496 (0.098–2.508)
Body mass index (kg/m ²)	0.024	0.446	1.024 (0.964–1.088)
Hepatitis B	0.891	0.019	2.436 (1.156–5.136)
Diabetes	-0.185	0.538	0.831 (0.461–1.498)
Hypertension	-0.271	0.31	0.762 (0.451–1.288)
Stroke	-0.696	0.248	0.499 (0.153–1.625)
Kidney disease	0.809	0.145	2.246 (0.758–6.656)
Cardiovascular disease	0.037	0.946	1.038 (0.351–3.069)
Pulmonary disease	-0.971	0.349	0.379 (0.050–2.883)

Abbreviations: HR, hazard ratio; SE, standard error.

Conclusions: In this study, hepatitis b increase mortality in osteoporotic vertebral fracture patients. So we should pay attention to this group in osteoporotic management.

References:

[1] Chong-Shan Wang et al. *Am. J. Trop. Med. Hyg.*, 66(4), 2002, pp. 389–393.

Acknowledgements: We thank Kaohsiung Chang Gang Memorial Hospital for data support.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2017-eular.3614

FRIDAY, 16 JUNE 2017

Other orphan diseases

FRI0582 OPEN-LABEL, MULTICENTER, DOSE-ESCALATING PHASE II CLINICAL TRIAL ON THE SAFETY AND EFFICACY OF TADEKINIG ALPHA IN ADULT ONSET STILL'S DISEASE

C. Gabay¹, B. Fautrel², J. Rech³, F. Spertini⁴, E. Feist⁵, I. Koetter⁶, E. Hachulla⁷, J. Morel⁸, T. Schaevebeke⁹, M.A. Hamidou¹⁰, T. Martin¹¹, B. Hellmich¹², P. Lamprecht¹³, H. Schultze-Koops¹⁴, A. Sleight¹⁵, E. Schiffrin¹⁵.
¹Rheumatology, University Hospitals of Geneva, Geneva, Switzerland; ²Rheumatology, Pitié Salpêtrière University Hospital, Paris, France; ³Department of Medicine 3, Rheumatology and Immunology, Universitätsklinikum Erlangen, Erlangen, Germany; ⁴Division of Immunology and Allergy, CHUV, Lausanne, Switzerland; ⁵Rheumatology, Charité Universitätsmedizin, Berlin; ⁶Asklepios Klinikum, Hamburg, Germany; ⁷CHRU de Lille - Hôpital Claude Huriez, Lille; ⁸Department of Rheumatology, University and CHU of Montpellier, Montpellier; ⁹Rheumatology, Hôpital Pellegrin, Bordeaux; ¹⁰CHU de Nantes, Nantes; ¹¹University Hospital of Strasbourg, Strasbourg, France; ¹²Klinik für Innere Medizin, Rheumatologie und Immunologie, Medius Klinik, Kirchheim unter Teck; ¹³Rheumatology, University of Lübeck, Lübeck; ¹⁴Medizinische Klinik und Poliklinik IV, LMU, München, Germany; ¹⁵Ab2 Bio, Lausanne, Switzerland

Background: Adult Onset Still's Disease (AOSD) is a severe systemic autoinflammatory disease. Elevated interleukin (IL)-18 levels correlate with AOSD disease activity, suggesting a central role

Objectives: To examine the safety and efficacy of IL-18 blockade in AOSD with the administration of Tadekinig alpha, a recombinant human IL-18 binding protein

Methods: Patients (Pts) with AOSD according to Yamaguchi criteria were included if they had been previously treated for at least one month with corticosteroids (CS). The pts received either 80 mg (group 1) or 160 mg (group 2) Tadekinig alpha sc three times weekly for 12 weeks. Oral CS and/or conventional DMARDs were allowed at stable dose in combination with Tadekinig alpha. After 3 wks, group 1 pts qualified as non responders (CRP levels did not decrease by $\geq 50\%$ from baseline values or absence of temperature normalization) could be up-dosed to 160 mg for an additional 12 weeks. The primary endpoint was safety and secondary endpoints included early signs of efficacy

Results: 23 pts were included (10 and 13 in group 1 and group 2, respectively). 6 pts of group 1 were up-dosed to 160 mg. Baseline characteristics and safety and efficacy results are described in Table 1. Most adverse events (AEs) (47) were considered as mild or moderate. Three serious AEs (SAE) were reported including two that were considered not related to the study drug and one possibly related according to the investigator (toxic optic neuropathy). Four premature discontinuations were related to AEs, including 3 cases of injection site reactions and 1 SAE. Systemic response as defined by $\geq 70\%$ decrease of serum C-reactive protein (CRP) or normalization of CRP and ferritin levels was obtained in 2/10 and 6/13 pts of groups 1 and 2, respectively in the ITT analysis, and in 2/9 and 6/9 pts of groups 1 and 2, respectively, in the PP analysis.

Table 1. Baseline characteristics and safety and efficacy results

	Group 1 (80 mg) (N=10)	Group 2 (160 mg) (N=13)
Age, median (IQR)	49.5 (34.3–58.8)	35 (30–58)
Median disease duration, months (IQR)	25.5 (8.7–44.2)	11.6 (2.1–37.6)
Pts with previous conventional DMARDs	6	7
Pts with previous biological DMARDs	4	5
Pts with SAEs	2	1
Discontinuations due to AEs*	1	3
Discontinuations due to lack of response*	1	2
Arthritis improvement ($\geq 20\%$ reduction in tender and swollen joint count)**	3/9	4/10
Skin rash resolution**	6/7	2/5
CRP level reduction by $\geq 70\%$ **	2/10	6/13
Normalization of ferritin levels**	0/7	1/5

*Number of Pts, **Denominator corresponds to Pts with clinical manifestations or abnormal values at baseline.

Conclusions: IL-18 blockade by Tadekinig alpha has an acceptable safety profile and is efficacious, in particular at the 160 mg dosage, in patients with refractory AOSD

Disclosure of Interest: C. Gabay Grant/research support from: AB2 Bio, Roche, Pfizer, Consultant for: AB2 Bio, Roche, Pfizer, MSD, BMS, AbbVie, Sanofi, B. Fautrel: None declared, J. Rech: None declared, F. Spertini: None declared, E. Feist: None declared, I. Koetter: None declared, E. Hachulla: None declared, J. Morel: None declared, T. Schaevebeke: None declared, M. Hamidou: None declared, T. Martin: None declared, B. Hellmich: None declared, P. Lamprecht: None declared, H. Schultze-Koops: None declared, A. Sleight Employee of: Ab2 Bio, E. Schiffrin Employee of: Ab2 Bio

DOI: 10.1136/annrheumdis-2017-eular.2336