count and/or ultrasounds) were enrolled in this 5-years prospective observational study. Clinical assessments as well as joint ultrasound (US) and MRI were performed at baseline and after 5 years. Each patient underwent a non-dominant hand-wrist US examination using a Logiq 9 with a linear probe operating at 14 MHz. Synovitis was defined as the presence of synovial hypertrophy and/or the presence of power Doppler signal (PD). A non-dominant hand-wrist MRI study with a 0.3 T extremity dedicated machine to evaluate the presence of bone erosions (BE) and bone marrow edema (BME) was also performed. Coronal and axial T1-weighted gradient-echo images and coronal STIR images were acquired. The images acquired were scored according to the RAMRIS scoring system for RA by a trained radiologist unaware of the clinical picture and diagnosis.

Results: Fifty-seven pts were enrolled (female 91.2%, mean age 44±12.2 years, mean disease duration 15.9±9 years);43 (75.4%) completed the follow-up, 3 died (5.2%) and 11 (19.4%) were lost to follow-up. At baseline, 7 (12.3%)satisfied criteria for Jaccoud arthropathy (JA) and 7 (12.3%) had a recent onset arthritis (<1 year of duration). 22 pts (28.6%) showed clinical signs of synovitis, 56 (98.2%) presented positive hand-wrists US (synovitis) and in 14 (24.6%) PD signal was also recorded. Six pts (11.76%) already showed erosions at standard X-Ray, while MRI revealed at least one BE in 30 and 54 patients respectively, for a cumulative mean erosive burden of 9.2 erosions (range 1–63). After 5 years of follow-up, 34 pts consented to repeat the assessment; 11 (33%) had JA and 18 (29%) were still presenting clinical signs of synovitis; 28 pts (82.3%) showed synovitis at US with PD in 7 cases (20.5%). The final mean erosive burden resulted 12.3 (range 2–82) with a significant increase from the baseline evaluation (p=0.001). Overall, 16 pts accrued joint damage. Interestingly, erosion progression was observed also in 12 pts with negative joint count but positive US at baseline. The presence of PD at US and BME at baseline was associated with higher erosive burden at follow-up (p=0.03 and p=0.02 respectively)

Conclusions: Arthritis in SLE can be persistent over time and progress to joint damage even in a short tem period despite treatment; normal joint count at physical examination but US findings of synovitis can be associated with erosion progression over time. The presence of PD at US and bone marrow edema at MRI are associated with a more severe damage progression. US and MRI can be a valid help for the clinician to identify patients at higher risk of severe damage to be treated with a more aggressive therapeutic strategy targeting arthritis.

Disclosure of Interest: None declared
DOI: 10.1136/annrheumdis-2017-eular.6016

FRI0641 WRIST INFLAMMATION AS ASSESSED BY MAGNETIC RESONANCE IMAGING IS ASSOCIATED WITH PATIENT-REPORTED PHYSICAL IMPAIRMENT, GLOBAL DISEASE ACTIVITY AND PAIN IN EARLY RHEUMATOID ARTHRITIS: LONG-TERM RESULTS FROM TWO RANDOMIZED CONTROLLED TRIALS

D. Glinatsi¹, J.F. Baker², M.L. Hetland^{1,3}, K. Hørslev-Petersen^{4,5}, B.J. Eijbjerg⁶, K. Stengaard-Pedersen⁷, P. Junker⁸, T. Ellingsen⁸, H.M. Lindegaard⁸, I. Hansen⁷, T. Lottenburger⁹, J.M. Møller¹⁰, L. Ørnberg¹, A. Vestergaard¹¹, A.G. Jurik^{7,12}, H.S. Thomsen^{3,10}, T. Torfing⁸, S.M. Bisgaard¹, M.B. Axelsen¹, M. Østergaard^{1,3}. ¹Rigshospitalet, Glostrup, Denmark; ²Corporal Michael J. Crescenz VA Medical Center, Philadelphia, PA, United States; ³University of Copenhagen, Copenhagen; ⁴King Christian 10th Hospital for Rheumatic Diseases, Gråsten; ⁵University of Southern Denmark, Odense; ⁶Slagelse Hospital, Slagelse; ⁷Aarhus University Hospital, Aarhus; ⁸Odense University Hospital, Odense; ⁹Vejle Hospital, SLB, Vejle; ¹⁰Copenhagen University Hospital Herlev, Herlev; ¹¹Hvidovre University Hospital, Hvidovre; ¹²Aarhus University, Aarhus, Denmark

Background: Studies in established rheumatoid arthritis (RA) have shown that radiographic progression is associated with increasing health assessment questionnaire (HAQ) score. However, most studies have failed to demonstrate this association at the early stage of the disease. In addition, little is known about how specific pathologies, e.g. joint inflammation, tenosynovitis and joint damage, contribute to different patient-reported outcomes (PROs).

Objectives: To examine the association between MRI wrist inflammation and damage with PROs in patients with early RA.

Methods: MRIs of the wrist and hand were obtained from 210 early RA patients participating in two investigator-initiated, randomized, controlled studies (CIMESTRA/OPERA), which aimed at achieving inflammatory control by use of conventional and/or biologic drugs combined with intra-articular injection of glucocorticoids. The image-sets were assessed according to the RA MRI scoring system (RAMRIS) for inflammation (synovitis, tenosynovitis, oostetis) and damage (bone erosions, joint space narrowing) at baseline (n=210), 1 (n=206) and 5 (n=113) years follow-up. Data from the two studies were pooled and assessed for associations between MRI features and HAQ, patient global visual analogue scales (VAS-PtGlobal) and VAS pain using Spearman correlation for status and change scores, univariate and multivariable linear regression analyses for change scores and generalized estimating equations for status and change scores. MRI features were further tested for trends against specific hand-related HAQ questions using the Jonckheere trend test.

Results: MRI inflammation, but not damage, showed statistically significant associations with HAQ, VAS-PtGlobal and VAS pain for status and change scores, independently of swollen joint count and level of C-reactive protein.

Synovitis and tenosynovitis were the MRI features most consistently associated with PROs, particularly VAS-PtGlobal and VAS pain (Table 1). MRI synovitis and tenosynovitis mean scores increased with the level of difficulty to cut meat and open a milk carton (p<0.01), and similar patterns were seen for other hand-related HAQ items. As an additional analysis, the metacarpophalangeal joints were included in the analyses, but this did not strengthen the associations between MRI pathology and PROs.

Table 1 Univariate and multivariable generalized estimating equations

	Status				Change			
	Univariate		Multivariable		Univariate		Multivariable	
	B (95% CI)	p	B (95% CI)	p	B (95% CI)	p	B (95% CI)	p
HAQ*								
MRI synovitis	0.03 (0.01;0.05)	0.01	-0.01 (-0.03;0.01)	0.48	0.08 (0.04;0.12)	<0.001	-0.01 (-0.04;0.03)	0.72
MRI tenosynovitis	0.02 (0.00;0.05)	0.05	0.02 (0.00;0.03)	0.04	0.07 (0.04;0.09)	<0.001	0.02 (0.00;0.03)	0.05
MRI oostetis	0.01 (0.01;0.02)	<0.001	0.01 (0.00;0.02)	0.01	0.02 (0.01;0.04)	0.001	0.00 (0.00;0.02)	0.04
MRI erosion	0.01 (-0.10;0.09)	0.96			0.04 (-0.13;0.21)	0.66		
MRI JSN	0.07 (-0.01;0.15)	0.08			-0.05 (-0.25;0.16)	0.64		
VAS-PtGlobal*								
MRI synovitis	0.05 (0.01;0.09)	0.01	0.00 (-0.04;0.04)	0.89	4.13 (2.55;5.71)	<0.001	1.79 (0.34;3.23)	0.02
MRI tenosynovitis	0.04 (0.01;0.07)	0.01	0.03 (0.00;0.06)	0.06	2.12 (1.14;3.09)	<0.001	-0.08 (-1.08;0.92)	0.88
MRI oostetis	0.01 (0.00;0.02)	0.10	0.00 (-0.01;0.01)	0.84	0.90 (0.22;1.58)	0.01	0.21 (-0.25;0.66)	0.37
MRI erosion	0.01 (-0.14;0.16)	0.92			-1.55 (-8.80;5.71)	0.68		
MRI JSN	0.09 (-0.07;0.24)	0.27			-2.77 (-10.78;5.24)	0.50		
VAS pain*								
MRI synovitis	0.05 (0.01;0.08)	0.01	0.00 (-0.04;0.04)	0.97	4.84 (3.20;6.08)	<0.001	2.20 (0.87;3.53)	0.001
MRI tenosynovitis	0.04 (0.01;0.07)	0.003	0.03 (0.00;0.06)	0.02	2.19 (1.30;3.09)	<0.001	-0.02 (-0.91;3.53)	0.96
MRI oostetis	0.01 (0.00;0.02)	0.16			0.89 (0.21;1.57)	0.01	0.21 (-0.27;0.68)	0.39
MRI erosion	-0.03 (-0.18;0.12)	0.69			-2.33 (-9.18;4.51)	0.50		
MRI JSN	0.08 (-0.08;0.23)	0.33			-1.24 (-9.09;6.61)	0.76		

Association between magnetic resonance imaging (MRI) parameters and patient-reported outcomes for status scores and change scores. All generalized estimating equations (GEE) were adjusted for age and sex. MRI parameters with a univariate p-value ≤0.10 were included in multivariable GEE analysis where C-reactive protein levels and swollen joint counts were incorporated. *Log-transformed for status scores. Abbreviations: HAQ: Health assessment questionnaire, JSN: Joint space narrowing, VAS: Visual analogue scale, PtGlobal: Patient global.

Conclusions: MRI-assessed inflammation, but not damage, in the early RA wrist is associated with patient-reported physical impairment, global disease activity and pain, and the amount of wrist inflammation influences physical hand function.

Disclosure of Interest: D. Glinatsi: None declared, J. Baker: None declared, M. Hetland: Grant/research support from: BMS, AbbVie, Pfizer, UCB-Nordic, MSD, K. Hørslev-Petersen: Consultant for: AbbVie and UCB, B. Eijbjerg: None declared, K. Stengaard-Pedersen: None declared, P. Junker: None declared, T. Ellingsen: None declared, H. Lindegaard: None declared, I. Hansen: None declared, T. Lottenburger: None declared, J. Møller: None declared, L. Ørnberg: None declared, A. Vestergaard: None declared, A. G. Jurik: None declared, H. Thomsen: None declared, T. Torfing: None declared, S. Bisgaard: None declared, M. Axelsen: Grant/research support from: Abbvie, M. Østergaard: Grant/research support from: Abbvie, BMS, Janssen and Merck, Consultant for: Abbvie, BMS, Boehringer-Ingelheim, Celgene, Eli-Lilly, Centocor, GSK, Hospira, Janssen, Merck, Mundipharma, Novartis, Novo, Orion, Pfizer, Regeneron, Schering-Plough, Roche, Takeda, UCB, and Wyeth

DOI: 10.1136/annrheumdis-2017-eular.1555

FRI0642 THE POTENTIAL VALUE OF POSITRON EMISSION TOMOGRAPHY (PET)-SCAN IN SYSTEMIC SCLEROSIS FOR THE QUANTITATIVE ASSESSMENT OF INTERSTITIAL LUNG DISEASE

D.M. Peelen¹, B. Zwezerijnen², E.J. Nossent³, L.J. Meijboom⁴, O.S. Hoekstra², C.J. van der Laken¹, A.E. Voskuyl¹. ¹Rheumatology; ²Nuclear Medicine; ³Pulmonology; ⁴Radiology, VU University Medical Center, Amsterdam, Netherlands

Background: Interstitial lung disease (ILD) in systemic sclerosis is treated by immunosuppressive drugs (e.g. cyclophosphamide), aimed at reduction of inflammatory response. Differentiation between inflamed and non-inflamed fibrotic tissue might help to develop treatment stratification, with the aim of improving the prognosis of (subgroups of) SSc-ILD patients. ¹⁸F-Fluoro-Dexoxyglucose Positron Emission Tomography (¹⁸F-FDG PET) scan might be a promising tool to detect inflamed lung areas, as formerly shown in a semi-quantitative setting.[1, 2]

Objectives: This study aims to investigate the potential role of ¹⁸F-FDG PET –scan for the quantitative assessment of metabolically active SSc related ILD.

Methods: ¹⁸F-FDG PET –scans of 22 patients with systemic auto-immune disease, including 9 with SSc, 9 with systemic lupus erythematosus (SLE) and 4 with primary Sjögren's syndrome (pSS), were retrospectively analyzed. FDG-uptake was quantitatively measured within 2cm-sized Regions of Interest (ROI's) at apical, medial and basal lung levels. A total of 22 ROI's were drawn in each patient. SUVmean values of all ROI's were corrected by the medial SUVmean bloodpool value. Subsequently, the average of 6 posterior basal SUVmean values