SAT0127 UNFAVORABLE BODY COMPOSITION ALREADY AT THE ONSET OF CLINICAL ARTHRITIS

S.A. Turk¹, D. van Schaardenburg^{1,2}, M. Boers³, S. de Boer¹, C. Fokker¹, W.F. Lems^{1,3}, M.T. Nurmohamed^{1,3}, ¹Amsterdam Rheumatology and immunology Center/ Reade; ²Amsterdam Rheumatology and immunology Center/ Academic Medical Center; ³Amsterdam Rheumatology and immunology Center/ VU University Medical Center, Amsterdam, Netherlands

Background: Rheumatoid arthritis is associated with an increased cardiovascular (CV) risk. There is mounting evidence that inflammation is involved in the pathogenesis thereof (1). An unfavorable body composition is, independently, associated with an increased CV risk, and often present in established arthritis patients. This unfavorable composition is a loss of muscle mass (sarcopenia), in the presence of a stable or even increased (abdominal) fat mass (sarcopenic obesity) (2).

Objectives: Currently it is unknown when this unfavorable body composition develops. Therefore, we compared body composition in DMARD-naïve early arthritis patients with non-arthritis controls and explored the association with disease activity and traditional CV risk factors.

Methods: A total of 317 consecutive early arthritis patients and 1268 age, gender and ethnicity matched non-arthritis controls (3) had a Dual-energy X-ray absorptiometry scan to assess lean (muscle) mass index (LMI), fat mass index (FMI), fat mass distribution (arms and legs versus trunk) and android to gynoid fat mass ratio. For the obesity definition, the cut offs of Gallaher et al. were applied. For the sarcopenia definition, cut offs based on the control group were applied (mean minus two times SD). In addition, a disease activity score, erythrocyte sedimentation rate, lipid profile (total cholesterol, HDL, LDL and triglycerides) and blood pressure assessments were done.

Results: Early arthritis patients (84% fulfilling the 2010 ACR/EULAR criteria) had a significantly lower mean LMI ($p<0.01$) compared with controls. Female patients had a higher mean FMI ($p<0.01$) and more fat was distributed to the trunk in patients (females $p<0.01$, males $p=0.07$) (table). The prevalence of an unfavorable body composition (i.e. sarcopenia and sarcopenic obesity) was higher than in controls, for females 5.0% versus 1.3%, OR: 4.2, $p<0.01$ and for males 8.2% versus 1.5%, OR: 5.7, $p<0.01$ (figure). A higher FMI, more android fat and more fat distributed to the trunk were associated with higher blood pressure and lipid levels (and lower HDL levels). There was no clear relationship between body composition parameters and disease activity.

Table 1. Body composition of early arthritis patients and non-arthritis controls

	Arthritis females	Control females	Arthritis males	Control males
Fat mass index (mean, SD)	27.6 (5.3)	26.6 (4.6)*	26.8 (4.4)	27.4 (3.6)
% of fat distributed to the trunk (mean, SD)	52.2 (6.3)	49.4 (6.5)*	58.8 (6.4)	57.6 (5.6)
Android to gynoid fat mass ratio (mean, SD)	0.5 (0.2)	0.5 (0.2)	0.7 (0.2)	0.8 (0.2)*
Lean mass index (mean, SD)	6.6 (0.9)	6.7 (0.8)*	8.0 (1.1)	8.6 (1.0)*
Obese (n,%)	99 (45.2)	334 (38.4)	50 (52.1)	222 (57.7)
Sarcopenic (n,%)	11 (5.0)	11 (1.3)*	8 (8.2)	6 (1.5)*
Sarcopenic obesity (n,%)	2 (0.9)	2 (0.2)	4 (4.2)	3 (0.8)*

n = number, SD = standard deviation, % = percentages *p-value <0.05 between arthritis and non-arthritis controls.

Figure. Prevalence of sarcopenia and sarcopenic obesity in early arthritis patients compared with non-arthritis controls.

Conclusions: Patients at the clinical onset of arthritis more often have an unfavorable body composition (sarcopenia and sarcopenic obesity) than non-arthritis controls. An unfavorable body composition was associated with higher blood pressure and lipid levels (and lower HDL levels).

References:

- [1] Curr Vasc Pharmacol 2010; 8(2):285–292.
- [2] Age (Dordr) 2015; 37(6):121.
- [3] Eur J Epidemiol 2015; 30(8):661–708.

Disclosure of Interest: None declared
DOI: 10.1136/annrheumdis-2017-eular.2174

SAT0128 CLINICAL FEATURES AND EVOLUTION OF PULMONARY FUNCTION IN A SINGLE-CENTER COHORT OF PATIENTS WITH RHEUMATOID ARTHRITIS RELATED INTERSTITIAL LUNG DISEASE

S.C. Rodriguez-García, R.A. Castellanos-Moreira, M.V. Hernandez-Miguel, J. Ramirez, A. Cuervo, J. Cañete, R. Sanmarti-Sala. *Rheumatology, Hospital Clinic Barcelona, Barcelona, Spain*

Background: Interstitial lung disease (ILD) is an extra-articular manifestation of Rheumatoid Arthritis (RA), has been related with a poor prognosis. However, there is a lack of clinical data regarding its evolution with different treatment approaches.

Objectives: Our aim was to assess clinical features and evolution of ILD in a cohort of RA patients from a tertiary hospital

Methods: Single-centre retrospective observational study including all patients diagnosed with ILD with a previous or posterior diagnosis of RA evaluated from January 2007 to December 2016 in the Rheumatology Department of a university hospital.

Variables analysed: age, sex, smoking status, duration of RA, presence and titers of ACPA (antiCCP2) and rheumatoid factor (RF), presence of erosions, use of glucocorticoids (GC), synthetic and biological DMARDs before and after ILD diagnosis and presence of other extra-articular manifestations.

We also analysed: duration of ILD, HRCT (High Resolution Computed Tomography) patterns and pulmonary function tests (PFT) at baseline and during follow-up to assess the evolution of the disease. The PFT results were evaluated according to the American Thoracic Society/European Respiratory Society definitions (1).

Results: Twenty-one patients (18 women) were included, mean age 68.4±11.9, mean RA duration 12.3±8.2 years. 9.5% were current smokers and 33.3% former smokers. 61.9% were ACPA+ (52.4% with high basal titers, median ACPA titers: 698) and 81% RF+ (47.6% with high basal titers). 71.4% had erosions and 5 patients (23.8%) had extra-articular manifestations, mainly rheumatoid nodules. RA diagnosis was made after that of ILD in 4 patients; mean elapsed time 1.56±1.26 years. In the other 17 patients, the mean duration of RA until ILD diagnosis was 11.2±7.2 years.

Currently 17 subjects (85%) receive GC; 4 (19%) in monotherapy and 11 (57.9%) synthetic DMARDs (5 methotrexate, 3 leflunomide (LEF), 1 hydroxychloroquine (HCQ) and 2 with double therapy (HCQ/AZA and HCQ/LEF)).

Four patients are currently treated with biological therapy (2 abatacept (ABA), 1 rituximab and 1 etanercept (ETN)), all of them in monotherapy.

Five patients received TNFi prior to ILD diagnosis (3 ETN, 1 infliximab and 1 adalimumab). Biological therapy was withdrawn in 3 cases and switched to ABA in the two remaining ones.

The median duration of EPID was 2.5 (range 0.7–11.9) years. HRCT patterns were: non-specific interstitial pneumonia (NSIP) 52.9%, usual interstitial pneumonia (UIP) 17.6%, cryptogenic organized pneumonia (COP) 11.8%, and other patterns 17.7%.

14 patients had baseline and follow-up PFT: 9 (64.3%) remained stable; 3 (21.4%) improved and 2 (14.3%) worsened (1 had received previous ETN). There was a trend for worsening of PFT in patients with UIP pattern.

No patient died from ILD during this period

Conclusions: In our RA-ILD cohort, NSIP was the most common HRCT pattern and PFT remained stable in most patients during follow-up. Five patients (23.8%) had received TNFi prior to ILD diagnosis and only one of them showed worsening of PFT during ETN treatment, with clinical improvement after withdrawal.

References:

- [1] Pellegrino R et al. Eur Respir J. 2005;26:94868.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2017-eular.3284

SAT0129 EXPLORATION OF COMORBIDITY INDICES IN AN ETHNIC RHEUMATOID ARTHRITIS SUBSET

S. Dowell¹, G. Kerr², C. Swearingen³, M. Quinones¹, J. Berrian¹, S. Hochberg¹ on behalf of EMRAC Working Group. ¹Rheumatology, Howard University Hospital; ²Rheumatology, Washington VAMC, Washington; ³NYU, New York, United States

Background: Comorbidity is common in ethnic patients with rheumatoid arthritis (RA) and often impact choice of DMARD therapy and clinical outcomes. In addition to simple summation of comorbidities, composite indices have been proffered as better predictors of poor clinical outcomes in patients with rheumatoid arthritis¹. However, their applicability in ethnic RA subsets is unknown.

Objectives: To compare composite to mean comorbidity indices in ethnic subsets with rheumatoid arthritis.

Methods: Patients enrolled in the Ethnic Minority RA Consortium (EMRAC), with at least 6 months of followup data were analyzed. At enrollment, sociodemographic data and RA disease status, rheumatic disease comorbidity index (RCDI)¹ and comorbidity count (COUNT)¹ were analyzed amongst ethnic subsets. Spearman correlation was estimated between RCDI and COUNT while Poisson regression

was used to model the differences in the comorbidity indices between race groups.

Results: 453 EMRAC subjects were analyzed; 342 (81.4%) were female, average age was 58.9 ($\pm$ 15.1) years, average duration 13.3 ($\pm$ 11.1) years and average follow-up length 2.1 ($\pm$ 1.4) years. Individual comorbidity frequencies as well as comorbidity indices are summarized in Table 1. Spearman correlation between overall RDCI and COUNT was 0.90 (95% CI [0.88, 0.91], $P < 0.001$). Hispanics, however, had significantly lower comorbidity indices scores than other race groups (RDCI Hispanic vs White $P = 0.003$, Hispanic vs Black $P = 0.004$, Hispanic vs Other $P = 0.038$; COUNT Hispanic vs White $P = 0.012$, Hispanic vs Black $P = 0.004$, Hispanic vs Other $P = 0.151$).

Table 1. Clinical Characteristics of EMRAC study cohort

	White	Black	Hispanic	Other	Total
N	102	195	85	71	453
Hypertension [N (%)]	36 (35.3%)	79 (40.5%)	24 (28.2%)	31 (43.7%)	170 (37.5%)
Coronary Artery Disease [N (%)]	12 (11.8%)	14 (7.2%)	3 (5.9%)	2 (2.8%)	31 (6.8%)
Other Heart Disease [N (%)]	11 (10.8%)	6 (3.1%)	2 (2.4%)	2 (2.8%)	21 (4.6%)
Fracture [N (%)]	11 (10.8%)	8 (4.1%)	2 (2.4%)	2 (2.8%)	23 (5.1%)
Psychiatric Disease [N (%)]	5 (4.9%)	10 (5.1%)	1 (1.2%)	0 (0.0%)	16 (3.5%)
Diabetes [N (%)]	11 (10.8%)	33 (16.9%)	9 (10.6%)	8 (11.3%)	61 (13.5%)
Cancer [N (%)]	9 (8.8%)	8 (4.1%)	5 (5.9%)	3 (4.2%)	25 (5.5%)
Stomach [N (%)]	8 (7.8%)	9 (4.6%)	8 (9.4%)	7 (9.9%)	32 (7.1%)
Stroke [N (%)]	4 (3.9%)	6 (3.1%)	2 (2.4%)	2 (2.8%)	14 (3.1%)
Hyperlipidemia [N (%)]	19 (18.6%)	29 (14.9%)	9 (10.6%)	12 (16.9%)	69 (15.2%)
Chronic Bronchitis [N (%)]	10 (9.8%)	17 (8.7%)	5 (5.9%)	5 (7.0%)	37 (8.2%)
Pulmonary Fibrosis [N (%)]	2 (2.0%)	5 (2.6%)	4 (4.7%)	0 (0.0%)	11 (2.4%)
Kidney Disease [N (%)]	3 (2.9%)	5 (2.6%)	0 (0.0%)	2 (2.8%)	10 (2.2%)
RAPID3	13.1 (6.8)	12.5 (7.4)	14.8 (6.9)	13.7 (7.2)	13.3 (7.1)
Enrollment					
RDCI	1.8 (2.1)	1.7 (2.0)	1.2 (1.7)	1.6 (1.8)	1.6 (1.9)
COUNT	1.0 (1.3)	1.0 (1.3)	0.7 (0.9)	0.9 (1.1)	0.9 (1.2)

Mean (SD) reported unless otherwise noted.

Conclusions: In an ethnic RA subset, the RDCI and COUNT correlated well across ethnic groups. The comorbidity indices were not associated with RA disease activity at enrollment as assessed by the RAPID3. Hispanics had lower RDCI and COUNT scores compared to other race groups but also presented at a younger age which could account for this difference. More studies are needed to determine the predictive value of these indices in determining poor outcomes and mortality in this ethnic subset.

References:

- [1] England, BR et al. Arthritis Care Res. 2015 Jun; 67(6): 885–872.
- Disclosure of Interest:** S. Dowell Grant/research support from: Bristol Myers Squibb, Pfizer, Genentech, Speakers bureau: Horizon Pharma, G. Kerr Grant/research support from: Bristol Myers Squibb, Pfizer, Genentech, C. Swearingen Grant/research support from: Bristol Myers Squibb, Pfizer, Genentech, M. Quinones Grant/research support from: Bristol Myers Squibb, Pfizer, Genentech, J. Berrian Grant/research support from: Bristol Myers Squibb, Pfizer, Genentech, S. Hochberg Grant/research support from: Bristol Myers Squibb, Pfizer, Genentech

DOI: 10.1136/annrheumdis-2017-eular.5842

SAT0130 EVALUATION OF THE FACTORS RELATED WITH RISK OF FALLING AND FEAR OF FALLING IN PATIENTS WITH RHEUMATOID ARTHRITIS

S. Unsal Delialioğlu¹, G. Kilinc Kamaci¹, S. Ozel¹, F. Yurdakul², H. Bodur².

¹Ankara Physical Therapy and Rehabilitation Training and Research Hospital;

²Ankara Numune Training and Research Hospital, Ankara, Turkey

Background: People with rheumatoid arthritis (RA) have an increased risk of falls (1,2). The identification of predictive and potentially modifiable risk factors is essential for the development of effective falls prevention strategies (3).

Objectives: The aim of the study is to determine the relationship between falling risk and falling fear with balance, lower extremity muscle strength, functional status, disease activity and pain in patients with Rheumatoid Arthritis (RA)

Methods: A total of 124 patients (92 females, 32 males) who were diagnosed with RA according to the American College of Rheumatology (ACR) diagnostic criteria were included in the study. The fear of falling was assessed by the Fall Efficacy Scale (FES-I) and the shortened form (Short FES-I). Disease activity was assessed using the Disease Activity Score-28 (DAS28), the functional status was assessed with Health Assessment Questionnaire (HAQ). While lower extremity muscle strength was determined using Chair Stand Test, Four Balance Test was used to determine the balance. The pain intensity was determined using 10 cm Visual Analogue Scale-Pain (VAS-pain).

Results: The mean age of the patients is 54.75 $\pm$ 10.96 years, the mean duration of illness is 14.10 $\pm$ 10.99 years. A statistically significant relationship was found between FES-I, Short FES-I scores and HAQ (respectively $r:0.776$, $p:0.000$; $r:0.783$, $p:0.000$), VAS rest ($r:0.397$, $p:0.000$; $r:0.405$, $p:0.000$), VAS motion ($r:0.542$, $p:0.000$; $r:0.519$, $p:0.000$), DAS28 values ($r:0.216$, $p:0.181$; $r:0.260$, $p:0.004$). There was a significant negative correlation between FES-I, Short FES-I scores and Chair Stand Test (respectively $r:-0.644$, $p:0.000$; $r:-0.652$, $p:0.000$), Four Balance Test ($r:-0.597$, $p:0.000$; $r:-0.611$, $p:0.000$). There was a statistically significant correlation between the number of falls in the last year and age, duration of illness, HAQ, DAS28, VAS scores ($p < 0.05$)

Conclusions: The risk of falling and fear of falling is associated with balance, lower extremity muscle strength, disease activity, functional status, and pain in patients with RA. Therefore; the risk of falling and fear of falling should be determined and necessary measures must be taken in patients with RA.

References:

- [1] Böhler C, Radner H, Ernst M, Binder A, Stamm T, Aletaha D, Smolen JS, Köller M. Rheumatoid arthritis and falls: the influence of disease activity. *Rheumatology* (Oxford). 2012 Nov;51(11):2051–7.
- [2] Armstrong C, Swarbrick CM, Pye SR, O'Neill TW. Occurrence and risk factors for falls in rheumatoid arthritis *Ann Rheum Dis*. 2005 Nov;64(11):1602–4.
- [3] Stanmore EK, Oldham J, Skelton DA, O'Neill T, Pilling M, Campbell AJ, Todd C. Risk factors for falls in adults with rheumatoid arthritis: a prospective study. *Arthritis Care Res* (Hoboken). 2013 Aug;65(8):1251–8.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2017-eular.2022

SAT0131 MEDIAN NERVE THICKNESS RELATED TO RENAL IMPAIRMENT IN RHEUMATOID ARTHRITIS

S. Hannawi¹, I. AlSalmi². ¹Rheumatology, Ministry of Health and Prevention of UAE, Dubai, United Arab Emirates; ²Nephrology, Royal Hospital, Muscat, Oman

Background: Autoimmune processes, contribute to the burden of kidney disease. The reported kidney disease prevalence in patients with rheumatoid arthritis (RA) ranges from 5%–50% based on studies of different designs. Historically, patients with RA-associated renal amyloidosis had higher mortality rates relative to the general population. The occurrence of haematuria, proteinuria, or CKD has been reported to be associated with a 3- to 4-fold increased risk of death. Subclinical decreased kidney function has been identified as an independent risk factor for cardiovascular (CV) events with increase mortality in patients with RA. On the other hand, both rheumatoid arthritis and renal impairment have been reported to be associated with increase prevalence of carpal tunnel syndrome.

Objectives: To establish the median nerve thickness (measured by ultrasound) in RA and its relation to renal function.

Methods: 120 RA patients were recruited through a specialized rheumatology clinic. The US measurements were performed by the same person. Patients were sitting with their forearm resting in a supinated position on a small table. The US probe (an 8–16 MHz linear array transducer) was held as lightly as possible to avoid disturbing the anatomy of the nerve. The median nerve was examined at the entrance of the carpal tunnel, between the pisiform bone and the tubercle of the scaphoid bone, where the distal volar crease is an external pisiform landmark. A continuous trace was made just within the hyperechogenic boundary of the nerve. The cross-sectional area of the median nerve was calculated directly by the software of the US equipment. Each median nerve was measured three times, and the mean value was used for further analyses. Modification of Diet in Renal Disease (MDRD) equation used to estimate the Glomerular Filtration Rate (GFR). The average of the right and the left areas of the median nerve were used when exploring bivariate correlations to the renal variables (Pearson's correlation coefficients). All statistics were performed using STATA programme.

Results: the average median nerve thickness was 9.79 $\pm$ 2.6 mm² (Range 1.5–22.25). The average GFR was 122 $\pm$ 20 ml/min (59.6–286). Thickness of the median nerve was positively associated with the age of the participants ($p=0.03$, CI: 0.00, 0.08), body mass index ($p=0.04$, CI: 0.00, 0.21), uric acid level ($p=0.033$, CI: 0.00, 0.01), and urine microalbumin ($p=0.04$, CI 0.00, 0.01). GFR showed no significant relation the thickness of the median nerve.

Conclusions: RA patients without symptoms or clinical signs have a median nerve thickness that is positively correlated to the level of microalbumin and uric acid. Whether sonographic examination of the median nerve would be helpful in predicting who is going to have a deteriorated renal function need to be explored in a larger study.

References:

- [1] Boers, M., et al., Subclinical renal dysfunction in rheumatoid arthritis. *Arthritis Rheum*, 1990 33(1): p. 95–101.
- [2] Daoussis, D., et al., Cardiovascular risk factors and not disease activity, severity or therapy associate with renal dysfunction in patients with rheumatoid arthritis. *Ann Rheum Dis*, 2010. 69(3): p. 517–21.
- [3] Pathan, E. and V.R. Joshi, Rheumatoid arthritis and the kidney. *J Assoc Physicians India*, 2004 52: p. 488–94.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2017-eular.1660

SAT0132 THE RHEUMATOID ARTHRITIS FLARE QUESTIONNAIRE (RA-FQ): RESULTS OF RASCH ANALYSIS AND FEEDBACK ON REAL-WORLD APPLICATIONS FROM INTERNATIONAL RA PATIENTS AND CLINICIANS

S.J. Bartlett^{1,2}, S.P. Barbic³, V.P. Bykerk⁴, B. Fautrel⁵, F. Guillemin⁶, A. den Broeder⁷, R. Alten⁸, R. Christensen⁹, E. Choy¹⁰, D. Furst¹¹, S. Hewlett¹², A. Leong¹³, L. March¹⁴, T. Woodworth¹¹, B.I.O. Clifton² on behalf of CATCH, STPR, & DRESS Investigators. ¹McGill University, Montreal, Canada; ²Johns Hopkins, Baltimore, United States; ³University of British Columbia, Vancouver, Canada; ⁴HSS, New York, United States; ⁵Pierre & Marie Curie University, Paris;