DOI: 10.1136/annrheumdis-2017-eular.6490

SAT0209 OBSERVATIONAL STUDY ON THE EFFECTS OF IL-6 INHIBITOR THERAPY ON MYOSTATIN IN PATIENTS WITH RHEUMATOID ARTHRITIS

M.J. Chapman¹, R.P. Narayanan², A. Cross³, R. Moots³, J. Wilding², N. Goodson³. ¹Department of Rheumatology, Aintree University Hospital; ²Department of Obesity and Endocrinology; ³Institute of Ageing and Chronic Disease, University of Liverpool, Liverpool, United Kingdom

Background: Rheumatoid cachexia (RC), a phenotype of increased adiposity, reduced lean mass and insulin resistance can occur in rheumatoid arthritis (RA). It is most prevalent in those with high disease activity and is associated with increased cardiovascular morbidity and mortality. Tocilizumab (TCZ), a monoclonal anti-IL-6 antibody, successfully reduces disease activity in RA. It has also been associated with improved insulin sensitivity, increased lean mass and hyperlipidaemia. The serum concentration of myostatin, a myokine released by skeletal muscle, is increased in most disease states associated with cachexia. Moreover, its knockout and inhibition increase lean mass. However, its role in RA has not been fully elucidated.

Objectives: To investigate the effect of tocilizumab treatment on serum myostatin in patients with RA.

Methods: 19 patients with RA (16 female, 3 male) mean age 49.6yrs, mean DAS28 6.1, median disease duration 10yrs (range 0 to 30), mean BMI 27.4, received 13 IV infusions of TCZ 8mg/kg every four weeks as part of a 52 week, single-centre, open-label study (ACT-NEUT [1]). 8 had previously been exposed to a biological agent, 4 were on regular steroids, and all patients received concurrent methotrexate. Serum myostatin was measured at 0, 1, 3 and 6 months of treatment using ELISA. BMI, serum lipid profile, CRP, ESR and DAS28 were measured at each visit (0, 1, 3 and 6 months). Data were analysed using STATA 14. Change between 0, 3 and 6 months was analysed using Wilcoxon signed rank test, statistical significance was confirmed using mixed model analysis adjusting for BMI, age and gender. Linear regression with adjustment for age, BMI and gender assessed correlation at baseline.

Results: DAS28, CRP and ESR all significantly decreased with TCZ treatment. A significant increase in BMI (25.5 (IQR 21.6–31.7) vs 26.0 (IQR 21.4–32.5) $p=0.0052$) between baseline and 6 months and serum triglycerides between baseline and 3 months (1.3 (IQR 1.3–2.2) vs 1.6 (IQR 1.3–2.6) $p=0.028$) was seen with TCZ. Baseline serum myostatin concentrations were negatively correlated with baseline DAS28 ($r^2=0.39$ $p=0.038$). Treatment with tocilizumab for 6 months resulted in a significant increase in serum myostatin from baseline (2.7ng/mL (IQR 2.0–3.1) vs 3.3ng/mL (IQR 3.0–5.0) $p<0.001$) [Fig. 1]. Moreover, a significant association was seen between baseline myostatin and change in cholesterol at 3 months ($r^2=0.51$ $p=0.029$), when adjusting for baseline cholesterol, age, gender, BMI and change in BMI.

Fig 1. Serum myostatin levels significantly increase after six months of tocilizumab in patients with rheumatoid arthritis. Values shown are actual serum values for individual patients ($n=19$ at baseline $n=15$ at 6 months). Black lines represent increasing serum values, grey lines decreasing values with treatment. Circles represent median±IQR. Change evaluated by Wilcoxon signed rank test $p<0.001$

Conclusions: We have demonstrated a significant correlation between myostatin and DAS28 and a significant change in myostatin with tocilizumab treatment. It is possible that IL-6 blockade results in a rise in myostatin. This might attenuate any improvement in muscle wasting. Future studies should measure body composition and muscle function to help understand the changes we have observed.

References:

[1] Wright HL, Cross AL, Edwards SW, Moots RJ. Effects of IL-6 and IL-6 blockade on neutrophil function in vitro and in vivo. *Rheumatology*. 2014 Jul;53(7):1321–31.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2017-eular.5831

SAT0210 IMPROVEMENT OF DISEASE ACTIVITY IN A 5-YEAR COHORT OF RHEUMATOID ARTHRITIS PATIENTS TREATED UNDER TREAT TO TARGET RECOMMENDATIONS RECEIVING BIOLOGICAL THERAPY

P. Santos-Moreno¹, D. Gomez¹, E. Castillo¹, R. Giraldo¹, G. Ballesteros¹, L. Villarreal², J. Bello³, D. Buitrago-Garcia³. ¹Rheumatology; ²Psychology and processes; ³Epidemiology, Biomab, Center for Rheumatoid Arthritis, Bogota, Bogota, Colombia

Background: Treat to Target (T2T) strategy becomes from the need to develop therapeutic targets and tools to achieve defined outcomes in rheumatoid arthritis (RA), this strategy has become recognized as a standard of good practice embodying the principle that rapid attainment of remission, or low disease activity, can halt joint damage and maintain good quality of life. However, there is no direct comparison between biologics in cohorts of patients with long-standing RA using T2T approach in real-life settings, which could have implications in treatment decisions and health economics.

Objectives: The aim of this study was to describe global change in Disease Activity Score 28 (DAS28) using T2T strategy for a 5 year period in patients with biological therapy in a large cohort of patients from a Colombian specialized in RA center with multidisciplinary approach.

Methods: A descriptive dynamic cohort study was performed. Records of patients using biological DMARD treatment (Anti-TNF and others) from specialized in RA center were reviewed; those patients were followed-up under T2T standards. Clinical follow-up was according to DAS28 as follows: every 3–5 weeks (DAS28 >5.1), every 7–9 weeks (DAS28 ≥3.1 and ≤5.1), and every 11–13 weeks (DAS28 <3.1). Therapy had to be adjusted with DAS28 >3.2 unless patient's conditions don't permit it. We divided patients in three groups: low disease activity (LDA), moderate disease activity (MDA) and severe disease activity (SDA) patients. Descriptive epidemiology was done, percentages and averages were calculated; the median of each variable was analyzed using t-Student assuming normality for DAS28 distribution and the level activity disease was analyzed using Pearson's statistics.

Results: During 60 month period we included 695 patients with biological therapy, 85% were female and 15% male. Mean age was 58 years ± 11. At beginning mean DAS28 was 4.1±1.1 and 20% of patients were in SDA, and 55% in MDA. At the end of 5 year period mean DAS was 3.2±0.89 and only 7% of patients were in SDA and 33% in MDA. The median was analyzed using t-Student assuming normality for DAS28 distribution. It showed statistical significant improvement ($p<0.00$).

ACTIVITY LEVEL	TIME 0 n(%)	2011-2012 n(%)	2013-2014(%)	2015-2016(%)
REM	-	50 (7)	166 (24)	217 (31)
LDA	178(26)	236 (34)	181(26)	198(28)
MDA	382 (55)	361(52)	278 (40)	234(74)
SDA	135 (19)	98(14)	70(10)	46(7)

Conclusions: There was a global improvement of DAS28 in a cohort of RA patients receiving biological therapy followed and treated under recommendations of T2T approach demonstrating that with the strategies mentioned above it is possible to obtain a good control of the activity of the disease.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2017-eular.5579

SAT0211 REAL WORLD DATA OF RITUXIMAB EFFECTIVENESS IN RHEUMATOID ARTHRITIS: DIFFERENCES BETWEEN BIOLOGIC-NAIVE PATIENTS AND PREVIOUSLY EXPOSED TO BIOLOGICS

R.M. Ferreira, R. Fonseca, D.R. Gonçalves, F. Aguiar, T.M. Rocha, A. Bernardo, S. Pimenta, M. Bernardes, L. Costa. São João Hospital Centre, Porto, Portugal

Background: Rituximab is only approved for rheumatoid arthritis (RA) treatment in patients with an incomplete response or intolerance to others DMARDs, including TNF alpha inhibitors. It represents a significant advance in RA biologics arsenal due to its safety and efficacy profiles.

Objectives: To evaluate the effectiveness of rituximab in RA patients and to compare the response between first-line rituximab patients and those previously exposed to other biologics.

Methods: An observational retrospective study was conducted, including all the consecutive patients with diagnosis of RA under rituximab, followed at our Rheumatology department until December 2016. Demographic and clinical data were obtained by consulting the national database (Reuma.pt). DAS28 variations and EULAR response were measured at 6, 12 and 18 months. Parametric and non parametric tests were used for statistics (SPSS 20.0).

Results: We included 63 RA patients (81% of women), with a mean (SD) age of 61 (10) years and a mean disease duration of 19 (10) years; 86% rheumatoid factor positive and 87% anti-citrullinated peptide antibody (ACPA) positive. Bone erosions and extra-articular manifestations were present in 85.7% and 58.7% of the patients, respectively. At baseline, the mean DAS28 was 5.79 (65% and 29% of patients with severe and moderate disease activity, respectively, and 6% in clinical remission). Thirty patients were treated with rituximab as first-line therapy and 33 patients were previously exposed to other biologics. Combination therapy

with methotrexate (MTX) was observed in 48% and with others classic DMARDs in 30%, while 22.3% received rituximab monotherapy. First-line rituximab option was justified by lung involvement in 21%, past malignancy in 13%, recurrent infections in 5%, congestive cardiac failure in 3%, vascular involvement in 3% and untreated latent tuberculosis in 3%. In the group previously exposed to biologics, 13% switched therapy due to ineffectiveness and 87% due to adverse events. No significant differences were found between the 2 groups in terms of age, gender, concomitant use of MTX and baseline DAS28. The group previously exposed to biologics had a longer disease duration (mean 23 vs 15 years, $p=0.001$) and fewer patients with ACPA seropositivity (79% vs 97%, $p=0.035$). There was a significant reduction of DAS28 at 6, 12 and 18 months ($p<0.001$ for all). Fifty six percent of the patients achieved a EULAR response at 6 months, 46% at 12 months and 59% at 18 months. DAS28 variation at 6 months differed significantly between groups, with a better clinical response in naive biological patients comparing to those previously exposed to biologics (median 1.173 vs 0.477; $p=0.038$). There were no differences in terms of DAS28 variation at 12 and 18 months ($p=0.642$ and $p=0.135$, respectively) and in EULAR responses at 6, 12 and 18 months between the groups ($p=0.289$, $p=0.523$ and $p=1.000$, respectively).

Conclusions: Our study confirms the effectiveness of rituximab in RA patients and suggests a higher magnitude of response in naive biological patients at 6 months of RTX therapy. These findings put in perspective an extension of rituximab as a first-line biologic for RA treatment.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2017-eular.4494

SAT0212 SARILUMAB DOSE REDUCTION IN AN OPEN-LABEL EXTENSION STUDY IN RA PATIENTS WITH INADEQUATE RESPONSE TO TNF INHIBITORS

R. Fleischmann¹, P. Hrycaj², H. van Hoogstraten³, E.K. Mangan⁴, Y. Lin³, S. Jayawardena³, G.R. Burmester⁵. ¹Metrex Clinical Research Center, University of Texas Southwestern Medical Center, Dallas, United States; ²Department of Rheumatology and Clinical Immunology, Poznań University of Medical Sciences, Poznań, Poland; ³Sanofi Genzyme, Bridgewater; ⁴Regeneron Pharmaceuticals, Inc, Tarrytown, United States; ⁵Charité - University Medicine Berlin, Free University and Humboldt University Berlin, Berlin, Germany

Background: Sarilumab is a human mAb blocking the IL-6R α . In the phase 3 TARGET study (NCT01709578), sarilumab (150 or 200 mg SC q2w) + csDMARDs demonstrated efficacy in adults with active, moderate-to-severe RA and inadequate response or intolerance to ≥ 1 TNFi. Infections, neutropenia, injection site reactions, increased lipids, and increased transaminases were among the most common TEAEs. EXTEND (NCT01146652) is an open-label extension study evaluating long-term safety and efficacy of sarilumab in patients from the sarilumab clinical development program.

Objectives: To examine the effects of dose reduction of sarilumab 200 mg q2w to 150 mg q2w in patients from TARGET that occurred in EXTEND primarily for protocol-specified laboratory abnormalities.

Methods: Patients were switched to or initiated on sarilumab 200 mg q2w after enrolling in EXTEND. Per protocol, investigators could reduce the sarilumab dose from 200 mg q2w to 150 mg q2w for ANC ≥ 0.5 to 1.0 Giga/L, platelet count ≥ 50 to 100 Giga/L, or ALT ≥ 3 to 5 $\times$ ULN. Dose reductions were also performed at the investigator's discretion. Efficacy data from EXTEND were analyzed before and 24 weeks after dose reduction.

Results: As of the July 2016 interim analysis (N=452), dose reduction from sarilumab 200 mg q2w to 150 mg q2w had occurred in 14.6% of patients (n=66) from TARGET. The most common reasons for dose reduction were decreased ANC (8.8%; n=40) and increased ALT (3.3%; n=15). At the time of analysis, 80.3% of patients (n=53) whose dose was reduced were continuing treatment, with a median treatment duration of 1.6 years after dose reduction. Improvements in ANC and ALT were observed over the 6 months after dose reduction (Table 1). Efficacy was maintained 24 weeks after dose reduction (Table 2).

Table 1. ANC and ALT After Dose Reduction^a

	Before dose reduction n (%)	6 months after dose reduction n (%)
ANC		
≥ 0.5 to <1.0 Giga/L	25/40 (62.5)	5/39 (12.8)
<0.5 Giga/L	0	0
ALT		
>3 to $\leq 5 \times$ ULN	13/15 (86.7)	1/13 (7.7)
>5 to $\leq 10 \times$ ULN	1/15 (6.7)	0

^aDenominator represents patients who dose reduced because of decreased ANC or increased ALT who had ANC/ALT measured within specified time point.

Table 2. Efficacy After Dose Reduction

	Before dose reduction (N=60)	6 months after dose reduction (N=53)
ACR20 response rate, n (%)	49 (81.7)	43 (81.1)
Δ HAQ-DI, mean (SD)	-0.7 (0.6)	-0.8 (0.7)
Δ DAS28-CRP, mean (SD)	-3.0 (1.2)	-3.0 (1.3)
Δ CDAI, mean (SD)	-28.4 (12.5)	-28.6 (14.6)

Conclusions: In patients from TARGET whose sarilumab dose was reduced

from 200 mg q2w to 150 mg q2w during EXTEND, there was an improvement in laboratory abnormalities and continuation of treatment for the majority of patients. Improvements in signs and symptoms of RA and physical function were maintained after dose reduction.

Acknowledgements: This study was sponsored by Sanofi Genzyme and Regeneron Pharmaceuticals, Inc. Editorial support was provided by MedThink SciCom and funded by Sanofi Genzyme and Regeneron Pharmaceuticals, Inc.

Disclosure of Interest: R. Fleischmann Grant/research support from: AbbVie, Amgen, Ardea Biosciences, Bristol-Myers Squibb, Celgene, GlaxoSmithKline, Eli Lilly, Pfizer, Roche, Sanofi, and UCB, Consultant for: AbbVie, Akros Pharma, Amgen, AstraZeneca, Bristol-Myers Squibb, Janssen, Eli Lilly, Pfizer, Roche, and UCB, P. Hrycaj Consultant for: Pfizer, Celltrion, and Eisai, H. van Hoogstraten Shareholder of: Sanofi Genzyme, Employee of: Sanofi Genzyme, E. Mangan Shareholder of: Regeneron Pharmaceuticals, Inc, Employee of: Regeneron Pharmaceuticals, Inc, Y. Lin Shareholder of: Sanofi Genzyme, Employee of: Sanofi Genzyme, S. Jayawardena Shareholder of: Sanofi Genzyme, Employee of: Sanofi Genzyme, G. Burmester Grant/research support from: AbbVie, Bristol-Myers Squibb, MedImmune, Merck, Pfizer, Roche, and UCB, Consultant for: AbbVie, Bristol-Myers Squibb, MedImmune, Merck, Pfizer, Roche, and UCB, Speakers bureau: AbbVie, Bristol-Myers Squibb, Merck, Pfizer, Roche, and UCB
DOI: 10.1136/annrheumdis-2017-eular.5155

SAT0213 LOW DOSE IL-2 RESTORES DECREASED ABSOLUTE NUMBER OF REGULATORY T CELLS AND IMBALANCE BETWEEN TH17 AND REGULATORY T CELLS IN PATIENTS WITH RHEUMATOID ARTHRITIS

R. Jia¹, X. Li¹, C. Wang¹, C. Gao². ¹The Department of Rheumatology, the Second Hospital of Shanxi Medical University, Taiyuan, China; ²Pathology, Joint Program in Transfusion Medicine, Brigham and Women's Hospital/Children's Hospital Boston, Harvard Medical School, Boston, United States

Background: A therapeutic revolution in the past decade is still a considerable unmet need in the treatment of rheumatoid arthritis (RA). On the other hand, dysfunction of regulatory T cells (Tregs) has been considered to be a pivotal cause of RA and correction of this dysfunction to be a potential RA therapy¹. However, abnormalities of Tregs in patients with RA were reported controversially in previous studies², in which only proportion was measured and Tregs were defined using different protein markers.

Objectives: In this study, we measured both absolute numbers and proportions of CD4+CD25+Foxp3+ Tregs in peripheral blood of RA patients and investigated the effects of low-dose recombinant human IL-2 (rhIL-2) on Tregs and CD4+ effector T cell subsets in patients with RA.

Methods: Both absolute numbers and proportions of Treg and Th17 cells in peripheral blood, defined as the CD4+CD25+FOXP3+T or CD4+IL-17+ T cell population, were examined by flow cytometry in 342 patients with RA with different 28-joint Disease Activities (DAS28), including 75 who had never received disease-modifying antirheumatic drugs (DMARD) and 151 who were receiving or had received DMARD. Among these patients, 112 consented at enrollment to receive rhIL-2 treatment. Before and after treatment, the Th17 and Treg cells in peripheral blood were analyzed by flow cytometry.

Results: The absolute count of Treg cells in patients with RA was significantly low compared with that of healthy controls ($P<0.05$), but the absolute count of Th17 cell was no different between RA and healthy controls. After the course of rhIL-2 treatment, there was a significant increase in the absolute count of Th17 and Treg cells in the CD4+ T cell population ($P<0.01$), but Th17/Treg was significantly low after the treatment.

Conclusions: The data suggest that, besides proportion, the decrease of Treg cell number, defined as the CD4+CD25+FOXP3+population, may contribute to the pathogenesis of RA. Over the treatment of rhIL-2, there was a more significant increase in the absolute count of Treg cells than that of Th17, and consequently restore the balance of Th17/Treg.

References:

- [1] Miyara M, Ito Y, Sakaguchi S. TREG-cell therapies for autoimmune rheumatic diseases. *Nature reviews Rheumatology* 2014;10:543–51.
- [2] Morita T, Shima Y, Wing JB, Sakaguchi S, Ogata A, Kumanogoh A. The Proportion of Regulatory T Cells in Patients with Rheumatoid Arthritis: A Meta-Analysis. *PLoS one* 2016;11:e0162306.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2017-eular.2808

SAT0214 AN ANALYSIS OF INCREASING SPACING TIME FOR THE INTRAVENOUS ADMINISTRATION OF TOCILIZUMAB IN PATIENTS WITH RHEUMATOID ARTHRITIS

S. Ito¹, D. Kobayashi¹, C. Takai¹, Y. Nomura¹, A. Abe¹, H. Otani¹, H. Ishikawa¹, A. Murasawa¹, I. Narita², K. Nakazono¹. ¹Niigata Rheumatic Center, Shibata; ²Division of Clinical Nephrology and Rheumatology Niigata University Graduate School of Medical and Dental Sciences, Niigata, Japan

Background: Although the biological disease-modifying antirheumatic drugs (bDMARDs) are the most effective treatment for rheumatoid arthritis (RA), the costs are much higher than those of conventional synthetic DMARDs. A number