[3] Matucci-Cerinic M, Kahaleh B, Wigley FM. Review: evidence that systemic sclerosis is a vascular disease. *Arthritis and rheumatism* 2013; 65(8):1953–62.

[4] Pattanaik D, Brown M, Postlethwaite BC, Postlethwaite AE. Pathogenesis of Systemic Sclerosis. *Frontiers in immunology* 2015; 6:272.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2017-eular.3739

THURSDAY, 15 JUNE 2017

Cytokines and chemokines

OP0097

SYSTEMIC IFN TYPE I AND TYPE II SIGNATURES IN PRIMARY SJÖGREN'S SYNDROME REVEAL DIFFERENCES IN DISEASE SEVERITY

I. Bodewes¹, S. Al-Ali^{2,3}, C.G. van Helden¹, N.I. Maria¹, J. Tarn², D. Lendrem², M.W. Schreurs¹, E.C. Steenwijk¹, P.L.A. van Daele^{1,4}, T. Both⁴, S. Bowman⁵, B. Griffiths⁶, W.-F. Ng^{2,7}, M.A. Versnel¹. ¹*Immunology, Erasmus MC, Rotterdam, Netherlands*; ²*Musculoskeletal Research group, Newcastle University, Newcastle upon Tyne, United Kingdom*; ³*University of Basrah, Basrah, Iraq*; ⁴*Internal Medicine, Erasmus MC, Rotterdam, Netherlands*; ⁵*Rheumatology, University Hospital Birmingham, Birmingham*; ⁶*Newcastle upon Tyne Hospitals NHS Foundation Trust*; ⁷*National Institute for Health Research, Newcastle Biomedical Research Centre, Newcastle upon Tyne, United Kingdom*

Background: Local and systemic activation of interferons (IFNs) has been demonstrated in primary Sjögren's syndrome (pSS).[1–4] Type I IFNs are associated with higher disease activity and autoantibody levels.[5] Recent findings also show activation of interferon type II (IFN_γ) induced gene expression in salivary glands of pSS patients.[6, 7] Although IFN type I and II bind to different receptors they induce partially overlapping gene expression patterns. Understanding the relative contribution of IFN type I and type II may deepen our knowledge in pSS pathogenesis and promote a stratified approach to therapeutic development.

Objectives: Determine IFN type I and II inducible gene expression in patients with pSS and correlate this to disease manifestations.

Methods: In whole blood of 50 pSS patients modular IFN scores were determined using real-time quantitative PCR followed by principal component analysis. Subsequently, five indicator genes per module were analysed in two independent European cohorts with a total of 141 patients.

Results: Three groups were distinguished: without IFN activation (19–47%), with IFN type I (53–81%) and with IFN type I+II activation (35–55%). Patients with IFN activation (I or I+II) have a higher presence of auto-antibodies, IgG levels and lower lymphocyte counts compared to IFN negative patients. The biological domain of the EULAR Sjögren's Syndrome Disease Activity Index (biological-ESSDAI) was higher in patients with IFN activation, while total-ESSDAI scores were not significantly different. 66–67% of the IFN type I positive patients had an additional IFN type II inducible gene expression. Patients with IFN type I+II activation have significantly higher IgG levels and lower lymphocyte counts compared to patients with only IFN type I activation. There were no differences in fatigue or dryness, but pain scores were lower.

Conclusions: pSS patients can be stratified according to their systemic IFN activation patterns. IFN activation (I or I+II) is present in patients with the highest activity of the biological-ESSDAI. These data raise the possibility that the biological-ESSDAI rather than total-ESSDAI score may be a more sensitive endpoint for trials targeting either type I or type II IFN pathways.

References:

[1] Wildenberg, et al. *EJL*. 2008; 38(7):2024–2033.

[2] Gottenberg, et al. *PNAS*. 2006; 103(8):2770–2775.

[3] Hjelmervik, et al. *A&R*. 2005; 52(5):1534–1544.

[4] Emamian, et al. *Genes Immun*. 2009; 10(4):285–296.

[5] Brkic, et al. *ARD*. 2013; 72(5):728–735.

[6] Hall, et al. *PNAS*. 2013; 72(5):728–735.

[7] Hall, et al. *A&R*. 2012; 109(43):17609–17614.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2017-eular.3688

THURSDAY, 15 JUNE 2017

Progress in biological treatment of RA

OP0098

REMISSION AND MAINTENANCE OF EFFICACY IN A PHASE 2B STUDY OF VOBARILIZUMAB, AN ANTI-INTERLEUKIN 6 RECEPTOR NANOBODY, IN PATIENTS WITH MODERATE-TO-SEVERE RHEUMATOID ARTHRITIS DESPITE TREATMENT WITH METHOTREXATE

T. Dörner¹, M. Weinblatt², P. Durez³, R. Alten⁴, K. Van Beneden⁵, E.J. Dombrecht⁵, K. De Beuf⁵, P. Schoen⁵, R.K. Zeldin⁵. ¹*Rheumatology and Clinical Immunology, Charité University Hospitals, Berlin, Germany*; ²*Division of Rheumatology, Immunology and Allergy, Brigham and Women's Hospital,*

Boston, United States; ³*Department Rheumatology, Cliniques Universitaires St-Luc, Sint-Lambrechts-Woluwe, Belgium*; ⁴*Schlosspark-Klinik, Berlin, Germany*; ⁵*Abylnx NV, Zwijnaarde, Belgium*

Background: Vobarilizumab is a Nanobody® consisting of an anti-IL-6R domain and an anti-human serum albumin domain in development for treatment of RA. The efficacy and safety were assessed in a 24-week double-blind global phase 2b study in patients with active RA on a stable background of MTX. Main efficacy and safety results were previously reported [1].

Objectives: To report the impact of treatment with vobarilizumab on secondary efficacy endpoints including SDAI and CDAI remission and the sustained response at 4 consecutive visits based on ACR50, ACR70 and DAS28_{CRP}.

Methods: Patients were randomized to receive subcutaneously administered placebo or 1 of 4 dose regimens of vobarilizumab in addition to MTX. SDAI and CDAI remission at Week 24 was evaluated, as was maintenance of efficacy as defined by sustained DAS28_{CRP} <2.6 responses at 4 consecutive visits (i.e., at Weeks 12, 16, 20 and 24). In addition, a post-hoc analysis was performed on sustained ACR50 and ACR70 responses from Week 12 through Week 24. Proportions of patients achieving response for these endpoints were summarized by treatment group. Subjects with missing values were analyzed as non-responders.

Results: A total of 345 patients were randomized. Demographics and baseline characteristics were similar across groups with mean baseline DAS28_{CRP} between 5.8 and 6.2. At Week 24, up to 19% and 20% in the vobarilizumab groups reached CDAI and SDAI remission, respectively vs. 10% and 9% who received placebo (Table 1).

Table 1. Percentage of patients with CDAI and SDAI remission at Week 24

	Placebo (N=69)	Vobarilizumab 75mg q4w (N=69)	Vobarilizumab 150mg q4w (N=70)	Vobarilizumab 150mg q2w (N=68)	Vobarilizumab 225mg q2w (N=69)
CDAI remission (≤2.8)	10	15	19	12	19
SDAI remission (≤3.3)	9	10	19	15	20

At Week 24, up to 61% and 45% of the patients in the vobarilizumab groups achieved an ACR50 or ACR70 response, respectively (39% and 17% on placebo). Approximately one third of the randomized patients in the 3 highest treatment groups had a sustained ACR50 response from Week 12 through Week 24 (Table 2). Sustained remission defined by DAS28_{CRP} <2.6 at 4 consecutive visits, i.e. at weeks 12, 16, 20 and 24, was observed in 20% to 25% of the patients in the 3 highest dosing arms compared with 3% of those receiving placebo.

Table 2. Percentage of patients with sustained efficacy response at 4 consecutive visits (Weeks 12, 16, 20 and 24)

	Placebo (N=69)	Vobarilizumab 75mg q4w (N=69)	Vobarilizumab 150mg q4w (N=70)	Vobarilizumab 150mg q2w (N=68)	Vobarilizumab 225mg q2w (N=69)
DAS28 _{CRP} <2.6	3	4	20	25	20
ACR50	16	14	29	31	39
ACR70	4	7	11	13	13

Conclusions: In patients with active RA, treatment with vobarilizumab at the 3 highest dose regimens in addition to MTX had a positive and sustained impact on disease activity through Week 24 as defined by clinically relevant efficacy endpoints.

References:

[1] Weinblatt et al. (Annual Scientific Meeting, Canadian Rheumatology Association, 2017).

Disclosure of Interest: T. Dörner Consultant for: Abylnx, M. Weinblatt Consultant for: Abylnx, P. Durez Consultant for: Abylnx, R. Alten Consultant for: Abylnx, K. Van Beneden Employee of: Abylnx, E. Dombrecht Employee of: Abylnx, K. De Beuf Employee of: Abylnx, P. Schoen Employee of: Abylnx, R. Zeldin Employee of: Abylnx

DOI: 10.1136/annrheumdis-2017-eular.3732

OP0099

SAFETY, TOLERABILITY AND INITIAL SIGNS OF EFFICACY OF THE FULLY HUMAN IMMUNOCYTOKINE DEKAVIL (F8IL10): A NOVEL THERAPEUTIC APPROACH FOR RHEUMATOID ARTHRITIS

M. Galeazzi¹, G. Sebastiani², R. Voll³, J. Wollenhaupt⁴, O. Viapiana⁵, J. Dudler⁶, E. Selvi¹, C. Baldi¹, M. Bardelli¹, B. Bannert³, S. Finzel³, C. Specker⁷, P. Sarzi Puttini⁸, F. Bootz⁹, D. Neri¹⁰ on behalf of the DEKAVIL Study Group. ¹*University Hospital, Siena*; ²*San Camillo-Forlanini Hospital, Rome, Italy*; ³*University Medical Center, Freiburg*; ⁴*Schoen-Klinik, Hamburg Eilbek, Germany*; ⁵*University Hospital, Verona, Italy*; ⁶*Cantonal Hospital, Fribourg, Switzerland*; ⁷*University Hospital, Essen, Germany*; ⁸*Luigi Sacco Hospital, Milan, Italy*; ⁹*Philochem AG, Otelfingen, Switzerland*; ¹⁰*PhiloGen SpA, Siena, Italy*

Background: The antibody-based targeted pharmacodelivery of cytokines by means of immunocytokines has the potential to enhance therapeutic activity at the site of disease while sparing healthy tissues. Dekavil (F8IL10) is a fully human immunocytokine composed of the vascular targeting antibody F8 (specific to