elevated HDAC9 expression and to a decreased Foxp3 acetylation. Lower acetylated Foxp3 determines an instable phenotype of GARP-deficient Tregs and their failure to properly regulate an immune reaction.

Acknowledgements This work supported by DFG grants SK59/09-1 and Schu1683/10-1, and by BMBF Projekt Arthromark 01Ec1401B.

Disclosure of interest None declared

P089

CHARACTERISATION OF CHEMOKINE RECEPTORS AND MIGRATION OF REGULATORY B CELLS IN PATIENTS WITH RHEUMATOID ARTHRITIS AND IN HEALTHY DONORS

^{1,2,3}R Audo*, ^{1,2,3}K Schreiber, ^{1,2,3}B Combe, ^{1,2,3}J Morel, ^{1,2,3}C Daien. ¹IGMM UMR5535; ²Lapeyronie Teaching Hospital; ³Montpellier University, Montpellier, France

10.1136/annrheumdis-2018-EWRR2018.105

Introduction B cells are pathogenic players in the development of rheumatoid arthritis (RA). More recently, it has been shown that B cells can also have regulatory functions, mainly through the secretion of interleukin-10 (IL-10). We found a decrease of IL10 + B (B10) cells in patients with early Rheumatoid Arthritis (RA) and B10 were inversely correlated with disease activity (DAS28), inflammation and rheumatoid factor, suggesting a key role at the initial phase of RA. Chemokines and chemokine receptors (CR) orchestrate migration of immune cells in physiological and pathological conditions. We hypothesise that CR could be involved in B10 functions and characterisation.

Objectives We aim to assess expression of CR on B10 and to understand their migration into the joint.

Methods Peripheral blood mononuclear cells (PBMCs) from 10 RA patients and 10 controls (HC) were activated 24 hours with CpG to generate B10 cells. We compared expression of several CR between B10 cells and IL-10^{neg} B cells by FACS analysis. For functional test, B10 cells and IL-10^{neg} B were sorted by using secretion assay (Miltenyi) and the ability to migrate in response to CCL21, CCL22, CXCL11, CXCL12 or CXCL13 was evaluated by using migration assay in 5 µM Transwell chambers. The presence of B10 cells in RA synovial tissue was evaluated by immunohistochemistry.

Results We found that B10 cells from HC differentially expressed several CR than IL-10^{neg} B cells. There were a strong decrease of CXCR5 and CXCR7 expression, and a strong increase of CCR4 expression in B10 cells ($p < 0.001$) compared to IL-10^{neg} B cells. Functional impact of differential expression was tested by migration assay. Among chemokines tested, CXCL13, CXCR5-ligand, attracted more B10 cells than IL-10^{neg} B cells ($n = 7$). More importantly, our results suggest that CR receptor profile on B10 cells is different between HC and RA ($n = 10$). Preliminary results showed that only a few B10 can be found in RA synovial tissue among B cells. One can hypothesised that the different expression profile of CR on B10 cells of RA patients might explain a defect in joint migration, and thus promoting uncontrolled inflammation.

Conclusions B10 cells have a specific CR profil compared to IL-10^{neg} B cells. The differential expression of CR on RA B10 compared to HC might explain a defect in joint migration. Understanding of B10 migration is an important issue and could be used in the future to drive regulatory B cells into the joints as a new therapeutic approach for RA.

Disclosure of interest None declared

P090

DIFFERENCE BETWEEN PALINDROMIC RHEUMATISM AND RHEUMATOID ARTHRITIS AT THE LEVEL OF GENE EXPRESSION

S Ajajib*, A Droop, K Mankia, P Emery, F Ponchel. Leeds Institute of Rheumatic and Musculoskeletal Medicine, University of Leeds, Leeds, UK

10.1136/annrheumdis-2018-EWRR2018.106

Introduction Palindromic rheumatism (PR) is characterised by recurrent, episodic attacks of articular inflammation, which resolve completely without residual joint damage. Whether PR should be considered as a prodrome of rheumatoid arthritis (RA) or as a distinct syndrome remains unclear, over 70 years since first described.¹

Objectives This project investigates whether the pathogenesis of PR and RA is similar by comparing gene expression profiles from PBMCs in PR, healthy control (HC), and early RA patients.

Methods PBMCs were obtained from drug naïve PR patients during palindromic attack (flare, $n = 10$) and between attacks (non-flare, $n = 12$). 12 age/sex matched, drug naïve RA patients were selected from the RA-MAP project along with 12 HC. The Illumina Human-HT-12/v4 Expression BeadChip microarray and a standard data analysis workflow, were used to establish a list of differentially expressed genes (DEGs). The STRING database was used to characterise functional associations of the top DEGs between groups.² Only the highest confidence (0.900) interactions were selected based on prior information from text-mining, experiments and databases.

Results Comparing RA and HC, STRING analysis revealed 112 interactions (from 378 DEGs); including expected associations with genes such as TNF, AP1 (JUN/FOS), chemokines/receptors and a node including several Interferon responsive genes. Between PR non-flare and HC, 185 interactions (from 458 DEGs) were observed, featuring numerous endoplasmic reticulum and ribosomal proteins, in addition to high IL-1beta/high IL-1R1, lower IL-10 and higher SOCS3. Interestingly, lower Ubiquitin C (UBC) was found to be a prominent feature of many of the identified interactions. Comparing inflammation between RA and PR flare, 176 interactions (from 476 DEGs) included an enhanced endoplasmic/ribosomal signature, lower ubiquitination but higher sumolation and other stress-associated proteins. SCOC3, was no longer present at the top of the DEGs list when both conditions were inflammatory. The IFN signature identified in RA was still highlighted in the associations. Finally, comparing PR non-flare and flare (10 matched pairs), 125 interactions (from 434 DEGs) were observed, primarily from pathways related again to stress, ubiquitination and DNA/RNA processing.

Conclusions Although preliminary, our findings suggest there are distinct gene expression programmes used in PR and RA, and that PR pathogenesis may involve several stress response mechanisms, which are not clearly associated with top DEGs in RA pathogenesis.

REFERENCES

- Hench P, Rosenberg EF. Palindromic rheumatism. A 'new' oft recurring disease of joints (arthritis, peri-arthritis, para-arthritis) apparently producing no articular residues. Report of thirty-four cases; its relation to angio-neural arthrosis', 'allergic rheumatism' and rheumatoid arthritis. *Arch Int Med* 1944;**73**:293-321.
- Szklarczyk D, Franceschini A, Wyder S, Forslund K, Heller D, Huerta-Cepas J, Simonovic M, Roth A, Santos A, Tsafou KP, Kuhn M, Bork P, Jensen LJ, von

Mering C. STRING v10: Protein-protein interaction networks, integrated over the tree of life. *Nucleic Acids Res* 2015 January;43:D447–52.

Disclosure of interest None declared

P091 CLUSTERIN IS INCREASED IN EARLY RHEUMATOID ARTHRITIS AND PREDICTS DISEASE ACTIVITY AND TREATMENT RESPONSE

^{1,2}T Lennerová*, ^{1,2}H Mann, ^{1,2}O Ružicková, ^{1,2}O Šléglová, ¹L Vernerová, ^{1,2}B Šumová, ^{1,2}K Pavelka, ^{1,2}J Vencovský, ^{1,2}L Šenolt. ¹Institute of Rheumatology, Prague, Czech Republic; ²Department of Rheumatology, 1st Faculty of Medicine, Charles University, Prague, Czech Republic

10.1136/annrheumdis-2018-EWRR2018.107

Introduction Clusterin (apolipoprotein J) is a molecular chaperone involved in a number of biological processes, such as inflammation and apoptosis. Recent data suggest its role in the development of autoimmune diseases and bone erosions.

Objectives The aim of this study was to analyse the serum levels of clusterin in patients with early rheumatoid arthritis (RA) and in healthy controls, and to investigate the association of clusterin with disease activity and treatment response.

Methods Clusterin serum levels were measured by ELISA (BioVendor) in 56 patients with early RA before and three months after treatment initiation, and in 56 age-/sex-matched healthy individuals. Disease activity was assessed by 28-joint Disease Activity Score (DAS28), Simplified Disease Activity Index (SDAI) and Clinical Disease Activity Index (CDAI). Treatment response was calculated based on the SDAI/CDAI definition. Receiver operating characteristic (ROC) curve analysis was performed to predict disease activity and treatment response after six months of therapy. Data are presented as mean ±SD.

Results Concentrations of clusterin at baseline were significantly higher in patients with early RA compared with healthy controls (75.1±12.4 vs 56.7±9.7, p<0.001). After three months of therapy, clusterin levels decreased and were comparable to those in healthy subjects (57.7±9.7 vs 56.7±9.7, p>0.05). Although clusterin levels did not correlate with disease activity at baseline, they positively correlated with CDAI and SDAI at month 3 (r=0.294, p=0.030 and r=0.269, p=0.047, respectively) and at month 6 after treatment initiation (r=0.318, p=0.021 and r=0.339, p=0.013, respectively). Using ROC analysis, clusterin baseline levels predicted moderate to high disease activity according to CDAI (AUC=0.829 (95% CI: 0.721 to 0.937), p<0.001) and SDAI (AUC=0.709 (95% CI: 0.548 to 0.869), p=0.019), and major treatment response after 6 months of therapy (AUC=0.696 (95% CI: 0.549 to 0.842), p=0.015 for both).

Conclusions We demonstrate here for the first time increased clusterin levels in patients with early rheumatoid arthritis and propose clusterin as a predictive biomarker for assessing disease activity and treatment response.

Acknowledgements Supported by the project of MHCR for conceptual development of research organisation 00023728, research project SVV 260 373 and project GAUK No. 5 34 217.

Disclosure of interest None declared

P092

MICROARRAY PATHWAY ANALYSIS COMPARING BARICITINIB AND ADALIMUMAB IN MODERATE TO SEVERE RHEUMATOID ARTHRITIS FROM A PHASE 3 STUDY

¹P Emery, ²PC Taylor, ³ME Weinblatt, ⁴Y Tanaka, ⁵EC Keystone, ⁶ER Dow, ⁶R Higgs, ⁶WL Macias, ⁶G Rocha, ⁶TP Rooney, ⁶DE Schlichting, ⁶SH Zuckerman, ⁷IB Mdnnes, ⁸T Holzkaemper*. ¹Leeds Musculoskeletal Biomedical Research Unit, the Leeds Institute of Rheumatic and Musculoskeletal Medicine, Leeds; ²Nuffield Department of Orthopaedics, Rheumatology and Musculoskeletal Sciences (NDORMS), University of Oxford, Oxford, UK; ³Center for Arthritis and Joint Diseases, Brigham and Women's Hospital, Boston, USA; ⁴First Department of Internal Medicine, School of Medicine, University of Occupational and Environmental Health, Kitakyushu, Japan; ⁵University of Toronto, Toronto, Canada; ⁶Eli Lilly and Company, Indianapolis, USA; ⁷Institute of Infection, Immunity and Inflammation, Glasgow, UK; ⁸Presenting on behalf of Eli Lilly and Company, Indianapolis, USA

10.1136/annrheumdis-2018-EWRR2018.108

Introduction In RA-BEAM (NCT01710358), baricitinib (BARI), an oral selective inhibitor of Janus kinase (JAK) 1 and JAK 2, showed significant improvements in patients (pts) with active RA who had an inadequate response to methotrexate compared to placebo (PBO) or adalimumab (ADA).

Objectives To analyse pathways modulated by BARI compared with ADA (both relative to PBO) through 12 wks of treatment.

Methods Pts (n=1307) were randomised 3:3:2 to PBO, BARI 4 mg QD, ADA 40 mg q 2 wks. Total RNA extracted from whole blood drawn at baseline (BL), wk4, and wk12 was analysed using the GeneChip Human Transcriptome Array 2.0 (Affymetrix). Data were analysed using a mixed effects model on a log2 transformed response.

Results There was little overlap of the immune pathways modulated by both BARI and ADA at wk4 with no significant overlap by wk12. BARI downregulated JAK/Signal Transducer and Activator of Transcription (STAT) signalling pathways, like those induced by IFNs, IL-6, GM-CSF, IL-5, and IL-3. Expression of interferon responsive genes (IRGs) was downregulated by BARI and upregulated by ADA. BARI reduced IRGs by 75% at wk4 in pts that had high IFN gene expression at BL. ADA modulated complement pathways. Of interest, STAT transcripts were reduced at wk4 by BARI (STAT1, 2, 3, 5A, 5B, 6); by wk12 several STATs (STAT 1, 2, 5A) did not differ from PBO. Additional differences were noted in the number of genes modulated by each treatment. BARI modulated more genes than ADA at wks 4 and 12; BARI resulted in more gene modulation at wk12 than at wk4, whereas ADA gene modulation was similar at wks 4 and 12. Both the numbers and types of genes modulated by BARI diverged further from ADA at wk12 than at wk4.

Conclusions Gene expression profiling showed significant differences between BARI and ADA treatments. BARI and ADA modulated JAK/STAT or complement pathways, respectively, and the drugs had opposite effects on interferons, indicating different and possibly complementary mechanisms of action of each targeted therapy.

Acknowledgements Study support:

Eli Lilly and Company and Incyte Corporation. Encore of ACRA/ARHP-2017 Annual Scientific Meeting, Nov 4–8, 2016; San Diego, CA, USA.

Disclosure of interest None declared