Objectives To address the limitations of existing studies, and evaluate the effect of a controlled, long-term (24 week intervention, 24 week follow-up), intensive (1 hour physiotherapy twice weekly, and home-exercise for 1 hour 5x weekly), tailored physiotherapy program to improve muscle strength, endurance and deep stabiliser system, and quality of life/disability in cohorts with a substantial number of IIM patients.

Methods All patients fulfilled the Bohan and Peter 1975 diagnostic criteria, had muscle involvement, and were consecutively recruited between 2014–2016. At months 0,3,6,12 all patients were assessed by a physician (physical examination, MITAX, MYOACT, and MDI), and a physiotherapist blinded to intervention (MMT-8 – muscle strength test, FI-2 – muscle endurance test), patients filled out patient reported outcomes (PRO)/questionnaires (HAQ, SF-36, Beck's depression inventory-II (BDI-II), PROs assessing nutrition and fatigue), body composition was analysed using densitometry (iDXA Lunar) and bioelectric impedance (BIA2000-M), and patients provided blood for routine laboratory analysis and biobanking. Normality of data was tested and inter-group analysis performed with 2-way ANOVA and intra-group analysis by Friedman's test with Dunn's post hoc test.

Results 27 IIM patients were recruited into the intervention group (IG) and 27 patients into the control group (CG). Compared to observed statistically significant deterioration in CG over the period of months 0–6, we found statistically significant improvement in FI-2, MMT8, HAQ, BDI-II. Only numerical improvement in IG compared to numerical deterioration in CG, which has not reached statistical significance, was observed in SF-36 and fatigue PROs.

Conclusions Our physiotherapy program led to a significant improvement in muscle strength, endurance, and function. Additionally, patients expressed a decrease in the level of depression. The improvements were clinically meaningful in a substantial proportion of patients.

Acknowledgements Supported by AZV-16–33574A.

Disclosure of interest None declared

P100 THE LINK BETWEEN ANGIOGENESIS AND OSTEOGENESIS IN SPONDYLOARTHRITIS

^{1,2}MH Kaaji*, ^{1,2}JP van Hamburg, ^{3,4}G Kollias, ^{1,2}D Baeten, ^{1,2}L van Duivenvoorde. ¹Amsterdam Rheumatology and Immunology Centre; ²Department of Experimental Immunology, AMC/University of Amsterdam, Amsterdam, Netherlands; ³Division of Immunology, Biomedical Sciences Research Centre 'Alexander Fleming', Vari; ⁴Department of Physiology, National and Kapodistrian University of Athens, Athens, Greece

10.1136/annrheumdis-2018-EWRR2018.116

Introduction Spondyloarthritis is characterised by inflammation, extensive angiogenesis and pathological osteogenesis. Transmembrane (tm) TNF transgenic (tg) mice¹ that overexpress tmTNF exhibit features of SpA, including chronic inflammation and pathological osteogenesis. tmTNF ligation to TNF receptor 2 in endothelial cells (ECs) can induce signal transduction pathways, that may promote these processes. Of note, angiogenesis and osteogenesis are coupled by EC differentiation towards a type H (CD31^{hi} endomucin^{hi}) phenotype.²

Objectives To investigate the link between pathological angiogenesis and osteogenesis in tmTNF tg mice and the potential contribution of noncanonical NF-κB signalling in ECs to this process and infiltration of immune cells into the bone marrow (BM).

Methods Vertebrae and ankles from 6 and 12 weeks and 8 months old tmTNF tg mice or sex-matched non-tg littermates

(n=18) were prepared by cutting 60 μm thick cryosections for confocal imaging.

Results tmTNF tg mice exhibited ectopic osteogenesis which was not observed in non-tg littermates. Immunostainings showed that type H vessels are in the vicinity of the ectopic osteogenesis and osterix⁺ osteoprogenitors. At six weeks of age, osterix⁺ cells are located throughout the ectopic lesion, while at eight months, osterix⁺ cells are only present at the border of the lesion. Furthermore, there is increased osteogenesis and a different vessel architecture within the vertebrae of tmTNF tg mice compared to non-tg littermates that progresses with age. Non-tg littermate vertebrae only have physiological osteogenesis, which is in the metaphysis and periosteum. In addition, tmTNF tg mice also exhibit altered bone marrow (BM) architecture containing extensive lymphoid aggregates, which predominantly consisted of B220⁺ B cells.

Conclusions tmTNF overexpression in mice leads to development of type H vessels associated with ectopic osteogenesis. In addition, extensive lymphoid aggregates develop in the BM. Current studies are aimed at identification of signalling pathways in ECs that contribute to these processes.

REFERENCES

1. Alexopoulou L, et al. *Eur J Immunol* 1997;**27**(10):2588–92.
2. Kusumbe AP, et al. *Nature* 2014;**507**(7492):323–328.

Disclosure of interest None declared

P101 DOT1L INHIBITION INCREASES DERMAL FIBROBLAST PROLIFERATION BUT HAS NO EFFECTS ON IN VITRO OR IN VIVO COLLAGEN DEPOSITION IN MODELS OF FIBROSIS

^{1,2}N Berghen*, ^{1,2}E De Langhe, ^{1,2}R Lories. ¹Department of Development and Regeneration, KU Leuven; ²Division of Rheumatology, UZ Leuven, Leuven, Belgium

10.1136/annrheumdis-2018-EWRR2018.117

Introduction Fibrosis is a key process in Systemic Sclerosis, a devastating disease that is not well understood. The role of epigenetic disease mechanisms is increasingly explored. DOT1L, the unique H3K79-methyltransferase, methylates histone 3 at the Lysine residue at position 79, and regulates gene expression. Inhibition of DOT1L has cell-type specific effects on Wnt-signalling, a pathway suggested to play an important role in fibrosis.

Objectives To study the role of DOT1L in fibrosis.

Methods Primary cell cultures of fibroblasts, isolated from healthy human skin and treated with DOT1L-inhibitor EPZ-5676 or vehicle for 14 days, were stimulated with TGFβ 5 ng/ml after starvation. Smooth muscle alpha 2 actin (ACTA2) gene expression was measured by RT-qPCR. Efficiency of DOT1L-inhibition was analysed with Western Blot for dimethylated H3K79. Picrosirius Red staining measured collagen I and III deposition. Proliferation was analysed with 5-Bromo-2'-deoxy-uridine (BrdU) labelling. 7–9 week old Col1a2; Cre-ERT²; DOT1L^{fl/fl} mice were injected intraperitoneally with tamoxifen (1 mg/day, 5 days) to induce a fibroblast-specific DOT1L knockout. Bleomycin (0.1 mg) or vehicle was injected subcutaneously (5 days/week, 4 weeks). Injected skin was analysed by OH-prolin assay for collagen content, and by histology for dermal thickness measurement.

Results The DOT1L-inhibitor EPZ-5676 reduced H3K79 dimethylation in all samples. TGFβ increased expression of ACTA2