P129 ABSTRACT WITHDRAWN

P130 INCREASE IN GLOBAL DNA METHYLATION AT 3 MONTHS OF METHOTREXATE USE IS NOT ASSOCIATED TO RESPONSE IN EARLY RA PATIENTS

¹HR Gosselt*, ¹MC de Rotte, ²JM Hazes, ³R de Jonge, ⁴SG Heil. ¹Clinical Chemistry, ²Rheumatology, Erasmus Medical Centre, Rotterdam; ³Clinical Chemistry, VU Medical Centre, Amsterdam; ⁴Clinical Chemistry, VU Medical Centre, Rotterdam, Netherlands

10.1136/annrheumdis-2018-EWRR2018.144

Introduction Methotrexate (MTX) is a first-line therapy in early Rheumatoid Arthritis (eRA). However, up to 40% of treated patients do not adequately respond to MTX. MTX interferes with the folate cycle, where it indirectly inhibits the global DNA methylation donor S-adenosylmethionine (SAM). Thus, we hypothesised that global DNA methylation changes during MTX use are associated with treatment response.

Objectives To examine whether there is a change in global DNA methylation ($\Delta\%$ Meth) upon MTX use and if this change is associated to MTX response (Δ DAS28) in eRA patients.

Methods DNA was isolated from whole blood (n=120) and Peripheral Blood Mononuclear Cells (PBMCs, n=83) of eRA patients, before and 3 months after MTX use. Samples were collected from the Treatment in the Rotterdam Early Arthritis Cohort (tREACH), a multicenter, stratified single-blind clinical trial of eRA patients. Selected patients received triple (MTX +SSZ + HCQ) or monotherapy (MTX) combined with corticosteroids. 7 CpG sites within Long-Interspersed Nuclear Elements (LINE-1), a proxy for global DNA methylation, were quantified by Sequenom EpiTyper. Paired t-tests or Wilcoxon Signed Rank tests were conducted to assess a change in methylation. Δ DAS28 score over 3 months was used as a measure for response. Associations between Δ DAS28 and $\Delta\%$ Meth were corrected for baseline DAS28 in a linear regression model.

Results In leukocytes, $\%$ Meth did not significantly change over time. However, in PBMCs $\%$ Meth in CpG1 (Δ Meth=2.61%, p=0.008), CpG2 (Δ Meth=0.73%, p=0.039) and CpG11.12 (Δ Meth=0.56%, p=0.016) significantly increased over 3 months of MTX use after Bonferroni correction. $\Delta\%$ Meth in CpG8.9 was significantly associated to the Δ DAS28 (B=0.29, p=0.039), yet this was no longer significant after Bonferroni correction. $\Delta\%$ Meth was not significantly associated to Δ DAS28 in any of the other LINE1 CpG sites tested.

Conclusions PBMC global DNA methylation in LINE-1 CpG sites increased upon 3 months of MTX use. However, this change in methylation is not associated to MTX response. Further research is needed to investigate the role of global DNA methylation in these patients.

Disclosure of interest None declared

P131 STUDY OF URATE TRANSPORTERS IN PRIMARY GOUT AND HYPERURICEMIA

^{1,2}K Pavelcova*, ^{1,2}L Petru, ¹J Zavada, ¹K Pavelka, ^{1,3}B Stiburkova. ¹Institute of Rheumatology; ²Department of Rheumatology, First Faculty of Medicine, Charles University; ³Department of Paediatrics and Adolescent Medicine, First Faculty of Medicine, Charles University and General University Hospital in Prague, Prague, Czech Republic

10.1136/annrheumdis-2018-EWRR2018.145

Introduction The urate transporters are one of the main genetic determinants of serum uric acid concentrations. Primarily nonsynonymous polymorphisms in *ABCG2* and *SLC2A9* are clinically important for the development of primary gout and hyperuricemia. In this study we investigated the effects of allelic variants of urate transporters in a cohort with primary gout and/or hyperuricemia.

Methods The cohort was recruited in the Institute of Rheumatology, gout was diagnosed in 165 subjects (151 men/14 women) and hyperuricemia in 58 subjects (39 men/19 women). Coding regions of *ABCG2*, *SLC2A9*, *SLC22A11*, *SLC22A8*, *SLC17A3*, and *SLC17A1* genes were amplified and sequenced directly.

Results In *ABCG2* gene, we detected nine non-synonymous variants: p.V12M, p.Q141K, p.R147W, p.T153M, p.K360del, p.F373C, p.T434M, p.S476P and p.D620N. The minor allelic frequency of p.Q141K is 0.23, whereas for the European population it is 0.09. In addition, this variant is associated with early onset of the gout.¹ We identified novel intron variant c.689+1G>A which is associated with two abnormal splicing variants, leading to premature introduction of the stop codon.²

In *SLC2A9* gene, seven missense variants were identified: p.A17T, p.G25R, p.T275M, p.D281H, p.V282I, p.R294H, and p.P350L. In *SLC17A3* gene, the analysis revealed allelic variants p.A100T and p.G279R. Non-synonymous variants p.V202M and p.R343L were found in *SLC22A11*. Only one missense variant (p.T269I) was identified in *SLC17A1*. In *SLC22A8* gene, sequencing revealed three non-synonymous variants p.R149C, p.V448I and p.R513G.

Conclusions Genetic variants of *ABCG2*, common and rare, increased the risk of gout. The precise impact of *SLC22A11*, *SLC22A8*, *SLC17A3*, and *SLC17A1* in the context of hyperuricemia and gout in our cohort is unclear. Although knowledge of renal urate handling has been increased, current evidence is insufficient to fully understand the precise mechanism governing the bi-directional transport of urate.

REFERENCES

1. Stiburkova B, et al. Functional non-synonymous variants of *ABCG2* and gout risk. *Rheumatology (Oxford)* 2017 November 1;56(11):1982–1992.
2. Stiburkova B, et al. Novel dysfunctional variant in *ABCG2* as a cause of severe tophaceous gout: Biochemical, molecular genetics and functional analysis. *Rheumatology (Oxford)* 2016 January;55(1):191–4.

Acknowledgements This study was supported by the grant from the Czech Republic Ministry of Health AZV 15-26693A.

Disclosure of interest None declared

P132 ABSTRACT WITHDRAWN

P133 HLA-ANTIGENS AND DISEASE MANIFESTATION IN A COHORT OF 600 SOUTHERN FRENCH PATIENTS WITH PSORIATIC ARTHRITIS

¹E Massy, ²C Picard, ²C Frassati, ²P Pedini, ³M Martin, ³I Auger, ^{1,3}J Roudier, ^{1,3}N Balandraud*. ¹Rheumatology, Hôpital Sainte Marguerite; ²Etablissement Français du sang, EFS; ³UMRs 1097 Immunogénétique Des Maladies Auto Immunes, INSERM and Aix Marseille University, MARSEILLE, France

10.1136/annrheumdis-2018-EWRR2018.146

Introduction Currently, there is no biomarker available to diagnose psoriatic arthritis (PsOA). Genome wide association studies showed that the majority of PsOA loci are shared with PsOC.¹ However, PsOA has a strong familial predisposition, more so than PsOC as was shown in an Icelandic population.² In the GWAS performed in UK and Germany, the strongest genetic locus is located within the major histocompatibility complex (MHC).^{3,4} No studies have been performed in southern Europe

Objectives The primary objective of our study was to focus on the HLA class I and II alleles found in a cohort of 600 French patients with PsOA clinically well characterised, compared to a control cohort. The secondary objective was to compare two clinical subsets of PsOA, one axial and one peripheral, to test which genotype determines these phenotypes.

Methods 600 patients from the Rheumatology department, St. Marguerite's Hospital, Marseilles, who fulfilled the CASPAR criteria for PsOA, underwent clinical, radiographic and laboratory investigations. HLA Class I and Class II alleles were genotyped. A cohort of 2346 healthy blood donors (HBD) was also tested.

Results

- Comparison between the PsOA population and controls showed one set of alleles significantly associated with PsOA; HLA-B*27, B*21(B*50), C*06 and with a weak significance HLA-A*01, A*25, B*13.
- Within the PsOA population, two genetically different subsets determine two different clinical subtypes: Peripheral disease was significantly associated with -C*06 in disequilibrium linkage with -B*13 and -DR*07, independently associated with HLA-A*01, -B*21 and -B*17. Axial disease was significantly associated with HLA-B*27 in disequilibrium linkage with C*01 and -C*02.

Conclusions This is the first and the largest study ever realised in southern Europe. It lead to conclude that PsOA is genetically heterogeneous. We found that PsOA is divided in two genetically and phenotypically relevant subgroups, one axial group with HLA-B*27 predominance very close to Ankylosis spondylitis and one peripheral group with HLA-C*06 predominance. Other HLA alleles within the MHC are also implied. The clinical utility of HLA typing in PsOA should be further addressed in larger studies.

REFERENCES

1. Stuart PE, Nair RP, Tsoi LC, et al. Genome-wide association analysis of psoriatic arthritis and cutaneous psoriasis reveals differences in their genetic architecture. *Am J Hum Genet* 2015;**97**:816–36. doi:10.1016/j.ajhg.2015.10.019
2. Chandran V, Schentag CT, Brockbank JE, et al. Familial aggregation of psoriatic arthritis. *Ann Rheum Dis* 2009;**68**:664–7. doi:10.1136/ard.2008.089367

3. Rahman P, Elder JT. Genetics of psoriasis and psoriatic arthritis: A report from the GRAPPA 2010 annual meeting. *J Rheumatol* 2012;**39**:431–3. doi:10.3899/jrheum.111242
4. Bowes J, Budu-Aggrey A, Huffmeier U, et al. Dense genotyping of immune-related susceptibility loci reveals new insights into the genetics of psoriatic arthritis. *Nat Commun* 2015;**6**:6046. doi:10.1038/ncomms7046

Disclosure of interest None declared

P134 THE LYMPHATIC SYSTEM PLAYS AN IMPORTANT ROLE IN THE MIGRATION OF PATHOGENIC T-CELLS TOWARDS SYNOVIAL JOINTS AND ENTHESES IN PSORIASIS

¹D Verhaegh, ²E Prens, ¹A Mus, ¹P Asmawidjaja, ¹N Davelaar, ³A Hofman, ³J-B Jaquet, ⁴M Kok, ⁵S Tas, ⁶H Yagita, ¹J Hazes, ¹E Lubberts, ⁴R Bisoendial*. ¹Rheumatology, ²Dermatology, Erasmus University Medical Centre; ³Plastic Surgery, ⁴Rheumatology, Maastad Hospital, Rotterdam; ⁵Rheumatology, Academic Medical Centre, Amsterdam, Netherlands; ⁶Immunology, Juntendo University School of Medicine, Tokyo, Japan

10.1136/annrheumdis-2018-EWRR2018.147

Introduction Psoriasis (PsO) is a common inflammatory skin disease that is characterised by acanthosis, oedema formation, immune cell infiltration, abnormal vascular proliferation, and remodelling of the lymphatic system. Up to ~30% of PsO patients are affected by psoriatic arthritis, a devastating form of arthritis, which leads to disability and deterioration of quality of life. Thusfar, the transition from PsO to PsA, particularly the factors that determine the propensity of pathogenic T-cells in psoriasis to colonise synovial membrane in joints and entheses are poorly understood. Lymphatic endothelial cells (LECs), core components of the lymphatic vascular system and lymph nodes, may control the transmission of pathogenic T-cells in PsO to other sites including synovial joints and enthesal regions by programming these T-cells in terms of their homing properties.

Objectives The focus of our proposal is to investigate whether LECs are capable of modulating the differentiation and homing receptor profile on CD4 memory T-cells.

Methods Human dermal lymphatic endothelial cells (LEC; 0.5×10^4), and fibroblast-like synoviocytes of a patient with PsA (FLS; 1.0×10^4) were pre-incubated for 3 days with PsA synovial fluid (PsA-SF; 0/10/20% v/v). Then, LEC or FLS were co-cultured with 2.5×10^4 allogeneic CD4⁺CD45RO⁺CD25⁻ T-cells that were sorted from peripheral blood mononuclear cells (PBMC) of 3 healthy donors with or without stimulation with α CD3 (0.3 μ g/ml) and α CD28 (0.4 μ g/ml).

Results By means of flow-cytometry based phenotypical analysis, dermal LECs appear to suppress the generation of Th17 cells from stimulated CD4 memory T-cells of healthy donors, as compared to co-culture experiments involving synovial fibroblasts. Initial studies to the underlying mechanisms suggest that blocking the non-canonical NF- κ B pathway downstream of the lymphotoxin beta receptor (LT β R) impairs the capacity of the LECs to suppress Th17 generation. In addition, Delta-like ligand 4 (DLL4) is differentially expressed in LECs and synovial fibroblasts after 3 days of co-culture. Earlier studies have shown that DLL4 may induce ROR γ t expression. Blocking DLL4 using a monoclonal antibody also impairs the capacity of the LECs to suppress Th17 generation.

Conclusions In our co-culture setup, LECs suppress Th17 generation from stimulated CD4 memory T-cells, and are thus directly involved in T-cell differentiation. The LT β R/non-