increase in prevalence of gout, the utilisation of urate lowering agents remained low. Only 1 in 4 patients with gout were prescribed urate lowering agents.

Acknowledgements: The Hospital Authority of Hong Kong

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2018-eular.6398

SAT0377 THE BURDEN OF MONOSODIUM URATE CRYSTALS ASSESSED BY DUAL-ENERGY CT AND ULTRASONOGRAPHY IS NOT CORRELATED TO CARDIOVASCULAR RISK

T. Pascari¹, B. Capon², A. Grandjean¹, J. Legrand², N. Namane², E. Houvenagel¹, L. Norberciak³, J.-F. Budzik². ¹Rheumatology; ²Radiology; ³Biostatistics, Hôpital Saint Philibert, Lomme, France

Background: Gout is associated with higher cardiovascular risk and increases with disease severity. It is not clear if the monosodium urate (MSU) crystal burden is associated with traditional cardiovascular risk factors.

Objectives: The objective of this study was to explore the relationship between the extent of MSU deposition assessed with ultrasonography (US) and dual-energy CT (DECT) and cardiovascular risk.

Methods: Gout patients naive of urate lowering therapy were included in this cross-sectional study to undergo DECT scans for the assessment of total MSU volume deposition of the knees and feet, and US to evaluate the number of joints with the double contour (DC) sign among the femoro-patellar, talo-crural and first metatarsophalangeal joints. Participants were screened for traditional cardiovascular risk factors and levels of the ACC/AHA 10 year-risk for heart disease or stroke was calculated. The primary endpoint was the Spearman correlation coefficient ρ between DECT MSU volume and cardiovascular risk.

Results: A total of 50 patients predominantly male (46/50) aged 62.6 years (± 14.1) were included. Participants had gout duration of 9.5 years (± 11.8), had experienced 4.1 flares (± 6.3) over the past year, had serum urate (SU) levels of 8.1 mg/dL (± 2.3), and 35/50 had at least one US tophus of 1.5 cm³ (± 1.8). The volume of MSU deposits with DECT was 3.9 cm³ (± 11.2) for the feet and 2 cm³ (± 4.4) for the knees. Overall, 28 patients presented with the metabolic syndrome. Correlations between DECT volumes of MSU deposits of the knees, feet, and knees+feet were poor with ρ respectively of 0.23, 0.03 and 0.21. The was no correlation between the number of joints with the DC sign and cardiovascular risk (ρ of 0) and the correlation was poor with SU levels ($\rho=0.09$). Patients with the metabolic syndrome had similar DECT volume of MSU deposits than those without ($p=0.29$).

Conclusions: This study suggests that the association of gout with traditional cardiovascular risk factors is not related to the extent of the monosodium urate crystal burden.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2018-eular.3822

SAT0378 THE TREND OF TREG AND TH17 CELLS CHANGES IN P2X7R-REGULATED ACUTE GOUTY ARTHRITIS MODEL RATS

X.-J. Dai¹, J.-H. Tao¹, X. Fang¹, Q. Lin¹, M.-Y. Li¹, Y. Xia¹, X.-M. Li¹, X.-P. Li¹, Y.-P. Wang². ¹Rheumatology and Immunology, The First Affiliated Hospital of University of Science and Technology of China, Hefei, China; ²Centre for Transplantation and Renal Research, Westmead Institute for Medical Research, The University of Sydney, NSW, Australia

Background: ATP may be the second causative signal for the onset of gout, which acts on P2 $\times$ 7R to regulate the development of acute gouty arthritis.¹ Both regulatory T cells and Th17 cells are important in the development and progression of inflammatory diseases.²

Objectives: To investigate the effect of P2 $\times$ 7R on Treg and Th17 cells in acute gouty arthritis model of rats and its role in acute gouty arthritis.

Methods: Eighty male SD rats were randomly divided into three groups: After establishment of acute gouty arthritis model, rats were given P2 $\times$ 7R agonist ATP, P2 $\times$ 7R inhibitor BBG and PBS, respectively. The rats were sacrificed at 6 hour, 12 hour, 24 hour, 48 hour and 72 hour after treatment. The spleens of the rats were grinded and the expression of Treg and Th17 cells were detected by flow cytometry. Comparison the levels and the ratio of Treg and Th17 cells at the different time points.

Results: (1) The expression levels of Treg and Th17 in the spleen: After treatment at 12 hour, The expression levels of Treg and Th17 in the ATP group were significantly higher than that in the BBG and control groups ($p=0.001$, 0.021 ; $p=0.01$, 0.025); The expression levels of Treg and Th17 in control group were higher than that in BBG group ($p=0.021$, 0.044); There were no significant differences in the three groups at 72 hour after treatment ($p=0.052$, 0.116). (2) The expression trend of Treg and Th17 in different time points: the level of Treg was increased at 6 hour, but decreased gradually at 12 h, 24 hour and then increased at 48 hour again; The level of Th17 was increased at 6 hour, 12 h and 24 hour, but decreased gradually at 48 hour, 72 hour. (3) The ratio of Treg/Th17 gradually decreased in the first three time points, increased at 48 hour and 72 hour in three groups. The ratio of Treg/Th17 in ATP group was lower than that in BBG group and control group at 12 hour, with significant difference ($p<0.05$). But at other four time points, the ratios were no significant differences among the three groups.

Conclusions: Activation of P2 $\times$ 7R decreased the ratio of Treg/Th17 in acute gouty arthritis rat model that showed an acute change trend along with the time, suggesting that P2 $\times$ 7R-regulated the ratio of Treg/Th17 cells affected acute gouty arthritis.

REFERENCES:

- [1] Tao JH, Zhang Y, Li XP. P2X7R: a potential key regulator of acute gouty arthritis. *Semin Arthritis Rheum* 2013;43(3):376–380.
- [2] Noack M. Th17 and regulatory T cell balance in autoimmune and inflammatory diseases. *Autoimmunity reviews* 2014;13(6):668–677.

Acknowledgements: This work was supported by grants from the National Natural Science Foundation of China (81671601).

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2018-eular.6136

SAT0379 VACUUM-ASSISTED CLOSURE VERSUS CONVENTIONAL WOUND CARE IN THE MANAGEMENT OF CHRONIC ULCERS IN PATIENTS WITH TOPHACEOUS GOUT: A PROSPECTIVE ANALYSIS

X. Pan, Z. Huang, M. Liu, D. Li, Z. Huang, Y. Huang, X. Huang, M. Liu, X. Li, X. Yao, Y. Wang, Q. Huang, X. Guo, S. Zheng, T. Li. *Department of Rheumatology and Clinical Immunology, Guangdong Second Provincial General Hospital, Guangzhou, China*

Background: With the rising epidemic of gout, an increasing number of patients suffer from chronic ulcers associated with tophaceous gout in China, causing poor quality of life and disability. Such ulcers are very difficult conditions to deal with. Vacuum-assisted closure (VAC) has been proved to be effective in treating a variety kinds of wounds such as diabetic foot ulcers. However, its use in chronic ulcers associated with tophaceous gout has been seldom reported.

Objectives: In the present study, we evaluated the use of VAC in the treatment of chronic ulcers associated with tophaceous gout in comparison to conventional wound care (CWC).

Methods: We performed a 12 week prospective study that included 13 patients treated with VAC and 14 patients treated with CWC. We collected the clinical outcomes of these patients and data from a satisfaction survey. Chronic ulcers were treated until wound closure, or until the end of 12 weeks. Study will discontinue when the ulcer worsens or remains unchanged by the end of week 4.

Results: Granulation tissue appeared in 12 (92.31%) patients by the end of week 2 in the VAC group, while it appeared in 6 (42.86%) patients by that time in the CWC group ($p=0.013$). 100% granulation was achieved in 11 (84.62%) patients by the end of week 8 in the VAC group as compared to 5 (35.71%) patients by that time in the CWC group ($p=0.018$). By the end of week 12, decreasing in wound size was achieved in 12 (92.31%) patients in the VAC group, while it was achieved in 10 (71.43%) patients in the CWC group ($p=0.326$). Among them, wound closure was achieved in 9 (69.23%) patients in the VAC group, while it was achieved in 3 (21.43%) patients in the CWC group ($p=0.021$). None developed local infection in both groups during the treatment. More patients in the VAC group were satisfied with treatment as compared to the CWC group.