Abstract SAT0384 – Figure 1

Abstract SAT0384 – Figure 2

Conclusions: Respondents tended to believe that ReA may be decreasing and the causes may be changing. In most cases the causative organism was thought to be unknown followed by GI and STIs. Interestingly, full triad ReA was thought to be linked with recurrent ReA. The results suggest several avenues for exploring clinical and epidemiological relationships in ReA.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2018-eular.6087

SAT0385

MULTIPLE INFECTION IS INDEPENDENTLY RELATED TO DEATH IN ADULT PATIENTS WITH HEMOPHAGOCYTIC SYNDROME: ETIOPATHOGENICALLY-DRIVEN MULTIVARIATE ANALYSIS IN 151 PATIENTS

S. Retamozo^{1,2}, P. Brito-Zerón^{1,3}, B. Kostov⁴, P. Moral-Moral⁵, A. Martínez-Zapico⁶, C. Díaz-Pedroche⁷, G. Fraile⁸, P. Pérez-Guerrero⁹, E. Fonseca¹⁰, A. Robles¹¹, M. Vaquero-Herrero¹², M.A. Calvo¹³, M.J. Forner¹⁴, C. Morcillo³, J. Larrañaga¹⁵, M. Rodríguez-Carballo¹⁶, M. Ruiz-Muñoz¹⁷, R. Hurtado-García¹⁸, S. Prieto-González¹, A. Aljibe Aguilar⁵, L. Caminal-Montero⁶, P. Hernández-Jiménez⁷, C. Rodríguez Fernández¹⁹, P. Castro¹, V. Morell Massó², A. Flores-Chavez^{1,20,21}, M. Ramos-Casals¹ on behalf of Reghem-geas-semi Study Group.
¹Hospital Clínic, Barcelona, Spain; ²Inicsa, Unc, Conicet, Córdoba, Argentina; ³Hospital CIMA-Sanitas; ⁴IDIBAPS, Barcelona; ⁵Hospital Universitari i Politècnic La Fe, Valencia; ⁶HUCA, Oviedo; ⁷Hospital Universitario 12 de Octubre; ⁸Hospital Ramón y Cajal, Madrid; ⁹Hospital Puerta del Mar, Cadiz; ¹⁰Hospital de Cabuñes, Gijón; ¹¹Hospital La Paz, Madrid; ¹²Complejo Asistencial Universitario de Salamanca, Salamanca; ¹³Hospital Río Hortega, Valladolid; ¹⁴Hospital Clínico de Valencia, Valencia; ¹⁵Hospital Xeral, Vigo; ¹⁶Hospital Mutua de Terrasa, Barcelona; ¹⁷Hospital Universitario Fundación Alcorcón, Madrid; ¹⁸Hospital Vega Baja, Orihuela; ¹⁹Unidad de Gestión Clínica de Medicina Interna, Hospital Puerta del Mar de Cádiz, Cadiz, Spain; ²⁰Mexican Institute for Social Security (IMSS), Guadalajara; ²¹University Center for Biomedical Research (CUIB), University of Colima, Colima, Mexico

Objectives: To characterise the etiologies and clinical features at diagnosis of patients with hemophagocytic lymphohistiocytosis (HLH) and correlate these baseline features with survival using an etiopathogenically-guided multivariate model.

Methods: The HLH Study Group of the Spanish Group of Autoimmune Diseases (GEAS) was formed in 2013 with the aim of collecting a large series of Spanish adult patients with HLH with substantial experience in the management of patients with systemic diseases. By October 2017, the database included 151 consecutive patients who fulfilled at least 5 of the 8 criteria proposed by the Histiocytosis Society in 2004.

Results: The cohort consisted of 151 patients (91 male, mean age 51.4 years). After a mean follow-up of 17 months, 80 (53%) patients died. With respect to the HLH-dependent variables, adenopathies (HR 0.63, p=0.040), low platelets (HR 3.39, p=0.008), leukopenia (HR 1.81, p=0.047), severe hyponatremia (HR 1.61, p=0.042), disseminated intravascular coagulation (HR 1.87, p=0.034), bacterial infection (HR 1.99, p=0.025), mixed microbiological infections (HR 3.42, p=0.008) and >1 infectious trigger (HR 2.95, p=0.003) were associated with death. Time-to-event analyses for death identified a worse survival curve for patients with neoplasia (p<0.001), mixed microbiological (p=0.019) and >1 (p=0.011) infections and glucocorticoid monotherapy (p=0.021). After adjusting for confounding variables, platelets<100,000/mm³ (HR 2.64), severe hyponatremia (HR 1.88), >1 infectious trigger (HR 3.43) and mixed microbiological infection (HR 2.96) remained significant. Multivariate Cox proportional hazards regression analysis identified >1 infectious trigger (HR 2.60, 95% CI 1.16 to 5.84) as the only variable independently associated with death.

Conclusions: The mortality rate of adult patients diagnosed with HLH exceeds 50%. Infection with >1 microbiological agent was the only independent variable associated with mortality irrespective of the underlying disease, epidemiological profile, clinical presentation and therapeutic management.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2018-eular.4112

SAT0386

ADJUDICATION OF INFECTIONS FROM THE PHARMACOVIGILANCE IN JUVENILE IDIOPATHIC ARTHRITIS PATIENTS (PHARMACHILD) TREATED WITH BIOLOGIC AGENTS AND/OR METHOTREXATE: UPDATE ON RESULTS WITH A FOCUS ON OPPORTUNISTIC INFECTIONS

G. Giancane, J.F. Swart, E. Castagnola, A. Groll, G. Horneff, H.-I. Huppertz, D. J. Lovell, T. Wolfs, M. Hofer, E. Alexeeva, V. Panaviene, S. Nielsen, J. Antón López, F. Uettwiller, V. Stanevich, M. Trachana, F. De Benedetti, L.M. Ailioaie, E. Tsitsami, S. Kamphuis, T. Herlin, P. Doležalová, G. Susic, B. Flato, F. Sztajnbock, E. Fueri, F. Bovis, F. Bagnasco, A. Pistorio, A. Martini, N. Wulffraat, N. Ruperto on behalf of PRINTO. *Clinica Pediatrica e Reumatologia – PRINTO, Istituto Giannina Gaslini, GENOA, Italy*

Background: Pharmachild is a pharmacovigilance registry on children with JIA treated mainly with biologics±methotrexate (MTX). Little evidence exists in literature about the role of JIA or its immunosuppressive therapy in determining infections, especially caused by opportunistic pathogens.