of different AAV in the Polish population seems very similar to other European countries.

	GPA	MPA	EGPA	
Number	417	106	102	
Male/female	210/207	54/52	34/68	*, \$
Mean (median) age (yrs.); range	48,9 (51,4) ¹³⁻⁸⁰	61,5 (63,7) ¹⁹⁻⁸⁵	44,6 (44,5) ⁹⁻⁸⁵	*, \$
Organ involvement				
General n (%)	355 ^(85,1)	83 ^(78,3)	87 ^(85,3)	NS
Skin n (%)	140 ^(33,6)	24 ^(22,6)	51 ⁽⁵⁰⁾	*, \$
Eye n (%)	114 ^(27,3)	4 ^(3,8)	6 ^(5,9)	*, \$
ENT n (%)	315 ^(75,5)	18 ^(17,0)	82 ^(80,4)	*, \$
Respiratory n (%)	315 ^(75,5)	60 ^(56,6)	100 ^(98,0)	*, \$
Heart n (%)	55 ^(13,2)	11 ^(10,4)	43 ^(42,2)	*, \$
GI n (%)	46 ^(11,0)	12 ^(11,3)	25 ^(24,5)	*, \$
Renal n (%)	252 ^(60,4)	101 ^(95,3)	25 ^(24,5)	*, \$
Urinary n (%)	8 ^(1,9)	3 ^(2,8)	0	NS
CNS n (%)	45 ^(10,8)	2 ^(1,9)	8 ^(7,8)	*, \$
Neurological n (%)	75 ^(18,0)	20 ^(18,9)	50 ^(49,0)	*, \$

p<0,05; # – GPA vs MPA, * – GPA vs EGPA, \$ – MPA vs EGPA

Conclusions: This is the first multicenter retrospective study of the Polish patients suffering from with AAV. Demographic characteristics and disease manifestations of AAV patients in POLVAS registry follow the same pattern as those from other European countries.⁴

REFERENCES:

- [1] Jennette JC, et al. *Arthritis Rheum* 2013 Jan;65(1):1–11.
- [2] Padjas A, et al. *Pol Arch Med Wewn* 2014;124(3):143–4.
- [3] Musiał J, Wójcik K. *Pol Arch Intern Med* 2017 Jan 31;127(1):71–72.
- [4] Bajema, et al. *Kidney Int Rep* 2017;1018–1031.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2018-eular.1828

SAT0534

ANNUAL INCIDENCE OF GIANT CELL ARTERITIS IN URBAN AND RURAL AREAS IN WESTERN NORWAY 1972–2012

L.K. Brekke¹, B.-T. S. Fevang², G. Myklebust³. ¹Hospital for Rheumatic Diseases, Haugesund; ²Dept. of Rheumatology, Haukeland University Hospital, Bergen; ³Dept. of Rheumatology, Hospital of Southern Norway, Kristiansand, Norway

Background: Giant cell arteritis (GCA) is the most common vasculitis in adults. The etiology is not fully understood, and environmental factors which may influence the incidence are poorly investigated.

Objectives: To determine the potential influence of rural or urban residence on the incidence of GCA during a 41 year period.

Methods: Hospital-based retrospective cohort study including patients diagnosed with GCA in Bergen Health Area during 1972–2012. Patients were identified through computerised hospital records using the International Classification of Diseases (ICD)-coding system. Clinical information was extracted by review of the patients' medical journals. The patients' residential address was obtained from the population register in Norway. Municipalities were classified as urban (code 1 and 2) or rural (code 3 thru 6) using the Statistics Norway 2017 classification of centrality. The background population data was obtained from Statistics Norway (www.ssb.no). Wilcoxon signed-rank test was used for statistical comparison.

Results: The inclusion process have been published previously.¹ For the computing of incidence 743 patients were included. Among these there were 536 (72%) females (mean age 73.4 years, SD 8) and 207 (28%) males (mean age 72.6, SD 9). 493 patients (66%) had a positive temporal artery biopsy. 484 patients (65%) had a residential address in a municipality classified as urban and 259 (35%) in a rural municipality. The number of inhabitants (aged ≥50 years) in urban areas (i.e. city of Bergen) was 60 910 in 1972 and 81 972 in 2012. The corresponding number of inhabitants in rural areas was 30 320 and 51 401. The overall annual cumulative incidence of GCA was 16.7 (95% CI 15.5–18.0) per 100 000 persons ≥50 years. The mean annual incidence for urban municipalities was 17.1 (95% CI 15.9–18.4) per 100 000 ≥50 years. The corresponding incidence for rural areas was 16.1 (95% CI 14.9–17.3), p=0.46. With regards to biopsy-proven GCA, the overall annual incidence was 11.2 (95% CI 10.2–12.3). In urban and rural areas, the incidence of biopsy-proven GCA was 11.7 (95% CI 10.6–12.7) and 10.4 (95% CI 9.4–11.4) respectively, p=0.10. There were large fluctuations in annual incidence in both urban and rural areas (figure 1).

Conclusions: Annual cumulative incidence of GCA was slightly higher in urban than in rural areas in our study, but the difference was not statistically significant. This is in contrast to a previous study, which found GCA more prevalent in urban than in rural populations.² Further studies are required to determine whether there is a true difference in incidence of GCA in urban versus rural populations, and whether or not exposures to environmental factors may be involved in GCA pathogenesis.

REFERENCES:

- [1] Brekke LK, et al. Incidence of giant cell arteritis in Western Norway 1972–2012: a retrospective cohort study. *Arthritis Research & Therapy* (2017) 19:278. doi:10.1186/s13075-017-1479-6
- [2] Reinhold-Keller E, et al. Giant cell arteritis is more prevalent in urban than in rural populations: results of an epidemiological study of primary systemic vasculitides in Germany. *Rheumatology (Oxford)* 2000;39(12):1396–402.

Disclosure of Interest: L. Brekke Grant/research support from: MSD, B.-T. Fevang Consultant for: Lilly, Novartis, AbbVie, G. Myklebust: None declared

DOI: 10.1136/annrheumdis-2018-eular.3281

SAT0535

EULAR 2018 CORE SET OF DATA TO BE COLLECTED IN GIANT CELL ARTERITIS REGISTRIES AND DATABASES VIEWPOINTS FROM A EULAR TASK FORCE

L. Ehlers¹, J. Askling², J.W. Bijlsma³, M.C. Cid⁴, M. Cutolo⁵, B. Dasgupta⁶, C. Dejaco^{7,8}, W.G. Dixon⁹, N. Feltelius^{10,11}, A. Finckh¹², K. Gilbert¹³, S. Mackie¹⁴, A. Mahr¹⁵, E.L. Matteson¹⁶, L. Neill¹⁷, C. Salvarani¹⁸, W.A. Schmidt¹⁹, A. Strangfeld²⁰, R. van Vollenhoven²¹, F. Buttgereit¹. ¹Department of Rheumatology and Clinical Immunology, Charité University Hospital, Berlin, Germany; ²Department of Medicine, Karolinska Institutet, Stockholm, Sweden; ³Department of Rheumatology and Clinical Immunology, University Medical Centre Utrecht, Utrecht, Netherlands; ⁴Department of Autoimmune Diseases, Hospital Clinic, University of Barcelona, Institut d'Investigacions Biomèdiques August Pi I Sunyer, Barcelona, Spain; ⁵Research Laboratory and Academic Division of Clinical Rheumatology, Department of Internal Medicine, University of Genoa, Genoa, Italy; ⁶Southend University Hospital NHS Foundation Trust, Westcliff-on-Sea, Essex, UK; ⁷Department of Rheumatology, Medical University of Graz, Graz, Austria; ⁸Rheumatology Service, Hospital of Bruneck, South Tyrol Health Trust, Bruneck, Italy; ⁹Arthritis Research UK Centre for Epidemiology, Centre for Musculoskeletal Research, Institute of Inflammation and Repair, University of Manchester, Manchester, UK; ¹⁰Medical Products Agency, Uppsala, Sweden; ¹¹Cross-Committee Task Force on Registries, European Medicines Agency, London, UK; ¹²Division of Rheumatology, Geneva University Hospital, Geneva, Switzerland; ¹³Patient Representative from PMRGCAuk, London; ¹⁴NIHR-Leeds Musculoskeletal Biomedical Research Unit and Leeds Institute of Rheumatic and Musculoskeletal Medicine, University of Leeds, Leeds, UK; ¹⁵Department of Internal Medicine, Hospital Saint-Louis, University Paris 7 – Diderot, Paris, France; ¹⁶Division of Rheumatology and Department of Health Sciences Research, Mayo Clinic College of Medicine, Rochester, MN, USA; ¹⁷Patient Representative from PMR-GCA Scotland, Perth, UK; ¹⁸Division of Internal Medicine, Azienda Ospedaliera-Istituto di Ricovero e Cura a Carattere Scientifico di Reggio Emilia, Reggio Emilia, Italy; ¹⁹Immanuel Krankenhaus Berlin: Medical Center for Rheumatology Berlin-Buch; ²⁰Epidemiology Unit, German rheumatism research Centre, DRFZ, Berlin, Berlin, Germany; ²¹Amsterdam Rheumatology and Immunology Center (ACR), Amsterdam, Netherlands

Background: Giant cell arteritis (GCA) represents the most common form of primary vasculitis and can be associated with severe and potentially life-threatening complications. Due to its low prevalence, systematically collected data on course and outcome of this disease are scarce.

Objectives: The aim of this EULAR Task Force was to identify a core set of data items which can easily be collected from clinicians and facilitates examination of disease course and outcome.

Methods: A multidisciplinary EULAR task force group of 20 experts including rheumatologists, epidemiologists and patient representatives was assembled and breakout groups formed for a meeting at which items from a previously compiled collection of core parameters for a GCA registry were evaluated. Results were presented to the other group members following a structured process for discussion and consensus finding. The meeting was followed by several rounds of discussions to achieve consensus.

Results: A total of 95 items were identified, subdivided into the following categories: General, Demographics, GCA-related signs and symptoms, Other medical conditions, and Treatment. Suitable instruments and assessment intervals were determined for documentation of each item. To facilitate implementation of the recommendations in both primary care and scientifically oriented registers, a minimum core set of parameters was distilled, with supplemental items that can be added optionally depending on the designated purpose of individual registers.

Abstract SAT0535 – Table 1. A minimum core set of parameters to be collected in giant cell arteritis registries and databases

Item	Instrument
General ^{1,2}	
visit date	date
Demographics	
age ³	date of birth
GCA-related Signs & Symptoms ^{4,5}	
cranial ¹	
ocular involvement	
permanent partial visual loss / blindness	y/n (examination)
transient visual loss (amaurosis fugax)	y/n (interview)
diplopia	y/n (interview)
headache	y/n (interview)
Treatment ²	
glucocorticoids ^{6,7}	
current GC use	mg per day in prednisone equivalent, route of administration
tapering ongoing	y/n
recent GC use	
continuous (> 3 months) intake	y/n (interview)
immunosuppressants	
conventional synthetic DMARDs	current medication ^{8,9}
	historical treatment ^{8,9}
biological DMARDs	current medication ^{10,11}

¹ record at baseline
² record during follow-up
³ report with date in case it occurs
⁴ record every 3-6 months; acute changes need to be recorded whenever they occur (e.g. new medication, imaging finding, osteoporotic fracture)
⁵ record every 6-12 months; acute changes need to be recorded whenever they occur (e.g. new medication, imaging finding, osteoporotic fracture)
⁶ record annually; acute changes need to be recorded whenever they occur (e.g. new medication, imaging finding, osteoporotic fracture)

Conclusions: This core set intends to ensure that data from different GCA registries and databases can be compared for the dual purposes of clinical research and improving clinical care, thereby facilitating collaborative analyses.

Disclosure of Interest: L. Ehlers: None declared, J. Askling Grant/research support from: Abbvie, BMS, MSD, Pfizer, Roche, Astra-Zeneca, Eli Lilly, Samsung Bioepis, UCB, J. Bijlsma Consultant for: Roche, SUN, M. Cid Consultant for: Roche, M. Cutolo Consultant for: Mundipharma, Horizon, B. Dasgupta Grant/research support from: Napp, Roche, Consultant for: Roche, Servier, GSK, Mundipharma, Pfizer, Merck, Sobi, Speakers bureau: UCB, Merck, C. Dejaco Grant/research support from: Pfizer, MSD, Consultant for: MSD, Pfizer, UCB, AbbVie, Roche, Novartis, Lilly, Celgene, Merck, Sandoz, GSK, W. Dixon Consultant for: Bayer, N. Feltelius Employee of: Swedish Medical Products Agency, A. Finckh Consultant for: AbbVie, AB2Bio, BMS, Eli-Lilly, MSD, Pfizer, Roche, K. Gilbert Consultant for: PMRGCAuk, S. Mackie Grant/research support from: Roche, GSK, Consultant for: Roche, Sanofi, Chugai, PMRGCAuk, A. Mahr Consultant for: Roche-Chugai, E. Matteson Grant/research support from: Novartis, Bristol Meyer Squibb, Hoffman-La Roche, Genentech, Consultant for: Glaxo-Smith-Kline, Endocyte, L. Neill Consultant for: PMR-GCA Scotland, C. Salvarani Consultant for: Roche, W. Schmidt Consultant for: Roche, GlaxoSmithKline, Sanofi, A. Strangfeld Speakers bureau: AbbVie, BMS, Lilly, MSD, Pfizer, Roche, Sanofi-Aventis, UCB, R. van Vollenhoven Grant/research support from: AbbVie, BMS, GSK, Pfizer, and UCB, Consultant for: AbbVie, AstraZeneca, Biotest, BMS, Celgene, Crescendo, GSK, Janssen, Lilly, Merck, Novartis, Pfizer, Roche, and UCB, F. Buttgerit Grant/research support from: Horizon Pharma, Consultant for: Horizon Pharma, Mundipharma, Roche, Galapagos
DOI: 10.1136/annrheumdis-2018-eular.2428

SAT0536 MULTIPLE RELAPSES IN PATIENTS WITH GIANT CELL ARTERITIS

M. Banjari, E. LIOZON, S. Dumonteil, G. Gondran, S. Palat, H. Bezanahary, S. Nadalon, A.-L. Fauchais, K.H. Ly. *Internal Medicine, CHU Dupuytren Limoges, Limoges, France*

Background: Glucocorticoids (GC) remain the mainstay of treatment for patients with GCA. However, relapses and recurrences (events) occurred in 34% to 74.5%

of the patients.¹⁻³ Most studies compared mainly patients with disease relapses/recurrences to whom without any.

Objectives: To compare patients who had multiple events to those who presented only one in order to identify characteristics and predictive factors of multiple events in patient with GCA

Methods: From 1976 to 2016, we collected prospectively clinical, laboratory, and pathological data, and the treatments and outcomes, of consecutive patients who were diagnosed to have GCA based ACR classification criteria. We compared these data in patients who had more than one event (group 1), either a relapse or a recurrence, to those who had only one (group 2). Relapse was defined as occurrence of clinical symptoms and/or inflammatory parameters, attributable to GCA, which required increased medication during follow-up i.e. under GC therapy while new disease activity after a period of remission was defined as a recurrence.

Results: We included 494 patients for whom data was collected. Forty-seven patients were excluded from our cohort due to lack of data. Among the 447 patients, 228 (51%) presented at least one event. Mean age of this population was 73.3±7 years with female predominance (70%). The median follow-up was 75±55 months. Among these patients 139 (61%) had only one event while 89 (39%) presented more than one. The total number of events was 477 of which 89 recurrences. Events occurred during the first year of treatment in 74% of patients. The majority of these events occurred with a mean prednisone dose of 12.8 ±9 mg/day (median dose 10 mg/d). In univariate analysis, significant differences between both groups is illustrated in table 1. Multivariate analysis showed that ear pain and an elevated ESR were predictive factors of multiple events (OR: 2.45 and 1.93; p=0.04 and p=0.03 respectively). We also found that this risk is lower in patients over 80 years-old (OR: 0.35, p=0.008)

Abstract SAT0536 – Table 1

Characteristic	Group 1 (>1; n=89)	Group 2 (=1; n=139)	P
Age (years)	72.67	75.34	0.011
Cough (%)	20.7	10.9	0.042
Ear pain (%)	19.5	9.4	0.029
PMR (%)	11.3	4.3	0.041
Duration of CS (m)	53.7	26.3	0.0001
Complications of CS(%)	28.4	16	0.024
Steroid sparing treatment (%)	30.7	17.8	0.024
Remission(%)	35.6	62.1	0.0001
Death(%)	23.8	36.4	0.046

Conclusions: Patients with more than 1 event represent 7% to 46% of patients with event¹⁻³ which is consistent with our study (39%). The event occurred most often during the first year of GC therapy in our cohort (74% of patients) while it affects 24% to 50% of patients in other series for the same period.¹⁻³ This difference could be explained by the heterogeneity of GC protocols. We did not found any positive correlation with hypertension, diabetes, and deep vein thrombosis that seemed to be more frequent in patients with multiple events.² In our study, this group of patients appeared younger and presented more often with cough, ear pain, and polymyalgia rheumatica that preceded GCA. Logically, in these patients, corticosteroid therapy was longer and the use of GC-sparing agent was more common. Although getting remission was more difficult in these patients, the long-term prognosis is not poor.

REFERENCES:

- [1] Restuccia, et al. *Medicine* 2016.
- [2] Labarca, et al. *Rheumatology* 2015.
- [3] Kermani, et al. *J Rheumatol* 2015.

Disclosure of Interest: None declared
DOI: 10.1136/annrheumdis-2018-eular.7012

SAT0537 IS BALANCE AFFECTED IN BEHCET'S DISEASE AND WHAT ARE THE FACTORS THAT EFFECT THE BALANCE?

A. Ceilik, M. A. Meilkoğlu. *Department of Physical Medicine and Rehabilitation, Faculty of Medicine, Ataturk University, Erzurum, Turkey*

Background: Behcet's Disease (BD) is a multisystem vasculitis that has a broad range of manifestations. Balance as a complex task may be affected in BD and this may cause postural instability and fall risk.

Objectives: The aim of this study was to determine the fall risk in cases with BD with an objective computerised technique and to evaluate the potential related risk factors for falls in these cases.

Methods: After calculating sample size as 24 (with%95 confidence interval and± %5 standard deviation), 30 patients with BD (according to The Behcet's