REFERENCE:

- [1] Aksentijevich, et al. *N Engl J Med* 2009;360:2426–2437.

Disclosure of Interest: J. Kuemmerle-Deschner Grant/research support from: Novartis, SOBI, Consultant for: Novartis, SOBI, K. Hoernagel: None declared, S. Schlipf: None declared, S. Hansmann: None declared, A. Hospach: None declared, X. Liu: None declared, S. Benseler: None declared, A. Weber: None declared

DOI: 10.1136/annrheumdis-2018-eular.5620

SAT0615

SYMPATHETIC JOINT EFFUSION IN AN URBAN HOSPITAL

J.L. Barlow¹, I.J. Tan². ¹Department of Internal Medicine; ²Department of Internal Medicine; Section of Rheumatology, Temple University Hospital, Philadelphia, USA

Background: Sympathetic joint effusion (SJE) or sympathetic synovial effusion (SSE) is a rheumatologic entity that has not been well defined in the medical literature. It is a non-inflammatory synovial fluid collection that is associated with infection or inflammation of an adjacent anatomic structure.¹ The epidemiology and clinical characteristics of SJE/SSE are largely unknown. This knowledge gap has led to a lack of recognition and misdiagnosis by clinicians.

Objectives: Our study aimed to determine incidence, demographic information, and describe the clinical characteristics and potential triggering conditions for this presumptive reactive phenomenon.

Methods: We conducted a study of patients >18 years of age hospitalised at Temple University Hospital (TUH) between January 31, 2010 and December 10, 2016 who underwent diagnostic arthrocentesis for painful effusions. Individuals with synovial fluid white blood cell count (WBC) in the normal range of 200 WBC/mm³ or less were included. Patients with both non-inflammatory and inflammatory range synovial fluid of greater than 200 WBC/mm³ were excluded to limit confounders. Demographic and clinical data of 72 patients were included for detailed analysis.

Results: SJE/SSE was seen in 80/882 hospitalised patients (incidence of 9%). Seventy-two patients fulfilled inclusion criteria for detailed chart review. Demographic information revealed: male 46/72 (64%), female 26/72 (36%), African-American 38/72 (53%), Caucasian 16/72 (22%), Hispanic 10/72 (14%), undefined and other 8/72 (11%). Onset was typically acute, with 45/72 (63%) of patients developing symptoms within six days of arthrocentesis. All patients (100%) with SJE/SSE presented with painful effusion, and a minority had physical findings of warmth 23/72 (32%) or erythema 12/72 (17%). Interestingly, nearly a third of patients 21/72 (29%) were misdiagnosed with crystal or septic arthritis based solely on clinical exam, and empiric treatment was often administered prior to arthrocentesis. The most commonly affected joint was the knee 61/72 (85%), followed by the elbow 5/72 (7%), shoulder 3/72 (4%) and hip 3/72 (4%). Identifiable pathology in the affected limb was found in 29/72 (40%) of patients. Infection was the most common etiology, found in 17/29 (59%) of patients, and included cellulitis, abscess, osteomyelitis, septic bursitis, myositis, and necrotizing fasciitis. The majority of cases of SJE/SSE 23/29 (79%) were associated with concomitant infection, DVT or intramuscular hematoma in the affected limb which required specific therapeutic interventions.

Conclusions: Sympathetic joint effusion or sympathetic synovial effusion (SJE/SSE) is relatively common in hospitalised patients. SJE/SSE may be a sentinel sign for a more serious disorder affecting the same limb. Clinicians should maintain a heightened index of suspicion for SJE/SSE. A search for underlying infection, venous thrombosis, and intramuscular hematoma in the affected limb is warranted when encountering acute painful joint effusion with normal range synovial fluid WBC count.

REFERENCE:

- [1] Strickland RW, Raskin RJ, Welton RC. Sympathetic synovial effusions associated with septic arthritis and bursitis. *Arthritis Rheum* 1985 Aug;28 (8):941–3.

Acknowledgements: Dr. Daohai Yu, Ph.D.

Dr. Mark Weiner, M.D.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2018-eular.3303

SAT0616

TEN-YEAR RETENTION RATE OF INFLIXIMAB IN PATIENTS WITH BEHÇET'S DISEASE-RELATED UVEITIS

J. Sota¹, C. Fabiani², A. Vitale¹, G. Emmi³, G. Lopalco⁴, V. Venerito⁴, I. Orlando¹, R. Franceschini¹, B. Frediani¹, M. Galeazzi¹, F. Iannone⁴, G.M. Tosi¹, L. Cantarini¹. ¹University of Siena, Siena; ²Humanitas Clinical and Research Center, Milan; ³University of Florence, Florence; ⁴University of Bari, Bari, Italy

Background: To date, a few studies have reported the long-term efficacy of Infliximab (IFX) in Behçet's disease (BD)-related uveitis.^{1,2} Nevertheless, it is known that TNF- α inhibitor drugs may lose their efficacy over time due to the formation of anti-drug antibodies, causing secondary failure and influencing the retention rate of these agents.³ In this regard no previous study has specifically investigated IFX retention-rate in BD-related uveitis.

Objectives: To evaluate the 10 year drug retention rate of IFX in BD-related uveitis, the effect of a concomitant use of disease modifying anti-rheumatic drugs (DMARDs) on drug survival and differences according to the lines of biologic treatment.

Methods: Cumulative survival rates were studied using the Kaplan-Meier plot, while the Log-rank (Mantel-Cox) test was used to compare survival curves.

Results: Forty patients (70 eyes) were eligible for analysis. The drug retention rates at 12-, 24-, 60- and 120 month follow-up were 89.03%, 86.16%, 75.66% and 47.11% respectively. No differences were identified according to the use of concomitant DMARDs ($p=0.20$), while a statistically significant difference was observed in relation to the different lines of IFX treatment ($p=0.014$). Visual acuity improved from baseline to the last follow-up visit ($p=0.047$) and corticosteroids-sparing effect was observed ($p<0.0001$).

Conclusions: IFX retention rate in BD-uveitis is excellent and is not affected by concomitant DMARDs

REFERENCES:

- [1] Keino H, Okada AA, Watanabe T, et al. Long-term efficacy of infliximab on background vascular leakage in patients with Behçet's disease. *Eye (Lond)* 2014;28(9):1100–6.
- [2] Keino H, Okada AA, Watanabe T, et al. Efficacy of Infliximab for Early Remission Induction in Refractory Uveoretinitis Associated with Behçet Disease: A 2-year Follow-up Study. *Ocul Immunol Inflamm* 2017;25(1):46–51.
- [3] Fabbri M, Cantarini L, Caso F, et al. Drug retention rates and treatment discontinuation among anti-TNF- α agents in psoriatic arthritis and ankylosing spondylitis in clinical practice. *Mediators Inflamm* 2014;2014:862969.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2018-eular.1174

SAT0617

COMORBIDITY IN PATIENTS AFFECTED BY AVASCULAR NECROSIS OF THE FEMORAL HEAD AND IMPACT OF COEXISTENT DISEASES ON CONTRALATERAL JOINT INVOLVEMENT

K. Gudkevich¹, N. Martusevich¹, A. Murzich². ¹Belarusian State Medical University; ²RSPC for Traumatology and Orthopedics, Minsk, Belarus

Background: Avascular necrosis (AN) represents a significant problem of modern rheumatology. The highest incidence of this disease is in able-bodied men. The risk of disability among patients with avascular necrosis of the femoral head (ANFH) is high, especially in case of bilateral process, as AN tends to progress steadily and results in osteoarthritis due to untimely diagnosis. Patients with AN can be affected by other coexistent diseases, that can influence the disease presentation and management.

Objectives: This study aimed to analyse the prevalence and pattern of comorbidities in patients affected by ANFH and to determine if there is a differential pattern of comorbidities in patients with bilateral and unilateral ANFH.

Methods: This was a cross-sectional study, comprising 102 adult patients, aged 48,^{38–58} with a diagnosis of ANFH according to Ficat and Arlet criteria. Sociodemographic, anthropometric, clinical parameters of interest were collected from medical records. In order to characterise the burden of coexisting diseases or conditions Charlson Comorbidity Index was calculated. The data obtained were processed with STATISTICA 10.0 using descriptive and nonparametric statistics.

Results: Among a total of 102 patients, a significant prevalence of comorbidities (86,27%) was found. 78,85% of all patients suffered from 2 or more coexistent conditions. The most prevalent comorbidities were: cardiovascular diseases

(CVD) (53.92%), including hypertension (50.0%) and cardiosclerosis (34.31%), endocrine and metabolic disorders (66.67%), including hypercholesterolemia (65.70%) and obesity (32.35%), diseases of the digestive system (30.39%), dor-sopathies (22.55%). The number of comorbidities was associated with age ($p=0.42$, $p<0.001$) and body mass index ($p=0.29$, $p<0.001$).

In patients with bilateral ($n=44$), compared with those affected by unilateral ANFH ($n=58$), prevalence of younger age (43.5 (36–51) vs. 54,^{43–62} $p<0.01$) and male gender (odds ratio (OR) 2.99 (95% confidence interval (CI) 1.28–6.99), $p<0.05$) were detected. A history of (CVD) was more frequent in patients with unilateral ANFH (63.79% vs. 40.91%, OR 3.62 (95%CI 1.67–7.88), $p<0.05$), as well as hypertension (62.07% vs. 34.09%, OR 3.16 (95%CI 1.40–7.17, $p<0.01$). Patients with unilateral ANFH were more likely to have higher number of comorbidities (3.80^{1,17–6.43} vs. 2.89 (0.73–5.05)) and Charlson Comorbidity Index (0.72 (0.43–1.47) vs. 0.52 (0.43–1.87)).

Conclusions: ANFH is associated with high prevalence of comorbidities, especially CVD.

CVD, including hypertension, were more likely to be found in patients with unilateral ANFH, as well as higher comorbidity burden. It can be explained by the fact that patients with CVD received treatment according to national guidelines, including antihypertensive drugs, anticoagulants, statins, etc. This indicates that performing secondary prevention of CVD can be important in both CVD and ANFH, as such treatment can influence on intraosseous blood circulation in the contralateral joint.

REFERENCE:

- [1] Valderas JM, Starfield B, Sibbald B, Salisbury C, Roland M. Defining comorbidity: implications for understanding health and health services. *Ann Fam Med* 2009;7:357–363.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2018-eular.7405

SAT0618

RELATIONSHIP BETWEEN ARTERIAL STIFFNESS AND DISEASE DURATION IN BEHCET'S DISEASE

K. Ayar¹, B. Sensoy², M.E. Aslancı³, T. Teker⁴, S. Cekiç⁴. ¹Rheumatology, University of Health Sciences, Bursa Yüksek İhtisas Training and Research Hospital; ²Cardiology; ³Ophthalmology; ⁴Internal Medicine, University of Health Sciences, Bursa Yüksek İhtisas Training and Research Hospital, BURSA, Turkey

Background: There are few studies in the literature investigating arterial stiffness in Behcet's Disease (BD). The association of arterial stiffness with duration of disease is unknown.

Objectives: The aim of this study is to compare arterial stiffness by assessing pulse wave velocity (PWV) and augmentation index (AI) measurements in healthy controls (HC) and patients diagnosed as BD with low and high disease duration.

Methods: Seventy BD patients and 40 HC without diabetes mellitus, Hypertension or chronic renal failure were included in the study. Patients with BD who were diagnosed within 12 months were taken to the group with low disease duration (LDD) and the others to the group with high disease duration (HDD). Demographic data and lipid parameters of all participants were recorded and PWV and AI measurements were made from brachial artery. The clinical characteristics of the patients with BD were recorded and the Behcet's Syndrome Activity Scale (BSAS) questionnaire was filled in for all patients with BD. Findings were compared between LDD, HDD and HC groups.

Results: Age, gender and body mass indexes were not statistically different between groups ($p=0.134$, $p=0.948$, $p=0.929$ respectively). Augmentation index and PWV measurements were different between groups ($p=0.014$, $p=0.024$ respectively). In the HDD group, both PWV and AI were significantly higher than HC ($p=0.013$, $p=0.006$ respectively). In LDD group AI was significantly higher than HC however PWV was not ($p=0.047$, $p=0.919$ respectively). There was moderate correlation between PWV and disease duration (Rho:0.393; $p<0.001$) and there was no correlation between AI and disease duration ($p=0.287$)

Abstract SAT0618 – Table 1. Demographic data, parameters of arterial stiffness, lipid parameters in all participants and disease duration, BSAS scores in patients with BD.

	Healthy controls (n=40)	The time since the diagnosis of BD		P
		LDD ¹ (12 months <=) (n=32)	HDD ² (12 months <) (n=38)	
Age (year)	38.35±8.99	37.16±8.89	41.37±9.38	0.134
Gender (f/m)	23/17	20/12	19/19	0.948
BMI ³ (kg/m ²)	26.55±3.65	26.33±3.68	26.22±4.38	0.929
BSAS ⁴ (0-100)		42.97±21.59	42.87±22.83	0.985
Disease duration (month)		0 (0-6.7)	120 (36-186)	<0.001
Augmentation index (%)	22.53±7.80	26.63±9.72	28.87±11.24	0.014
Pulse wave velocity (m/s)	5.75 (5.3-6.4)	5.8 (5.3-6.5)	6.6 (5.6-7.0)	0.024
LDL (mg/dl)	124.25±29.99	114.89±29.41	115.80±27.78	0.593
T. Cholesterol (mg/dl)	204.25±36.80	191.82±42.66	188.80±29.62	0.426
Triglycerides (mg/dl)	105.62±35.46	127.82±77.22	125.40±58.38	0.507
HDL (mg/dl)	54.75±10.34	52.35±13.62	47.65±10.71	0.182

LDD: Low disease duration, HDD: High disease duration, BMI: Body mass index, BSAS: Behcet Syndrome Activity Scale, BD: Behcet Disease

Conclusions: The augmentation index is higher in patients with BD than HC regardless of the duration of the disease. Pulse wave velocity does not increase in the early period of the disease in patients with BD but increases with prolonged disease duration.

REFERENCES:

- [1] Yilmaz S1, Celik G, Esmen SE. Assessment of arterial stiffness in patients with inactive and active Behcet's disease. *Scand J Rheumatol*. 2014;43 (1):63–9. doi:10.3109/03009742.2013.809787. Epub 2013 Sep 9
- [2] Yıldırım A1, Karakaş MS2, Kılınc AY3, Altekin RE3, Yalçınkaya AS3. Evaluation of arterial stiffness and subclinical atherosclerosis in patients with Behcet's disease without cardiovascular involvement. *Türk Kardiyol Dern Ars* 2016;Oct;44(7):575–581. doi:10.5543/tkda.2016.06944

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2018-eular.2823

SAT0619

ASSOCIATIONS BETWEEN CLINICAL MANIFESTATIONS OF BEHCET'S SYNDROME AND WORK OUTCOMES: RESULTS FROM A UK CROSS-SECTIONAL ANALYSIS

L. Chadwick^{1,2}, N.J. Goodson^{1,2}, R.J. Moots^{1,2}. ¹Department of Rheumatology, Aintree University Hospital; ²Institute of Ageing and Chronic Disease, University of Liverpool, Liverpool, UK

Background: Behcet's syndrome (BS) is a multi-systemic auto-inflammatory disease which exhibits significant heterogeneity in clinical manifestations, including orogenital ulceration, skin rash, arthralgia and ocular, venous, gastro-intestinal and neurological involvement. BS affects adults of working age, usually presenting between 20–40 years,¹ and can therefore have a significant impact on work. The association between individual and cumulative clinical manifestations of BS on employment status has not previously been assessed.

Objectives: To 1) describe demographics, clinical manifestations, work outcomes and use of unemployment benefits in a UK BS cohort and 2) to explore the relationship between individual and cumulative manifestations of BS and work outcomes.

Methods: A cross sectional analysis was performed using the Liverpool Behcet's Centre of Excellence clinical database. Inclusion criteria were clinical characteristics meeting International Criteria for Behcet's Disease (ICBD) diagnostic criteria and recorded employment data. Patients meeting ICBD but thought unlikely to have BS on clinical review by our multi-disciplinary team were excluded. Demographics, clinical manifestations, work outcomes and use of employment related benefits were described. A binomial variable 'Out of work' was generated and defined as unemployment, sickness absence, or patients marked as retired but of working age, a carer or a homemaker if the patient had to stop work due to BS. Associations between Out of work and: 1) individual clinical manifestations of BS, and 2) number of clinical BS manifestations, were explored using multivariate logistic regression adjusted for age, gender, mean EQ5D score and socio-economic status. Odds ratios (OR) and 95% confidence intervals were calculated.

Results: 120 patients met inclusion criteria. Mean age was 41.1 years and 33 patients (28%) were male. The minimum number of clinical manifestations were two in order to meet ICBD; the frequencies were 100% oral ulceration, 94% genital ulceration, 71% arthralgia, 43% skin rash and 20% ocular, 7% neurological, 6%