Cr^{3+} , we found that bivalent cobalt ions and trivalent chromium ions had different effects on osteoblasts. While Co^{2+} reduced the secretion of collagen type 1 in osteoblast-like cells, Cr^{3+} did not impair the collagen type 1 production. The secretion of this protein was partially attributable to a reduced gene transcription. In contrast, the mineralization was not impaired by Co^{2+} . However, Cr^{3+} at a concentration of 250 μM resulted in a decrease of mineralisation by 89% compared to controls. Even at the concentration of 50 μM a reduced mineralization by 34% was observed. The increase of the phosphate source resulted in an increased mineralization, suggesting that the capture of phosphate by Cr^{3+} (formation of CrPO_4), which is required for the mineralization could explain the inhibitory effect of chromium.

Conclusions: Our data suggest that Co^{2+} and Cr^{3+} ions affect bone formation at various stages of differentiation. While Co^{2+} ions impair the formation of the bone organic matrix, Cr^{3+} ions affect the deposition of inorganic minerals. Whether the inhibitory effect of chromium ions on the mineralization is caused by soluble Cr^{3+} ions itself or by solid CrPO_4 needs to be clarified in further studies.

Acknowledgements: The study was supported by Stiftung Endoprothetik (S01/16).

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2018-eular.5534

Rheumatoid arthritis – etiology, pathogenesis and animal models

AB0108 RELATIONSHIP BETWEEN PROGRANULIN AND MUSCULOSKELETAL ULTRASOUND IN RHEUMATOID ARTHRITIS

A.A. Negm^{1,2}, M.H. Nassr³, H.M. Fathi³, N.A. Fouad³, O. Zaki⁴, S.H. Senara³.

¹Rheumatology unit, Dubai Hospital, Dubai Health Authority, Dubai, United Arab Emirates; ²Rheumatology, Faculty of Medicine, Al-Azhar University, Cairo;

³Rheumatology; ⁴Clinical Pathology, Faculty of Medicine, Fayoum University, Fayoum, Egypt

Background: Progranulin (PGRN) is a pluripotent, secretory growth factor, mainly present in cells of the epithelium, central nervous system and immune system. PGRN known to directly regulate regulatory T cells and recently found to be upregulated in Rheumatoid Arthritis (RA) and related to its activity. No attempts was made to explore the relation between PGRN and RA activity measured by clinical and Ultrasound score (USS).

Objectives: To explore PGRN levels in the serum of patients with RA, and its correlation with disease activity assessed clinically combined by ultrasonography (USS).

Methods: RA patients: 52 RA patients were included in this study. They were consecutively recruited from the outpatient clinic of Rheumatology Department for regular follow-up at Fayoum University Hospitals. Patients were classified according to the 2010 ACR/EULAR criteria. Control group: 20 age and sex matched healthy volunteers were recruited as controls for level of serum PGRN. Informed consent was obtained from all patients and controls for inclusion in the study.

All patients were subjected to full history taking, and clinical examination. Assessment of disease activity by modified DAS-28 score. Progranulin is measured by an ELISA Kit (Elabscience), this kit uses sandwich-ELISA as the method.

The ultrasound examination was performed with GE LOGIQ 7 PRO equipment using a linear transducer with 18 MHz frequency. The patients were examined according to the German US7 score in 7 joint areas.

Results: Demographic and clinical data of the RA group: Females represents 90.4% (47 of 52). Age ranged from 21–72 years, mean age was 42.8 ± 10.4 years. Disease duration ranged from 1 to 30 years, median 4 years. Their mean ESR level was 36.4 ± 16 (Range: 5–70). The median of Ultrasound score assessment was 8 while the mean was 8.73 ± 6.9 (Range: 0–33).

Serum PGRN concentrations in RA patients and healthy controls: The median of serum PGRN levels in RA patients was 65 ng/ml (mean, 92.54 ± 45.4) which is much higher in comparison to those in normal controls, median 31 ng/ml (mean, 32.74 ± 9.2); the difference showed high statistical significance, $p < 0.001$.

Correlation analysis: The relationships between PGRN levels and other continuous variables were analysed using Spearman's rank correlation. There was high statistically significant correlation between PGRN level and USS score ($r = 0.37$, $p = 0.006$). Also, there was high statistically significant correlation between PGRN level and ESR ($r = 0.67$, $p < 0.000$) and between PGRN level and DAS 28 ($r = 0.74$, $p < 0.000$). There was high statistically significant correlation between USS score and ESR level ($r = 0.54$, $p < 0.000$) and between USS score and DAS 28 ($r = 0.68$, $p < 0.000$).

Conclusions: The serum Progranulin levels were higher in the serum of RA patients than age and sex matched controls. It is positively correlated with diseases activity measured by DAS 28, ESR and Ultrasound activity measured by

German US7 score. Serum PGRN levels may be a useful biomarker in RA disease. Ultrasound positively correlated with ESR and DAS 28 in our cohort of RA patients.

REFERENCE:

- [1] Chen J, Li S, Shi J, Zhang L, et al. Serum progranulin unrelated with Breg cell levels, but elevated in RA patients, reflecting high disease activity. *Rheumatology international* 2016 Mar 1;36(3):359–64.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2018-eular.6984

AB0109 CHARACTERISATION AND COMPARISON OF SERUM BIOMARKERS IN PATIENTS WITH RHEUMATOID ARTHRITIS VERSUS OSTEOARTHRITIS USING MASS SPECTROMETRY

A. Circiumaru^{1,2}, C. Butnaru³, M. Bojinca¹, S. Petrescu⁴. ¹Department of Internal Medicine and Rheumatology, "Dr. I. Cantacuzino" Clinical Hospital; ²Department Molecular Biology of the Cell; ³Department of Bioinformatics and Structural Biochemistry; ⁴Institute of Biochemistry, Romanian Academy, Bucharest, Romania

Background: There is increasing interest in biomarkers in medicine, especially for their use as a diagnostic tool and in disease outcome prognosis. Rheumatoid arthritis (RA) is a chronic autoimmune disease with important debilitating outcome potential, therefore a biomarker panel may improve early diagnosis meaning better therapeutic management and lower social costs. Recently, mass spectrometry has emerged as one of the leading methods to identify new biomarkers for different pathologies as well as being able to determine their potential interactors in their cellular pathway.

Objectives: To characterise biomarkers using mass spectrometry, from sera of patients with RA and make a comparison between RA, osteoarthritis (OA) and healthy controls.

Methods: Blood was collected from 26 subjects, 12 patients with RA, 12 patients with OA, and 2 HC, centrifuged and sera was collected in dedicated tubes. Sera were separated by SDS-Page electrophoresis and stained with Coomassie staining. The samples were subjected to in gel digestion followed by LC/MS analysis. The obtained spectra were searched with Proteome Discoverer v1.4 using SEQUEST algorithm.

Results: A total of 985 proteins were identified, from which 323 were specific to the RA study group and 159 were characteristic to the OA group. Prevalent peptides found in the RA group were EGF-containing fibulin-like extracellular matrix protein 2, isoform DeltaLf of lactotransferrin, BPI fold-containing family B member 1 and isoform 2 of BPI fold-containing family A member 1. In the OA study group there was a prevalence in homeobox protein TGIF1, major facilitator superfamily domain-containing protein 9, methyltransferase-like protein 23, a fragment of the salivary acidic proline-rich phosphoprotein, serine/threonine-protein phosphatase 4 regulatory subunit 1, zinc finger protein 229.

Conclusions: Our results show a prevalence of peptides linked to cell migration, proliferation and activation in the RA sera as compared to peptides related to cell turnover and tissue senescence observed in OA sera. We believe further mass spectrometry studies of biomarkers in larger cohorts to be a promising tool for diagnosis and outcome prognosis in RA.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2018-eular.3918

AB0110 SERUM AND SALIVARY IGA1 AND IGA2 ACPA SUBCLASSES IN ESTABLISHED RHEUMATOID ARTHRITIS AND ASSOCIATIONS TO SMOKING

A. Svärd¹, K. Roos Ljungberg^{1,2}, A. Kastbom², T. Skogh², K. Martinsson². ¹Center for Clinical Research Dalarna, Falun; ²Department of Clinical and Experimental Medicine, Linköping University, Linköping, Sweden

Background: A prevailing hypothesis regarding the pathogenesis of rheumatoid arthritis (RA) involves an association between mucosal immunity and the development of RA. A pathogenetic role for antibodies against citrullinated peptides (ACPA) is assumed, and there are indications of mucosal immunisation in the lungs, the oral mucosa and the intestinal mucosa preceding overt RA.¹

IgA antibodies are produced both locally, at mucosal membranes, and systemically. Among the two IgA subtypes, IgA1 dominates in serum IgA2 in mucosal secretions.

We have previously reported the presence of ACPA of IgA isotype in both serum and saliva of RA patients,² and that IgA ACPA is associated with cigarette smoking.³