declared, H. Hammer: None declared, I. Olsen: None declared, T. Uhlig: None declared, D. van der Heijde: None declared, T. Kvien: None declared, E. Haavardsholm Grant/research support from: Pfizer, UCB, Roche, MSD, and AbbVie
DOI: 10.1136/annrheumdis-2018-eular.5318

OP0151
ROUTINELY RECORDED MUSCULOSKELETAL ULTRASOUND FINDINGS IMPACT CLINICIANS' DIAGNOSTIC BEHAVIOUR MAXIMALLY IN AUTOANTIBODY-SERONEGATIVE PATIENTS ATTENDING AN EARLY ARTHRITIS CLINIC

K. Iqbal^{1,2}, D.W. Lendrem^{1,2}, B. Hargreaves^{1,2}, B. Thompson³, J.D. Isaacs^{1,2}, A. G. Pratt^{1,2}. ¹National Institute for Health Research Newcastle Biomedical Research Centre, Newcastle upon Tyne Hospitals NHS Foundation Trust and Newcastle University; ²Institute of Cellular Medicine, Newcastle University; ³Musculoskeletal Directorate, Newcastle upon Tyne Hospitals NHS Foundation Trust, Newcastle upon Tyne, UK

Background: Musculoskeletal ultrasound (MSUS) is a popular assessment tool, but its contribution to diagnostic practice over and above standard clinical and laboratory parameters has proved difficult to quantify.
Objectives: A published 7-joint ultrasound algorithm¹ has been adapted for pragmatic application during 15 min screening appointments forming part of initial patient assessments in the Newcastle Early Arthritis Clinic (NEAC). Its additive contribution to diagnostic classification in this routine setting was appraised.
Methods: Detailed baseline clinical and laboratory parameters were recorded. Semi-quantitative MSUS scoring (0–3, grey scale and power Doppler) of the most symptomatic wrist (midline and ulnar dorsal longitudinal views), 2nd/3rd MCPs and PIPs and 2nd/5th MTPs (all longitudinal views) was recorded by sonographers, along with the 'sonographer's scan impression' ('definitely inflammatory,' 'possibly inflammatory' or 'non-inflammatory'). All MSUS findings were available to rheumatologist diagnosticians during subsequent consultations, and persistent inflammatory arthritis (PIA) was classified only where patients were started on ≥ 1 disease modifying anti-rheumatic drug (DMARD). Stepwise multiple logistic regression was employed to identify clinical variables that independently predicted IA diagnosis; the additive contribution of MSUS parameters to resultant models was assessed by comparing areas under receiver operator characteristic curves (AU ROCs).

Abstract OP0151 – Figure 1 ROC curves depicting additive discriminatory utility of MSUS 'scan impression' with clinical parameters alone (red), with respect to PIA diagnosis amongst seronegative early arthritis patients. Δ AU AOC=0.08; $p<0.001$

Results: 847 patients were enrolled (17% seropositive for anti-citrullinated peptide autoantibody, ACPA); final outcomes of PIA were recorded in 29% and 18% of the overall and ACPA-seronegative cohorts, respectively. SJC, CRP, age and ACPA status were non-redundant clinical/laboratory predictors of a PIA diagnosis by consulting rheumatologists in the overall cohort (AU ROC 0.85; 95% CI: 0.81 to 0.88), their discriminatory utility being diminished in the seronegative sub-cohort (AU ROC 0.78; 95% CI: 0.72 to 0.82). Although the additive contribution of

summed parameters from the 7-joint MSUS algorithm to the model was statistically significant ($p<0.001$) it was numerically small (Δ AU ROC 0.03 and 0.05 in the overall and seronegative cohorts, respectively). The 'sonographer's scan impression' was a potentially more useful diagnostic tool, its additive contribution to diagnostic outcome over clinical parameters alone being most evident in ACPA-negative patients where it increased the AU ROC by 10% (Δ AU ROC 0.08; $p<0.001$; figure 1).

Conclusions: In this large, un-blinded observational study, the clinical utility of a 15 min MSUS screen for diagnosing PIA requiring DMARDs was particularly evident amongst ACPA negative patients attending an EA clinic.

REFERENCE:
[1] Backhaus, et al. *Arthritis Rheumatol* 61:1194–201.

Acknowledgements: National Institute for Health Research Newcastle Biomedical Research Centre
Disclosure of Interest: None declared
DOI: 10.1136/annrheumdis-2018-eular.3893

OP0152
IS SONOGRAPHIC PHENOTYPE OF LATE-ONSET RHEUMATOID ARTHRITIS DIFFERENT FROM YOUNG-ONSET RHEUMATOID ARTHRITIS? RESULTS FROM THE BIRMINGHAM EARLY ARTHRITIS COHORT

R. Singh^{1,2}, I. Sahbudin^{1,2}, M. Maybury¹, K. Raza^{1,3}, A. Filer^{1,2}. ¹Rheumatology Research Group, Institute of Inflammation and Ageing, University Hospitals of Birmingham; ²Department of Rheumatology, University Hospitals Birmingham NHS Foundation Trust, Queen Elizabeth Hospital Birmingham; ³Department of Rheumatology, Sandwell and West Birmingham Hospitals NHS Trust, Birmingham, UK

Background: The incidence of rheumatoid arthritis (RA) is generally seen as a bimodal age distribution, which consists of young-onset RA (YORA) and late-onset RA (LORA). Although a retrospective study has reported some differences in sonographic changes between LORA and YORA, little is known whether there are any differences in the clinical and sonographic phenotypes between these two groups during the early disease phases.
Objectives: To compare the clinical and sonographic characteristics between YORA and LORA during the early phases of disease.
Methods: DMARD-naïve patients with clinically apparent synovitis of at least one joint and symptom duration of three months or less were included in the analysis. Patients underwent clinical, ultrasonography and radiological assessments at baseline and final outcomes were determined at 18 months; patients were classified as having RA (according to either the 1987 ACR and/or 2010 ACR/EULAR criteria), a non-RA persistent arthritis or a resolving arthritis. Sonographic assessment included MCP, PIP, wrist, MTP, knee, ankle, elbow joints and wrist, hand flexor, bicep tendon, anteromedial-lateral tendon compartments. The presence and absence of joint synovitis and tenosynovitis were recorded according to the EULAR/OMERACT consensus definition.

	Young-onset RA (n=37)	Late-onset RA (n=36)	Non-RA persistent arthritis (n=27)	Resolving arthritis (n=50)	P (YORA vs LORA)
Age (years)	50 (45–56)	67 (63–73)	39 (31–60)	46 (35–59)	<0.001
EMS (mins)	75 (30–120)	120 (60–240)	60 (10–180)	30 (0–68)	0.808
DAS-28 CRP	5.02 (3.60–5.73)	5.11 (4.49–6.10)	3.77 (3.02–4.95)	3.35 (2.98–4.43)	1.00
RF%					
High	15 (40.5)	10 (27.8)	1 (3.7)	2 (4)	0.568
Positive					
Low	5 (13.5)	7 (19.4)	4 (14.8)	1 (2)	
Positive					
Negative	17 (45.9)	19 (52.8)	22 (81.5)	47 (94)	
ACPA %					0.722
High	17 (45.9)	14 (38.9)	2 (7.4)	2 (4)	
positive					
Low	1 (2.7)	2 (5.6)	0 (0.0)	0 (0.0)	
positive					
Negative	19 (51.4)	20 (55.6)	25 (92.6)	48 (96.0)	
Symptom duration (weeks)	8 (4–10)	8 (6–10)	6 (4–8)	5 (4–8)	1.00
Bilateral MTP squeeze, n (%)	22 (59.5)	10 (27.8)	4 (14.8)	10 (20.0)	0.09

Results: 150 patients were included in the analysis. At 18 months, 37 patients developed YORA, 36 developed LORA, 27 developed non-RA inflammatory arthritis and 50 patients had resolving arthritis. The clinical characteristics

between YORA and LORA were not significantly different at initial presentation (table 1). The ultrasound characteristics differed between these two groups. LORA patients were more likely to have shoulder biceps tendon tenosynovitis (GS; $p=0.026$, PD; 0.037), elbow joint synovitis (GS; $p=0.010$, PD; $p=0.037$), MCP1 (PD; $p=0.032$) and MCP5 (GS; $p=0.035$) synovitis, compared to YORA patient. YORA patients were more likely to have MTP synovitis (GS; $p=0.013$) compared to LORA patients.

Conclusions: This is the first study to describe the difference of both clinical and sonographic inflammation of YORA and LORA in recent-onset DMARD-naïve RA patients in a longitudinal study. There are differences in US-detected joint and tendon inflammation despite similarities in clinical characteristics. The prognostic value of the differences in US pathology between these two groups should be further explored.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2018-eular.5805

OP0153

RESOLUTION OF SYMPTOMS OF PATIENTS WITH CLINICALLY SUSPECT ARTHRALGIA IS PARALLELED BY DISAPPEARANCE OF MRI-DETECTED SUBCLINICAL INFLAMMATION; A LONGITUDINAL STUDY

R.M. Ten Brinck, D.M. Boeters, H.W. van Steenberg, A.H. van der Helm-van Mil. *Rheumatology, LEIDEN UNIVERSITY MEDICAL CENTRE, Leiden, Netherlands*

Background: It is known that MRI-detected subclinical inflammation in patients with Clinically Suspect Arthralgia (CSA) precedes the development of clinically evident Inflammatory Arthritis (IA). However, there are no longitudinal studies in CSA patients that did not progress to IA. The course of joint symptoms in these patients is unknown. Furthermore, it is unknown if subclinical inflammation resolves spontaneously in parallel with disappearance of joint symptoms.

Objectives: To determine the course of joint symptoms and of MRI-detected subclinical joint inflammation, and mutual time relationships, in patients that presented with CSA but did not progress to IA during 2 year follow-up.

Methods: Between April 2012-April 2014, 149 patients were included in the CSA cohort. These patients had no clinically evident arthritis but recent-onset (<1 year) arthralgia of small joints that was clinically considered at risk for IA by the rheumatologist. None of the patients were treated with DMARDs. Patients underwent MRI at baseline and when arthritis developed or after 2 years (when follow-up ended). The patients that did not progress to IA within 2 years and had serial MRIs, as well as longitudinal clinical data and questionnaires (e.g. the HAQ), were studied ($n=60$). 1.5T MRIs of wrist, MCP2-5, and MTP1-5 joints were scored for synovitis, tenosynovitis and bone marrow oedema (BME) (summed in the total MRI inflammation score) according to RAMRIS, blinded for order in time and clinical information. Paired t-tests were used.

Results: After 2 year follow-up, 20 patients (33%) had complete resolution of symptoms whereas 40 patients (66%) did not. The remaining symptoms in the latter group were diagnosed as: persistent inflammatory arthralgia ($n=26$), osteoarthritis ($n=6$) and tendinomyalgia ($n=8$). In the total group, the mean total MRI-inflammation score decreased from 3.2 at baseline to 2.5 at 2 years ($p=0.046$). When the patients with persistent symptoms were studied, there were no significant changes in MRI-inflammation scores. Analysing the patients with resolution of symptoms separately, the total MRI-inflammation score decreased significantly (3.3 to 2.3; $p=0.019$). Also the total synovitis and tenosynovitis scores decreased significantly (1.3 to 0.82 and 0.83 to 0.30, respectively; $p=0.043$ and $p=0.029$), whereas BME scores remained unchanged (1.2 and 1.2; $p=0.87$). After 2 year follow-up the absolute MR-scores of these patients were compared to data from age-matched symptom-free controls; no differences were found for synovitis, suggesting normalisation. Physical functioning, measured with HAQ, remained similar in patients with persistent symptoms (0.56 to 0.51; $p=0.61$), whereas HAQ normalised in patients with resolution of symptoms (mean score of 0.15 at 2 years, $p=0.053$ compared to baseline).

Conclusions: Approximately one-third of the patients that presented with CSA and that did not convert to IA had resolution of symptoms, this was paralleled by normalisation of MRI-detected inflammation as well as physical functioning. This is the first study showing that subclinical inflammation in individuals at risk for developing RA can resolve spontaneously. The data suggest that subclinical inflammation was causally related to joint pain.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2018-eular.3436

OP0154

PROSPECTIVE OBSERVATIONAL STUDY TO EVALUATE THE USE OF MUSCULOSKELETAL ULTRASONOGRAPHY TO IMPROVE RHEUMATOID ARTHRITIS MANAGEMENT: INTERIM ANALYSIS OF THE ECHO STUDY

M. Stein^{1,2}, E. Rampakakis³, J.S. Sampalis^{2,3}. ¹McGill University, ²Canadian Rheumatology Ultrasonography Society (CRUS), Montreal; ³JSS Medical Research, St. Laurent, Canada

Background: Musculoskeletal Ultrasound (MSUS) has been shown to be superior to clinical examination in the detection of synovitis in patients with Rheumatoid Arthritis (RA), and can be used to improve diagnostic accuracy and potentially monitor disease changes in order to make treatment decisions aimed at optimising patient care. Since the creation of the Canadian Rheumatology Ultrasonography Society (CRUS) in 2010, an increasing number of rheumatologists has been trained in the use of MSUS.

Objectives: The overall study objective is to compare the effectiveness of MSUS to Routine Care (RC) as a disease management tool in patients with moderate-to-severe RA for whom a change in treatment is indicated. In addition, the predictive power of MSUS assessments has been assessed here.

Methods: 'Echo' is a prospective two-cohort, quasi-experimental study of patients diagnosed with active moderate-to-severe RA managed either with MSUS (within CRUS) or as per RC. To be eligible for the study patients must require a change in treatment as per the judgment of the treating physician. Patients are followed for 1 year with assessments at baseline, 3, 6, 9, and 12 months. Key outcome measures of interest include CDAI LDA/Remission, DAS-28 LDA/Remission, patient satisfaction (TSQM) and patient perception of participation in disease management (PAM-13).

Results: A total of 383 patients (71.5% female) with a mean (SD) age of 58.7 (11.7) years and disease duration of 7.0 (10.0) years were enrolled, without any significant differences between treatment groups.

At baseline, a greater proportion of patients in the MSUS group were treated with a biologic DMARD (bDMARD; 50.3% vs 35.8%, $p=0.004$); patients in the RC group were more likely to be treated with a non-biologic DMARD (nbDMARD; 84.2% vs 91.5%, $p=0.027$). Over time, a comparable proportion of patients in the two groups started/switched a bDMARD (21.6% vs 15.6%, $p=0.126$) or added/switched a nbDMARD (18.7% vs 23.6%, $p=0.248$). The overall number of treatment modifications was also similar between groups (3.0 vs 2.7, $p=0.236$).

Upon adjusting for age, gender, previous bDMARD treatment, and baseline parameter level, no differences between the two treatment groups with respect to CDAI LDA/Remission, DAS-28 LDA/Remission, and TSQM score were observed during follow-up. However, the PAM-13 score was significantly higher in the MSUS group (69.6 vs 64.2, $p=0.02$).

In the MSUS group, higher total US erosion score at baseline was associated with a lower rate of CDAI LDA at 12 months (OR=0.86; $p=0.047$); higher total PD synovitis score at baseline was associated with a lower rate of CDAI LDA at 6 months (OR=0.90; $p=0.010$), and; higher total synovitis GREY scale at baseline was associated with lower rates of DAS28 LDA (OR=0.93; $p=0.026$) and DAS28 remission (OR=0.94; $p=0.061$) at 6 months.

Conclusions: MSUS assessments can be useful predictors of future disease remission in patients with RA. MSUS may be associated with increased patient perception of participation in disease management and patient activation.

Acknowledgements: The Sponsors of this investigator-initiated study are the Canadian Rheumatology Ultrasonography Society (CRUS) and JSS Medical Research (in-kind support). The study is supported by an unrestricted grant by AbbVie.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2018-eular.3563

OP0155

ULTRASOUND AS AN OUTCOME MEASUREMENT TOOL FOR OPTIMISED MONITORING OF GOUT. VALIDATION OF THE OMERACT ULTRASOUND DEFINITIONS OF GOUT ELEMENTARY LESIONS

S.N. Christiansen, M. Østergaard, O. Slot, L. Terslev. *Center for Rheumatology and Spine Diseases, Copenhagen Center for Arthritis Research, Rigshospitalet, Glostrup, Glostrup, Denmark*

Objectives: To evaluate ultrasound (US) as an outcome measurement instrument for monitoring gout patients during urate lowering therapy using the OMERACT US Working Group's 2015 definitions of US elementary lesions in gout.

Methods: US examination (28 joints, 26 tendons) were performed in patients with microscopically verified gout who either initiated or increased urate lowering therapy. Joints and tendons were evaluated for the four OMERACT elementary lesions of gout (Double contour, Tophus, Aggregates and Erosions). Furthermore, subcutaneous (SC) oedema was registered and synovitis was graded by grey scale (GS) and colour Doppler (CD) (both graded 0-3). A sum score was