targeted-synthetic DMARDs. The use of any conventional synthetic DMARD was associated with lower HAQ-DI indices ($p: 0.0243$), while the use of any biologic DMARD or targeted-synthetic DMARD was related to greater functional impairment ($p: 0.0018$). By evaluating separately, abatacept ($p: 0.0046$), rituximab ($p: 0.0001$) and tocilizumab ($p: 0.0441$) were associated with higher levels of HAQ-DI. The results are summarised in table 1.

Abstract AB0200 – Table 1. Health assessment questionnaire-disability index score position and dispersion measurements and result of comparison between groups

Variable	Category	N	Mean	SD	Median	p-value
Rheumatoid factor	High	612	0.98	0.80	0.88	0.0244 ^c
	Low	247	0.89	0.73	0.75	
	Negative	235	0.82	0.74	0.75	
ACPs	High	262	0.96	0.77	0.88	0.3927 ^c
	Low	86	0.89	0.84	0.63	
	Negative	111	0.88	0.75	0.68	
Bone erosion	No	493	0.77	0.72	0.63	<0.0001 ^a
	Yes	599	1.06	0.80	1.00	
Methotrexate	No	373	0.96	0.80	0.88	0.4028 ^b
	Yes	739	0.91	0.76	0.75	
Lefunomide	No	735	0.93	0.77	0.88	0.8936 ^b
	Yes	377	0.93	0.78	0.88	
Hydroxychloroquine	No	993	0.94	0.78	0.88	0.2696 ^b
	Yes	119	0.85	0.75	0.75	
Sulfasalazine	No	1,058	0.93	0.78	0.88	0.6727 ^b
	Yes	54	0.87	0.75	0.81	
Conventional synthetic DMARDs	No	115	1.12	0.89	1.00	0.0243 ^b
	Yes	997	0.91	0.76	0.75	
Adalimumab	No	1,059	0.92	0.77	0.88	0.7728 ^b
	Yes	53	1.00	0.86	0.75	
Infliximab	No	1,064	0.93	0.77	0.88	0.8659 ^b
	Yes	48	0.94	0.78	0.81	
Etanercept	No	1,046	0.94	0.77	0.88	0.1317 ^b
	Yes	66	0.79	0.75	0.69	
Certolizumab pegol	No	1,095	0.93	0.78	0.88	0.3074 ^b
	Yes	17	0.72	0.67	0.50	
Golimumab	No	1,075	0.93	0.78	0.88	0.8372 ^b
	Yes	37	0.85	0.70	1.00	
Tocilizumab	No	1,058	0.92	0.78	0.75	0.0441 ^b
	Yes	54	1.08	0.66	1.00	
Abatacept	No	1,040	0.91	0.77	0.75	0.0046 ^b
	Yes	72	1.16	0.78	1.00	
Rituximab	No	1,063	0.91	0.77	0.75	0.0001 ^b
	Yes	49	1.34	0.77	1.00	
Tofacitinib	No	1,103	0.93	0.77	0.88	0.9211 ^b
	Yes	9	0.93	0.76	0.75	

Conclusions: In our prospective cohort, patients with high levels of RF, bone erosion, in use of any biologic DMARD or targeted-synthetic DMARD, abatacept, rituximab or tocilizumab had worse functional capacity indexes. When compared to non-use, the use of any conventional DMARD was associated with better rates of HAQ-DI.

REFERENCES:

- [1] Drossaers-Bakker KW, de Buck M, van Zeben D, et al. Long-term course and outcome of functional capacity in rheumatoid arthritis: The effect of disease activity and radiologic damage over time. *Arthritis Rheum* 1999;42:1854.
- [2] Scott DL, Pugnier K, Kaarela K, et al. The links between joint damage and disability in rheumatoid arthritis. *Rheumatology (Oxford)* 2000;39:122.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2018-eular.3556

AB0201

ASSOCIATION BETWEEN BASELINE CALPROTECTIN SERUM LEVELS AND RESPONSE TO BIOLOGICAL THERAPY IN PATIENTS WITH RHEUMATOID ARTHRITIS

A. Martínez-Feito¹, C. Plasencia¹, A. Jochems¹, V. Navarro-Compan¹, D. Pascual-Salcedo¹, B. Hernández-Breijo¹, P. Nozal², D. Peiteado¹, A. Villalba¹, A. Mezcueta², A. Balsa¹. ¹Immuno-Rheumatology research group; ²Immunology, University Hospital La Paz, Madrid, Spain

Background: Calprotectin (CLP) is an important proinflammatory factor of innate immunity released from activated granulocytes and macrophages during inflammation, which has also been identified in synovial fluid. It is a potential pro-inflammatory biomarker reflecting joint damage. CLP levels in synovial fluid are well correlated with levels in plasma, which allows measuring it easily in patients with rheumatoid arthritis (RA). However, the additional value of CLP over other biomarkers is unclear

Objectives: To study the association of baseline CLP serum levels with the clinical response and serum drug levels in patients under biological therapy at 6 and 12 months after starting treatment.

Methods: Prospective observational study including 109 patients with RA who started biological treatment (Infliximab(Ix); Adalimumab(Ada); Etanercept(Etn); Certolizumab(Ctz); Golimumab(Glm); Tocilizumab(Tcz) and Rituximab(Rtx) in a tertiary hospital since 1999 to 2016. Serum CLP levels were measured by ELISA

with the commercial kit CALPROLAB (Lysaker, Norway). Levels of biological drug and ADA were measured by capture and bridging ELISA respectively, except Certolizumab which was measured by Sanquin Diagnostic Service. Clinical response was assessed by DAS28. Serological and clinical parameters were evaluated at baseline, 6 and 12 months. For analysis, linear regression models adjusted for confounders (age, sex, BMI, DMARDs, smoking status) and with standardised beta as estimate were employed.

Results: Out of 109 studied patients, 22% patients received Ix, 6% Ada, 16% Etn, 18% Ctz, 5% Glm, 25% Tcz and 9% Rtx. Most patients were woman (84%), with a median (IQR) age of 62 (52.3–73.1) and disease duration of 9.5 (4.8–15.3). At baseline, CLP levels directly correlated well with CRP and DAS28 ($r=0.411$; $p<0.001$ and $r=0.308$; $p<0.001$). On the other hand, an inverse correlation between baseline CLP levels and the presence of drug at 6 ($\beta=-0.325$; $p=0.01$) and 12 months ($\beta=-0.397$; $p<0.001$) was also found. Moreover, in the adjusted analyses a trend for this correlation at 6 months ($\beta=-0.199$; $p=0.07$) and a significant correlation at 12 months ($\beta=-0.381$; $p<0.001$) was observed. Additionally, in the non-adjusted models, baseline CLP levels significantly directly correlated with the clinical response at 6 months ($\beta=0.194$; $p=0.04$) while the correlation at 12 months was not significant ($\beta=0.186$; $p=0.06$). However, after adjusting for confounders, no significant correlation was found at any study point ($\beta=0.213$; $p=0.06$ at 6 months and $\beta=0.206$; $p=0.07$ at 12 months).

Conclusions: In patients with RA initiating biological therapy, baseline CLP concentrations are inversely correlated with the circulating drug levels along the treatment. However, the additional value of CLP as a predictor of clinical response remains unclear.

Acknowledgements: This work has been supported by a SORCOM grant at 2016

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2018-eular.5537

AB0202

GLUCOCORTICOIDS IN THE INITIAL TREAT-TO-TARGET STRATEGY OF EARLY RHEUMATOID ARTHRITIS

A.L. Fedele, L. Petricca, B. Toluoso, S. Alivernini, M.R. Gigante, S.L. Bosello, C. Di Mario, G. Ferraccioli, E. Gremese. *UOC of Rheumatology, Fondazione Policlinico Universitario A. Gemelli - Università Cattolica del Sacro Cuore, Rome, Italy*

Background: As stated in the 2013 update of RA recommendations, glucocorticoids (GCs) should be used as bridging therapy for up to 6 months, ideally tapering them at earlier time points.¹

Objectives: To evaluate whether initial combination therapy with GCs and disease modifying anti-rheumatic drugs (DMARDs) influences clinical and radiological outcome in the real-life practice of a cohort of early rheumatoid arthritis (ERA) patients.

Methods: A total of 367 ERA patients with less than 12 months of disease duration were enrolled in the study. ERA patients fulfilled the 2010 ACR criteria for RA and were followed according to the treat-to-target strategy. The mean follow-up (FU) was 38.2±32.8 months. At baseline, and every three months, the ACR/EULAR core data set variables were recorded. At baseline and every year, hand and foot radiographs were examined according to modified Total Sharp score (mTSS). At each visit, clinical improvement and remission were evaluated according to EULAR criteria. The achievement of CDC (28-joint Disease Activity Score using C reactive protein <2.6, Health Assessment Questionnaire <0.5 and change from baseline in mTSS ≤0.5) was assessed every year of follow-up.

Results: At baseline 291 (71.9%) ERA patients started GCs at a dosage of 0.2 mg/Kg, gradually tapered and withdrawn as rapidly as clinically feasible. As expected, these patients presented higher values of acute phase reactants ($p=0.001$), and higher levels of disease activity scores ($p<0.001$) and disability index ($p=0.001$), compared to the 76 subjects (18.8%) who had not been prescribed GCs. Patients not treated with GCs were in higher percentages anti-citrullinated peptide antibody (ACPA) (75.0%) and IgM-rheumatoid factor (RF) (64.5%) positive, compared to subjects taking GCs (62.9%, $p=0.05$; 49.8%, $p=0.02$, respectively). There were no differences regarding age, disease duration, BMI, smoking habit and presence of erosions at onset. The mean duration of GC treatment was 7.5±7.9 months and the mean dosage during FU was <5 mg prednisone per day. During the FU, in the two groups a similar percentage of patients started a combination therapy with biological (b)-DMARDs and no differences were observed regarding radiographic progression.

A higher rate of remission defined according to DAS values, 2011 ACR/EULAR criteria, and CDC criteria, was registered in patients not treated with GCs compared to subjects who required corticosteroid therapy. ERA patients that didn't manage to stop GCs at the sixth month (38.8%) had a higher BMI ($p=0.04$) and a lesser chance of achieving remission defined according to DAS values, 2011 ACR/EULAR criteria, and CDC criteria during follow-up. Moreover, a higher percentage of them required a combination therapy with bDMARDs during FU ($p<0.0001$).

Conclusions: In our cohort ERA patients initially treated with GCs had higher disease activity scores at onset compared to subjects without GCs. The lesser chance of achieving remission and the higher rate of bDMARD therapy in ERA