patients not able to stop GCs, reflect a more aggressive disease, refractory to conventional drugs.

REFERENCE:

- [1] Smolen JS, et al. *Ann Rheum Dis* 2014;73:492–509.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2018-eular.7507

AB0203

CLINICAL PHENOTYPE AND ULTRASOUND CHARACTERISTICS OF RHEUMATOID ARTHRITIS FLARE AFTER DISCONTINUATION OF CONVENTIONAL SYNTHETIC DMARDS

A. Manzo, S. Bugatti, F. Benaglio, G. Sakellariou, R. Caporali, C. Montecucco. *Arthritis Remission Clinic, Division of Rheumatology, IRCCS Policlinico San Matteo Foundation/University of Pavia, Pavia, Italy*

Background: Current protocols based on early and intensive treatment with csDMARDs in rheumatoid arthritis (RA) have allowed the achievement of remission in a considerable proportion of the patients and opened the perspective, in selected cases, of a drug-free monitoring scheme. Treatment discontinuation can lead, however, to possible recurrence of joint inflammation and clinical flare. Understanding the dynamics acting upstream these events remains a fundamental research task with direct clinical and patho-biologic implications.

Objectives: To delineate the clinical, serological and ultrasonographic changes associated to a drug-free flare in patients discontinuing csDMARD after achievement of stable remission. Co-primary objective was to compare, through a retrospective analysis in the same patients, these changes with early features of the pathology at onset, before treatment introduction.

Methods: 92 RA patients in stable DAS28 remission following a DAS-steered treatment strategy with MTX were recruited in our Centre and introduced to a drug-free monitoring scheme according to the following inclusion criteria: a) treatment introduced within 12 months from symptoms' onset, b) at least 24 months of continuative treatment, c) DAS28 <2.6 for at least 6 months in the absence of glucocorticoids. After discontinuation, all patients were follow-up at three months intervals across 24 months through complete clinical, ultrasonographic (power Doppler ultrasound –PDUS- in hands-feet and tendons) and serological analyses. Treatment was re-introduced upon occurrence of moderate disease activity (DAS28 ≥3.2) in a single occasion.

Results: A total drug-free follow-up of 1398 person-months was analysed with a median (IQR) of 15^{6–24} months. Thirty-eight patients (27/38 in ACR/EULAR Boolean remission, 16/38 with PD score=0 at withdrawal visit) required treatment re-introduction after a median (IQR) time from discontinuation of 6^{3–9} months (range 3–18). DAS28 variations at re-treatment showed a mean (SD) increase of 2.26 (1.03), reflecting significant differences in all DAS components (p<0.001 for ESR, tender joint count, swollen joint count and GH). Clinical activity in flaring subjects was paralleled by average changes in synovial US, with increased PD scores in hands joints (median [IQR]: 3.5 [0.5–7] vs 1 [0–2], p<0.001), feet (1 [0–4] vs 0 [0–0.5], p=0.002) and tendons (0 [0–2] vs 0 [0–0], p=0.002), determining *ex-novo* PD positivity in 85.7% of PD negative patients (p<0.001). Despite stringent remission achieved at the time of discontinuation, no significant differences were observed between disease onset and drug-free flare in DAS28 (p=0.26), patient global assessment of disease activity (p=0.53) and synovial US scores (p=0.61 for grey scale, p=0.31 for PD) with recurrence of similar patterns of clinical joint involvement.

Conclusions: Drug-free clinical flare can occur over a wide temporal window, in the absence of detectable signs of inflammation at the time of treatment discontinuation. It can associate with *ex-novo* recurrence of US pathologic changes at joint and tendon level, reproducing some of the quantitative/qualitative features of disease onset.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2018-eular.7443

AB0204

RADIOGRAPHIC PROGRESSION OF LARGE JOINT DAMAGE IN PATIENTS WITH RHEUMATOID ARTHRITIS TREATED WITH BIOLOGICAL DISEASE-MODIFYING ANTIRHEUMATIC DRUGS (bDMARDs) AND ITS PREDICTIVE FACTORS: RESULTS OF 3 TO 4 YEARS FOLLOW-UP

A. Nakajima, K. Terayama, M. Sonobe, H. Takahashi, Y. Akatsu, S. Taniguchi, M. Yamada, K. Nakagawa. *Toho University Sakura Medical Center, Sakura, Japan*

Background: Although damage to large joints such as the shoulder, elbow, hip, knee, and ankle has a substantially larger impact on functional ability than

damage to the small joints of the hand and foot,^{1,2} large joints are not routinely monitored for progressive damage in patients with rheumatoid arthritis (RA). Furthermore, a little information is available regarding long-term follow-up results of radiographic progression of damage (RPD) to the large joints during treatment with biological disease-modifying antirheumatic drugs (bDMARDs).

Objectives: We investigated ratios of RPD to the large joints in RA patients treated with bDMARDs for 3 to 4 years and analysed association between RPD and patient backgrounds or Larsen grades of individual large joint.

Methods: Sixty-eight patients (naïve: 42, switch: 26) receiving bDMARDs (IFX: 5, ETN: 9, ADA: 7, GLM: 3, CZP: 5, TCZ: 28, ABT: 11) for 3 to 4 years or achieving bDMARDs-free status were included in this study. The mean age and disease duration at the start of bDMARDs was 62.7 year-old and 10.7 years, respectively, and baseline DAS28-ESR and HAQ was 4.61 and 1.014, respectively. A total of 311 joints including shoulder, elbow, hip, knee, and ankle were evaluated whether there was RPD during the observation period by comparing radiographs before and after the treatment.

Results: RPD was found in 22 patients (31.4%) and 32 joints (10.0%), and it occurred during the former half period in 17 joints and during the latter half period in 15 joints. Joints with Larsen grade (LG) III or more had significantly higher ratios of RPD than those with LG II or less (p<0.01). An ROC analysis was performed to calculate the cut-off value for the progressive damage, which was 2.5 (sensitivity: 50.0%, specificity: 79.6%), suggesting that exceeding LG III would be a risk factor for the RPD to the large joints. A multivariate logistic regression analysis revealed that stage and baseline HAQ were independent risk factors for RPD (cut-off value: 2.5, odds ratio: 8.864 for stage; cut-off value: 1.4375, odds ratio: 6.316 for baseline HAQ).

Conclusions: The stage and baseline HAQ were associated with RPD to the large joints, and progressive damage is expected to increase when the stage exceeds III and/or functional disability exceeds an HAQ score of 1.5. Progressive damage also increases when LG exceeds III. Treatment with bDMARDs should be started before stage, HAQ, and LG exceed III, 1.5, and III, respectively.

REFERENCES:

- [1] Drossaers-Bakker KW, Kroon HM, Zwiderman AH, Breedveld FC, Hazes JM. Radiographic damage of large joints in long-term rheumatoid arthritis and its relation to function. *Rheumatology* (Oxford) 2000;39:998–1003.
[2] Kuper HH, van Leeuwen MA, van Riel PL, Prevoo ML, Houtman PM, Lolkema WF, van Rijswijk MH. Radiographic damage in large joints in early rheumatoid arthritis: relationship with radiographic damage in hands and feet, disease activity, and physical disability. *Br J Rheumatol* 1997;36:855–60.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2018-eular.3408

AB0205

COMPARISON OF THORACIC HRCT AND SELF-REPORTED QUESTIONNAIRES IN THE ASSESSMENT OF PULMONARY INVOLVEMENT IN RHEUMATOID ARTHRITIS PATIENTS: PRELIMINARY RESULTS

A. Topcu¹, Y. Yalcinkaya², S. Karakurt³, H.H. Mursaloglu⁴, E. Albayrak⁴, N. Coşkun⁵, M. Demir¹, E. Dalkılıç⁵, N. İnanç². ¹Department of Internal Medicine; ²Division of Rheumatology, Department of Internal Medicine; ³Department of Chest Diseases; ⁴Marmara University Faculty of Medicine, Istanbul; ⁵Division of Rheumatology, Department of Internal Medicine, Uludağ University Faculty of Medicine, Bursa, Turkey

Background: Pulmonary involvement in rheumatoid arthritis (RA) is one of the extra-articular manifestations affecting morbidity and mortality during the course of the disease. Pulmonary function tests (PFTs) and thoracic high-resolution computerised tomography (HRCT) are the standard of care in the assessment of pulmonary involvement in RA. In this study, we aimed to compare the findings between self-reported questionnaires and HRCT to detect pulmonary abnormalities in RA patients.

Methods: Forty-two RA patients fulfilling ACR/EULAR classification criteria (2010) who had thoracic HRCT within 6 months of any symptom and/or any pathology on radiography of chest were included in the study. The patients were also assessed by modified Borg Scale, SF-36 Quality of Life Scale and Leicester Cough Questionnaire for the evaluation of respiratory symptoms.

Results: Demographics and clinical characteristics were summarised in table 1. Warrick score, assessing the severity and extent of alveolitis and fibrosis on thoracic HRCT, was evaluated in 15 patients with ILD (score range:4–28). DLCO values were lower in patients with Warrick score ≥1 (73±22% vs. 88±12%, p=0.019) while FVC were not found to be different. The findings of HRCT and self-reported questionnaires were summarised in table 2. Any relationship between self-reported questionnaires and Warrick scores was not detected. Presence of any parenchymal lesions was found to be associated with SF-36 total score (p=0.048). DLCO levels were found to be negatively correlated with SF-36 total scores (r=-0.470, p=0.006).