parameters (DAS28, ESR, CRP and HAQ) was observed ($p<0.05$). Likewise, the % of hypertensive and dyslipidemic patients increased during follow up ($p<0.05$), while the levels of cholesterol, triglycerides, glucose and diabetic patients remained similar. The presence of CV event was related to high levels of ESR, male sex and hypertension, ($p<0.05$).

Conclusions:

- The incidence of cardiovascular event in our recent onset RA cohort resembles that previously described in the literature, with the majority occurring during the first 5 years of follow-up.
- The development of hypertension and dyslipidemia occurs mainly during the first 5 years of follow-up.
- The presence of a cardiovascular event was related to male sex, high blood pressure and high ESR levels.

REFERENCES:

- [1] McInnes IB, Schett G. The Pathogenesis of Rheumatoid Arthritis. *N Engl J Med* 2011;365(23):2205–19.
- [2] Mielants H, Van den Bosch F. Extra-articular manifestations. *Clin Exp Rheumatology* 2009;27(Suppl 55):S56–S61.
- [3] Nurmohamed MT. Cardiovascular risk in rheumatoid arthritis. *Autoimmunity Reviews* 2009;8:663–7.
- [4] del Rincón ID, Williams K, Stern MP, Freeman GL, Escalante A. High incidence of cardiovascular events in a rheumatoid arthritis cohort not explained by traditional cardiac risk factors. *Arthritis Rheum* 2001;44(12):2737–45.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2018-eular.3140

AB0395

CORRELATION BETWEEN CARDIOVASCULAR RISK CALCULATORS IN A COHORT OF PATIENTS WITH RHEUMATOID ARTHRITIS

R. Castellanos-Moreira, S.C. Rodríguez-García, S. Mandelkova, V. Ruiz-Esqueda, B. Frade-Sosa, A.B. Azuaga-Piñango, S.D.C. Rodríguez-García, N. Sapena, C. Gonzalez-Delaurens, O. Camacho, J. Ramírez, M.V. Hernández, A. Cuervo, J. A. Gómez-Puerta, J.D. Cañete, R. Sanmarti. *Rheumatology, Hospital Clinic de Barcelona, Barcelona, Spain*

Background: EULAR recommends for patients with rheumatoid arthritis (RA) a cardiovascular (CV) risk assessment at least once every five years. Several risk calculators are available but most of them are based on data from general population, so they may underestimate the risk of RA patients. To obtain a more accurate estimation they should be adapted for RA by a 1.5 multiplication factor, unless the instrument includes RA as a variable. To date, there are no studies analysing the concordance and correlation between different CV risk calculators in spanish population.

Objectives: To analyse concordance and correlation of CV risk calculated using different tools and to determine the proportion of CV risk factors in a cohort of patients with RA.

Methods: We performed a cross-sectional study including patients with RA according to ACR/EULAR 2010 criteria, treated in a tertiary hospital. Individuals with previous CV events, diabetes and chronic kidney disease were excluded. Ten-year CV risk was obtained with the following calculators: Framingham-body mass index (FRS-BMI), Framingham-lipids (FRS-L), SCORE CV risk calculator (SCORE) and the Expanded CV Risk Prediction Score for RA (ERS-RA). A 1.5 multiplying factor was used to adapt the results for RA patients in those which didn't include it in their models. CV risk was categorised as low or high using 10% as a cut-off for FRS-BMI and FRS-L and 5% for SCORE y ERS-RA. Correlation was evaluated using Spearman correlation coefficient (Rho) and concordance with weighted kappa.

Results: We included 88 patients (18 male), 87.5% Caucasian, mean age 56.9 $\pm$ 10.6 years, mean disease duration 5.8 $\pm$ 3.3 years. 67% and 76% had positive RF and ACPA respectively and 15% presented extra-articular manifestations (rheumatoid nodules, interstitial lung disease, pleuropericarditis). Regarding CV risk factors: 14.8% were active smokers, 60.2% were overweight or obese, 25% had hypertension, 47.8% had dyslipidemia.

The median CV risk predicted at 10 years was 15% (1.5–70.20) for FRS-IMC, 11.8% (1.2–54.0) for FRS-L, 1.5% (1.1–16.5) for SCORE and 6% (0.6–27.8) for ERS-AR. 64 patients were categorised as low risk and 22 as high risk for ERS-AR. No significant differences were observed in the means of risk obtained by the different calculators or in the risk categories according to ERS-RA when stratified by disease duration (more or less than 5 years).

For the correlation between ERS-RA and FRS-BMI, FRS-L and SCORE, Spearman's Rho coefficients were 0.84, 0.79 and 0.80. ($p<0.005$).

For the concordance between ERS-RA vs FRS-BMI a weighted $\kappa=0.29$ (CI₉₅: 0.15–0.42) was obtained. For ERS-AR vs FRS-L, $\kappa=0.34$ (CI₉₅: 0.19–0.50) and for ERS-AR vs SCORE, $\kappa=0.70$ (CI₉₅: 0.52–0.87).

Conclusions: A strong correlation was observed between the CV risk calculators evaluated and good agreement only between ERS-AR and SCORE. Additionally, overweight, obesity and dyslipidemia were the most prevalent comorbidities.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2018-eular.7044

AB0396

RELATIONSHIP BETWEEN PATIENT GLOBAL ASSESSMENT AND PAIN ASSESSMENT IN THE DISEASE ACTIVITY INDEXES IN RHEUMATOID ARTHRITIS, AND ITS CORRESPONDENCE WITH SONOGRAPHIC ALTERATIONS

R. Menor Almagro, R.M. Martínez, E.M. Rubio, M.M. Fernández. *Rheumatology, Hospital Universitario Virgen del Rocío, Sevilla, Spain*

Background: Disease activity indexes in rheumatoid arthritis (RA) provide information of different parameters in a single value. It is very useful in the evaluation of the disease activity and its progression. The subjective component of these indexes can modify the index final value.

Objectives: The aim of our study was to describe the relation between the patient global assessment (PGA) and pain assessment (PA) with the sonographic alterations in AR.

Methods: Transversal study describing the ultrasonographic changes and disease activity indexes in AR. We included patients with RA according to ACR/EULAR classification criteria. They were sent by their usual clinician, in random selection. They were reviewed in the same day by a rheumatologist and blind sonographer. PGA, PA, HAQ, tender and swollen joint count, CDAI, SDAI and DAS28 were evaluated. The ultrasound were realized by grey scale and doppler in 12 joints (wrists, second to fifth MCF and fifth bilateral MTF).

Results: A total of 48 patients had PGA <50 and 37 PGA $\geq$ 50. The mean age was 53.2 $\pm$ 11.7 for group 1 and 56.5 $\pm$ 10.2 for group 2, women 64.6% and 83.8% respectively. The positive rheumatoid factor resulted in 72.9% vs 73%. The evolution of the disease in months presented a median of 96 (47.2–132.7) vs 108 [48–138], $p=0.564$. The Body Mass Index in the groups was 26.6 (23.5–29.7) and 26.0 (22.8–30.0), $p=0.647$, and the tobacco consumption of 11 patients group 1 and 12 in group 2 ($p=0.397$). No significant difference was found in the treatment at the time of the evaluation with methotrexate, median of 10 (0–15) vs 7.5 (0–15), $p=0.350$, or corticosteroids 1.4 (0–2.3) vs 2.5 (0–4.5), $p=0.05$. 23 patients in the group 1 and 19 in the group 2 were in biological therapy. The results referring to the patient's assessment of their disease, acute-phase reactants, clinical examination, ultrasound and disease activity indexes are presented in table 1.

Abstract AB0396 – Table 1

	Group 1 (PGA<50)	Group 2 (PGA $\geq$ 50)	P value
ESR: median (p25-p75)	9 (4,2-17)	8 (4,5-16)	0578
CRP: median (p25-p75)	2,4 (1,02-4,7)	5 (1-5,7)	0996
DAS 28 (CRP)	2,1 $\pm$ 0,97	3,37 $\pm$ 1,19	0000
DAS 28 (ESR)	2,6 $\pm$ 0,72	3,25 $\pm$ 0,92	0000
Pain assessment, 100 mm VAS: median (p25-p75)	20 (0-40)	70 (50-80)	0000
HAQ: average $\pm$ DE	0,8 $\pm$ 0,14	2,27 $\pm$ 0,18	0000
Tender joint count: median (p25-p75)	0 (0-2)	2 (0-3)	0001
Swollen joint count: median (p25-p75)	0 (0-1)	1 (0-3)	0090
CDAI: median (p25-75)	5 (1-8,8)	13 (9,4-17,4)	0000
SDAI: average $\pm$ DE	8,69 $\pm$ 5,82	17,67 $\pm$ 6,49	0000
Synovitis in grey scale: median (p25-p75)	1 (0-3)	2 (0-4)	0159
Synovitis in doppler: median (p25-p75)	0 (0-1)	1 (0-2,7)	0231

Conclusions: Disease activity indexes in AR have been established as a very useful measure, both in the homogenization of the results obtained in the research and in the daily clinical practice. In our study, we observed the modification in the values of disease activity indexes due to PGA and PA, even those in which its contribution is less decisive than other parameters, as it happens in the DAS 28.

Those variables no related in their value with the patient assessment of his disease status did not show statistical significance between the groups. The patients' valuation about activity of their AR and the modification of the indexes in which it participates in our study were not concordant with the echographic alterations found in grey scale and doppler echo.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2018-eular.7080