Methods: Transversal descriptive study. We included 13 patients diagnosed with SLE according to the criteria of the American College of Rheumatology (ACR) with positivity for antinuclear antibodies (ANA) and anti-DNA, 34 first-degree relatives and 50 healthy controls between the months of May 2016 and March 2017. None of the subjects evaluated received treatment with rituximab or belimumab. We analysed B-cells subsets (negative double, naïve B-cells, unswitched memory B-cells, switched memory B-cells) in all the subjects included in our study. The 95% confidence intervals were obtained for both the means and the percentage difference. The level of statistical significance was established at $p < 0.05$. The data was analysed with IBM SPSS software Statistics 19 and EPIDAT 4.1.

Results: 47 subjects were analysed between relatives and patients, of which 33 (70.20%) were women. 13 subjects (27.70%) were diagnosed with lupus. 100% of those diagnosed with lupus were women. The mean (X) and confidence intervals (95% CI) for the different subgroups (healthy subjects, subjects diagnosed with SLE, relatives of the first degree) is shown in table 1. In none of the subpopulations analysed in patients diagnosed with SLE in front of relatives of 1 st grade it has reached statistical significance. When analysing B cells subsets of the three groups of subjects, we did find statistically significant differences between unswitched memory B cells of healthy subjects and 1 st degree relatives (being lower in the group of healthy subjects) This finding has not been described in any previous study, although it should be noted that the sample before us is small.

Abstract OP0178 – Table 1

B-CELLS SUBSETS		HEALTHY	SLE	RELATIVES
Naïve (%)	Mean (CI95%)	70,4 (95% CI: 68,2 to 72,7)	69,3 (95% CI: 62,2 to 76,3)	65,8 (95% CI: 60,4 to 71,2)
Doubles Negative (DN) (%)	Mean (CI95%)	4,1 (95% CI: 3,5 to 4,6)	6,2 (95% CI: 4,8 to 7,6)	2,9 (95% CI: 1,9 to 4,0)
Unswitched (%)	Mean (CI95%)	9,6 (95% CI: 8,2 to 11,1)	13,0 (95% CI: 7,4 to 18,7)	17,2 (95% CI: 13,4 to 21,1)
Switched (%)	Mean (CI95%)	11,4 (CI95% =10,1-12,7)	11,5 (95% CI: 6,3 to 16,7)	14,0 (95% CI: 11,2 to 16,8)

Conclusions: There are quantitative differences between unswitched memory B-cells of healthy subjects and relatives of 1 st grade of SLE. More studies with a larger sample size are necessary to see the behaviour of the rest of B-type lymphocyte subpopulations

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2018-eular.7262

OP0179

MOLECULAR DOCUMENTATION OF THE CLONAL EVOLUTION OF A DIFFUSE LARGE B-CELL LYMPHOMA OUT OF CLONALLY EXPANDED RHEUMATOID FACTOR-EXPRESSING B CELLS IN A SJÖGREN'S SYNDROME PATIENT

R.J. Bende¹, L.M. Slot¹, T.A. Wormhoudt¹, M.J. Kwakkenbos², A.H. van Kampen³, A. Jongejan³, J.E. Guikema¹, C.J. van Noesel¹. ¹Pathology, Academic Medical Center; ²AIMM Therapeutics, AIMM Therapeutics; ³Clinical Epidemiology, Biostatistics and Bioinformatics, Academic Medical Center, Amsterdam, Netherlands

Background: Sjögren's syndrome (SS) is the autoimmune disease with the highest risk of lymphoma development. SS patients develop most frequently MALT lymphoma and to a lesser extent diffuse large B-cell lymphoma (DLBCL). Previously, we have shown that at least 40% of salivary gland MALT lymphomas express groups of near identical (also called stereotypic) B-cell receptors, which display *in vitro* mono-specific rheumatoid factor (RF) reactivity¹. Recently, we have analysed the B-cell immunoglobulin heavy variable (IGHV) repertoire in 4 SS labial salivary glands. In 2 out of 4 salivary glands only one minor stereotypic RF B-cell clone was detected. Interestingly, in one salivary gland of patient SG22 a highly expanded stereotypic RF-expressing B-cell clone was present, which was also detected in peripheral blood. Twenty six months later, a clonally-related DLBCL was diagnosed².

Objectives: To study the molecular alterations present in the RF-expressing B cell clones of salivary gland, peripheral blood and in the DLBCL of SS patient SG22.

Methods: From peripheral blood 4 RF-expressing immortalised B-cell clones with identical IGHV rearrangements were isolated. These 4 RF-clones were analysed by whole exome sequencing and the identified non-synonymous exome mutations were traced in the salivary gland and in the DLBCL, using targeted next generation sequencing.

Results: In total we identified 56 exome mutations in the 4 RF B-cell clones. Twelve non-synonymous mutations were shared between all 4 RF-clones, of which one was a missense mutation in *CARD11*, a well-known oncogenic mutation of DLBCL. In the salivary gland all shared 12 non-synonymous mutations were detected, whereas in the DLBCL only 4 of these mutations were detected.

Remarkably, the *CARD11* mutation was not detected in the DLBCL. The exome mutations were analysed for typical characteristics of induction by the B-cell specific mutator AID. This revealed that the large majority of mutations (80%) were (i) not located within 1,5 kB of the transcription initiation site of the genes, (ii) did not concern C/G nucleotides in AID hotspots (WRC/GYW) and (iii) were not in genes expressed by B cells. This lack of evidence for AID involvement, suggests that the exome mutations of the RF-clones were due to replication errors, obtained during their clonal expansion. Interestingly, as compared to the RF-clones of peripheral blood, the DLBCL accumulated 44 extra somatic mutations together in IGHV and IGKV. As expected, these new IGHV/IGKV mutations showed the characteristics of AID involvement, indicating that in an ancestor RF-clone, AID activity was induced.

Conclusions: We have documented the clonal outgrowth and diversification of a RF-specific B-cell clone in a SS patient and the evolution of this clone into a DLBCL. Our analysis are indicative for AID activity in the lymphoma cells, represented by accumulation of IGHV/IGKV mutations. However, we have no evidence that AID is instrumental in the accumulation of the non-IG mutations in the RF-clones from peripheral blood.

REFERENCES:

- [1] Bende, et al. *J. Exp. Med* 2005;201:1229–1241.
- [2] Bende, et al. *Arthritis Rheumatol* 2015;67:1074–1083.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2018-eular.4401

OP0180

TYPE I INTERFERON IS PRODUCED BY NON-HAEMATOPOIETIC TISSUE RESIDENT CELLS BUT NOT PDCs IN PRE-CLINICAL AUTOIMMUNITY AND SLE

A Psarras^{1,2}, A. Alase¹, A. Antanaviciute³, I. Carr³, M. Wittmann¹, G. Tsokos², P. Emery¹, E.M. Vital¹. ¹Leeds Institute of Rheumatic and Musculoskeletal Medicine, University of Leeds, Leeds, UK; ²Beth Israel Deaconess Medical Center, Harvard Medical School, Boston, USA; ³Leeds Institute for Data Analytics, University of Leeds, Leeds, UK

Background: Systemic Lupus Erythematosus (SLE) is characterised by persistently high type I interferon (IFN) activity. Plasmacytoid dendritic cells (pDCs) produce large amounts of IFNs in viral infection, although these immunogenic properties are usually strictly regulated. *In vitro*, pDCs are responsive to nucleic acids and they have therefore been postulated to be the main source of type I IFNs in SLE. However, their function is not fully established in human SLE.

Objectives: To investigate the dysregulated IFN axis in patients with pre-clinical autoimmunity and SLE.

Methods: Patients with SLE who met 2012 ACR/SLICC criteria were recruited. We also recruited healthy controls (HC) and therapy-naïve individuals presenting with ANA and 1–2 clinical symptoms, but not meeting ACR/SLICC criteria, of whom 17% progressed to SLE (At-Risk). IFN activity was evaluated by measuring a score of IFN-responsive genes in the PBMCs using TaqMan. pDCs were immunophenotyped as well as studied *in vitro* for production of proinflammatory cytokines and induction of T cell responses using flow cytometry. pDCs were sorted and sequenced using high-sensitive RNA sequencing. IFN expression was visualised in skin biopsies using *in situ* hybridisation. Keratinocytes were isolated from fresh skin biopsies and cultured *in vitro*; IFN production was measured by qPCR and ELISA.

Results: Most of the SLE and At-Risk patients had increased IFN activity, which correlated with disease activity and clinical features. In contrast, circulating pDCs were decreased in both SLE and At-Risk patients and their numbers did not correlate with any clinical features or IFN status. *In vitro* stimulation revealed that pDCs from SLE and At-Risk patients could not produce IFN- α and TNF- α upon stimulation with TLR9 or TLR7 agonists. In addition, they induced significantly less T cell activation and proliferation compared to HC pDCs. RNA-seq data analysis showed an upregulation of IFN-responsive genes in most of the SLE and At-Risk pDCs but not transcripts of any IFN subtypes. Other upregulated pathways were related to immune regulation and senescence. Phenotypically, SLE pDCs were characterised by upregulation of regulatory receptors and increased telomeric erosion. *In situ* hybridization revealed high IFN expression in the epidermis but not in lymphocyte-infiltrating areas of lesional biopsies from SLE patients. High expression of IFN was also observed in epidermis of At-Risk individuals without any signs of cutaneous inflammation. *In vitro* stimulation of freshly isolated keratinocytes also showed a notable increase in IFN production.

Conclusions: In SLE, non-haematopoietic tissue resident cells are a dominant source of IFN and this is present prior to clinically overt disease. Meanwhile, the professional IFN-producing pDCs have lost their immunogenic properties. These findings suggest an important role for tissue resident cells in autoimmunity and may facilitate novel therapeutic interventions.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2018-eular.5407

OP0181

PATIENT PERSPECTIVES OF PEOPLE WITH PRIMARY SJÖGREN'S SYNDROME: A MULTICENTRE QUALITATIVE EUROPEAN STUDY

J. Unger¹, M. Mattsson^{2,3}, R. Dragoi⁴, C. Boström³, F. Buttgeriet⁵, A. Lackner⁶, T. Witte⁷, B. Raffener⁸, P. Peichl⁹, J. Hermann⁶, C. Dejaco^{6,10}, T. Stamm¹¹. ¹FH JOANNEUM, University of Applied Sciences, Bad Gleichenberg, Austria; ²Sunderby Hospital, Department of Physiotherapy, Luleå; ³Karolinska Institutet, Department of Neurobiology, Care Sciences and Society, Stockholm, Sweden; ⁴Victor Babes University of Medicine and Pharmacy Timisoara, Timisoara, Romania; ⁵Charité University Medicine, Berlin, Germany; ⁶Medical University Graz, Graz, Austria; ⁷Medical University of Hannover, Hannover, Germany; ⁸Hospital of Bolzano, Bolzano, Italy; ⁹Evangelic Hospital of Vienna, Vienna, Austria; ¹⁰Hospital of Bruneck, Bruneck, Italy; ¹¹Medical University Vienna, Vienna, Austria

Background: Primary Sjögren's syndrome (pSS) is one of the most common systemic autoimmune disorders and leads to an impaired health related quality of life. However, treatment mainly focuses on the management of physical manifestations. Little is known about the lived experiences of people with pSS, including the impact on people's life, functioning and their social relationships.

Objectives: To explore the perspectives of people with pSS from different European countries with various cultural backgrounds in order to achieve a broad understanding of concepts that are important and meaningful to people with pSS. This study is a part of a project which aims to evaluate the coverage of the patient perspectives by patient reported outcome measures in pSS, which is funded by the Austrian Association of Rheumatology.

Methods: A multicentre focus group study was performed in five European countries, namely Austria, Germany, Italy, Romania and Sweden. Patients were recruited from the outpatient clinics of the local centres. Focus groups were chaired by a trained moderator and followed an interview guide which included questions about impairments and limitations in body structures, body functions, activities and participation as well as contextual factors and resources, such as coping strategies. Focus groups were audiotaped and transcribed. We conducted a content-analysis of each focus group and subsequently combined the extracted concepts from each country, using the International Classification of Functioning, Disability and Health as a frame of reference.

Results: A total of 12 focus groups was conducted in seven participating centres in five countries. Fifty people (48 women; 96%) with pSS participated in the focus groups (ranging from two to four groups per country). All focus groups had a total duration of 1030 min and resulted in 252 pages of transcript. From qualitative analysis we derived concepts meaningful to people with pSS from all countries, especially those concepts that were linked to a physical dimension. However, we identified differences in the description of these experiences in daily life, for example for pain-concerning sensations or for the impact on social relationships. Furthermore, the attitudes towards the treatment and towards the disease differed between the participants. People with pSS had various coping strategies, such as gaining more knowledge about the disease or utilising non-pharmacological treatment.

Conclusions: This is the first multicentre qualitative European study which investigated the patient perspectives in pSS with a cross-cultural understanding. Clinicians, health professionals and researchers need to know about the perspectives, experiences and needs of people with pSS in order to ensure a comprehensive treatment.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2018-eular.4699

THURSDAY, 14 JUNE 2018

The crystal maze – etiology and management

OP0182

GOUT AND DEMENTIA IN THE ELDERLY: A MEDICARE CLAIMS STUDY

J.A. Singh, J.D. Cleveland. University of Alabama at Birmingham, Birmingham, USA

Background: The pursuit of a link between gout/hyperuricemia and dementia has led to contradictory results. Most observational studies, including population-based studies, showed that hyperuricemia was associated with a higher risk of dementia and less cognitive dysfunction, while a few studies found hyperuricemia to be associated with a lower risk of dementia. Recently, a large French population-based study in the elderly (65 years or older) showed that hyperuricemia was associated with a higher risk of dementia and with MRI changes of ageing in the brain.

Objectives: To assess whether gout in the elderly is associated with a risk of incident dementia.

Methods: We used the 5% Medicare claims data for this observational cohort study. We used multivariable-adjusted Cox proportional hazard models to assess the association of gout with incident dementia, adjusting for potential confounders/covariates including demographics (age, race, gender), comorbidities (Charlson-Romano comorbidity index), and medications commonly used for cardiac diseases (statins, beta-blockers, diuretics, and angiotensin converting enzyme (ACE)-inhibitors) and gout (allopurinol and febuxostat).

Results: In our cohort of 1.23 million Medicare beneficiaries, 65 324 had incident dementia. The crude incidence rates in people without and with gout were 7.36 and 13.58 per 1000 person-years, respectively. In multivariable-adjusted analyses, gout was independently associated with a significantly higher hazard ratio of incident dementia, with a hazard ratio [HR] of 1.17 (95% CI, 1.13, 1.21); sensitivity analyses confirmed the main findings. Compared to age 65 to <75 years, older age groups were associated with 3.3 and 6.3-fold higher hazards of dementia; hazards were also higher for females, Black race or people with higher medical comorbidity.

Subgroup analyses indicated that gout was significantly associated with dementia in patients without key comorbidities (CAD, hyperlipidemia, CVD, diabetes, hypertension) with HR ranging 1.20–1.57, but not in patients with each of these comorbidities, except CAD, with HR 0.97–1.07 (table 1).

Abstract OP0182 – Table 1. Association of gout with incident dementia, in pre-defined subgroups of presence/absence of CAD, hyperlipidemia, CVD, diabetes or hypertension

	Multivariable-adjusted (Model 1)		Multivariable-adjusted (Model 1)	
	HR (95% CI)	P-value	HR (95% CI)	P-value
	No CAD		CAD	
Gout	1.20 (1.15–1.25)	<0.0001	1.07 (1.01–1.14)	0.032
	No Hyperlipidemia		Hyperlipidemia	
Gout	1.33 (1.27–1.40)	<0.0001	1.02 (0.97, 1.07)	0.56
	No CVD		CVD	
Gout	1.23 (1.18–1.28)	<0.0001	0.97 (0.90–1.05)	0.46
	No Diabetes		Diabetes	
Gout	1.24 (1.19–1.29)	<0.0001	1.04 (0.98–1.10)	0.23
	No Hypertension		Hypertension	
Gout	1.57 (1.46–1.68)	<0.0001	1.01 (0.97–1.05)	0.76

Gout*CAD p-value<0.0001; Gout*hyperlipidemia p-value<0.0001; Gout*CVD p-value<0.0001;

Gout*diabetes p-value<0.0001; Gout*hypertension p-value<0.0001;

Conclusions: Gout was independently associated with 17%–20% higher risk of incident dementia in the elderly. Future studies need to understand the pathogenic pathways involved in this increased risk.

Acknowledgements: This material is the result of work supported by research funds from the Division of Rheumatology at the University of Alabama at Birmingham and the resources and use of facilities at the Birmingham VA Medical Centre, Birmingham, Alabama, USA.

Disclosure of Interest: J. Singh Grant/research support from: Takeda, Savient, Consultant for: Savient, Takeda, Regeneron, Merz, Iroko, Bioiberica, Crealta/ Horizon and Allergan pharmaceuticals, WebMD, UBM LLC and the American College of Rheumatology, J. Cleveland: None declared

DOI: 10.1136/annrheumdis-2018-eular.7595

OP0183

ASSOCIATION BETWEEN HFE GENOTYPES AND CLINICAL SEVERITY CHARACTERISTICS OF CALCIUM PYROPHOSPHATE CRYSTAL ARTHRITIS

F. Perez-Ruiz, S.P. Chinchilla, J. Atxotegi. Rheumatology Division of Cruces Hospital and University of the Basque Country, Baracaldo, Spain

Background: Several metabolic disturbances that reduce the activity of pyrophosphatases have been associated with development of calcium pyrophosphate crystal arthritis (CPPCA), but there is scarce data on their influence on clinical manifestations, as such disease-specific variables are not recorded in most available databases.

Objectives: To evaluate factors associated to severity of clinical joint involvement in patients with definite CPPCA.

Methods: Transversal study with prospective recruitment of cases (patients with CPPCA diagnosis confirmed by microscopy showing CPP crystals within