Conclusions: Fatigue, depression and anxiety have a greater influence on QoL in SLE than disease activity and damage. Of these, fatigue appears to be the most important variable which influences QoL.

REFERENCES:

- [1] Schmeding A, Schneider M. Fatigue, health-related quality of life and other patient-reported outcomes in systemic lupus erythematosus. *Best Pract Res Clin Rheumatol.* 2013;27(3):363–375.
- [2] Maruish ME (Ed). *NEW SF36v2 User Guide, 3rd Edition.*; 2011.

Acknowledgements: Mr. John Michael Raj, Department of Biostatistics, St. John's National Academy of Health Sciences, Bengaluru

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2018-eular.6467

AB0555 IS HIGHER FITNESS ASSOCIATED WITH BETTER HEALTH-RELATED QUALITY OF LIFE IN WOMEN WITH SYSTEMIC LUPUS ERYTHEMATOSUS?

B. Gavilán-Carrera¹, J. García da Silva², J.A. Vargas-Hitos³, P. Morillas-de-Laguna¹, A. Rosales-Castillo³, L.M. Sáez-Urán³, C. Montalbán-Méndez³, A. Soriano-Maldonado^{1,4}. ¹Department of Physical Education and Sport, ²Department of Personality, Evaluation and Psychological Treatment, University of Granada; ³Systemic Autoimmune Diseases Unit, Department of Internal Medicine, "Virgen de las Nieves" Hospital, Granada; ⁴Department of Education, University of Almería, Almería, Spain

Background: Patients with systemic lupus erythematosus (SLE) present reduced health-related quality of life (HRQoL).¹ As an adjunct to traditional medical therapy, higher levels of physical fitness seem to be associated with better symptomatology and HRQoL.² However, a comprehensive characterisation of the association of different components of physical fitness with the different dimensions of HRQoL in this population is lacking and would provide valuable information for the design of prospective studies.

Objectives: The aim of this study was to investigate the association of different physical fitness components (flexibility, muscle strength and cardiorespiratory fitness (CRF) with the dimensions of HRQoL.

Methods: This cross-sectional study included a total of 70 women with SLE (age 42.5; SD 13.9). The back scratch test, 30 s chair stand test and 6 min walk test (n=49) (from the Senior Fitness Test battery) as well as the handgrip strength test were used to assess physical fitness. The median value of each fitness test [flexibility (0.125 cm), upper muscle strength (24.2 kg), lower muscle strength (15 repetitions) and CRF (575 metres)] was used to consider high ($\geq$ median value) or low ($<$ median value) fitness. HRQoL was assessed through the 36-item Short-Form Health Survey (SF-36). Partial correlation was used to examine the association of physical fitness components with all dimensions of the SF-36. For those dimensions significantly associated with fitness, analyses of covariance compared the differences in HRQoL between patients with low and high fitness level. All analyses were controlled for age, body mass index, depression and fatigue.

Results: Flexibility was correlated with physical function, social functioning and physical component (r_{partial} between -0.26 and 0.28 ; all, $p<0.05$). Patients with a high level of flexibility only presented better physical function than those with a low level ($p=0.03$). Upper and lower muscle strength were correlated with physical function, physical role, bodily pain and physical component (r_{partial} between 0.28 and 0.50 ; all, $p<0.05$) with differences between groups of low and high muscle strength (upper and lower body) in all those dimensions ($p<0.05$). CRF was correlated with physical function, bodily pain and physical component (r_{partial} between 0.39 and 0.65 ; all $p<0.01$), with differences in physical function and physical component between groups of low and high CRF ($p<0.05$).

Conclusions: The main findings of the present study suggest that flexibility, muscular strength and cardiorespiratory fitness are associated with different dimensions of HRQoL, particularly with those related to self-reported physical health, in women with systemic lupus erythematosus. Future prospective research should assess the predictive role of fitness regarding disease activity and function.

REFERENCES:

- [1] Chaigne B, et al. *BMC Immunology.* 2017 Mar. 18:17.
- [2] Balsamo S, et al. *BMC Musculoskeletal Disorders.* 2013 Sept (8):14, 263.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2018-eular.6611

AB0556

THE DIAGNOSTIC AND PREDICTIVE SIGNIFICANCE OF NAILFOLD VIDEO CAPILLAROSCOPY IN CTD PATIENTS WITH PULMONARY ARTERIAL HYPERTENSION

B. Jiang, X. Xu, K. Zhou, C. Wang, L. Geng. *The Affiliated Drum Tower Hospital of Nanjing University Medical School, Nanjing, China*

Background: Pulmonary arterial hypertension (PAH) is a serious complication of connective tissue diseases (CTD), defined as an elevation in mean pulmonary artery pressure 25 mmHg at rest. The 1 year, 3 year survival rates of CTD-PAH are only 69% and 30% respectively. Therefore, to early prediction and diagnosis of PAH is very critical for prolonging survival of CTD patients. Nailfold video capillaroscopy (NVC), is an imaging technique for the study of microcirculation. Until now, capillary abnormalities in systemic sclerosis (SSc) are well documented, but rare study focus on the association between the abnormality of capillary with PAH in CTD patients.

Objectives: Through estimating the correlation of the abnormality of nailfold capillary with pulmonary arterial hypertension (PAH) and other organs injury to find out its importance on early diagnosis and prediction for PAH in the patients with connective tissue disease (CTD).

Methods: Nailfold video capillaroscopy (NVC) was performed using a videocapillaroscope to test the fingers capillaries of 123 hospitalised CTD patients with and without PAH. The parameters of NVC were estimated including: the inter, outer and top loop diameter, loop number, misshaped or giant capillaries, flowing speed, haemorrhage, erythrocyte aggregation, et al. The organ involvement like interstitial lung disease (ILD) and cardiovascular complications were also considered. Mann-Whitney test was used to estimate the difference.

Results: 1) The number of capillary loops in patients with SSc was much lower, but morphologic score, liquid score, and total score were all higher than other CTD patients (Tab). 2) The loop number was significantly decreased but abnormal scores of morphology, liquid, diameter, misshaped loop number, flowing speed, erythrocyte aggregation were increased in CTD-PAH patients especially in SLE-PAH compared with non-PAH patients (Fig). 3) ILD relates the abnormality of peripheral circulation such like capillary malformation, decreased loop number, diameter and blood flow, increased haemorrhage, but the influence of PAP may be larger than ILD. 4) Down-regulated loop number, increased morphology, liquid and total score, and abnormal diameter, malformation, erythrocyte aggregation were found in CTD patients with cardiovascular complications, especially in CTD-PAH patients and SLE patients. 5) Age doesn't directly influence the capillaries of CTD patients, but in young and middle age patients, the difference between PAH and non-PAH groups were more evident.

Abstract AB0556 – Table 1. Comparison of number of nail fold capillaries in patients with various CTD

disease	SSc	SLE	SS	PM/DM	RA
loop number	4.688 ± 1.301	5.915 $\pm 1.512^{***}$	5.875 $\pm 1.495^{**}$	5.717 $\pm 1.49^{**}$	6.417 $\pm 1.429^{**}$
morphologic score	4.117 ± 0.293	2.389 $\pm 0.342^{**}$	2.378 $\pm 0.376^{***}$	2.293 $\pm 0.346^{***}$	2.5 $\pm 1.253^{*}$
liquid score	1.942 ± 0.402	0.5277 $\pm 0.101^{***}$	0.6541 $\pm 0.111^{***}$	0.7367 $\pm 0.125^{**}$	0.5333 $\pm 0.217^{*}$
total score	8.321 ± 0.874	4.628 $\pm 0.612^{***}$	4.297 $\pm 0.569^{***}$	4.54 $\pm 0.572^{***}$	4.633 $\pm 1.972^{***}$

Abstract AB0556 – Figure 1. Loop number and other indexes of nail fold capillaries in patients with various CTD and CTD-PAH with non-PAH CTD

Conclusions: In all CTD, the capillaries in SSc patients are most severely damaged. The disease of capillaries in CTD patients correlates with PAH, ILD and cardiovascular complications. So, NVC could be a predictive detection method for PAH and cardiopulmonary disease in CTD patients.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2018-eular.3187

AB0557

HAEMATOLOGICAL INVOLVEMENT (CYTOPENIA) AT THE TIME OF THE DIAGNOSIS IS ASSOCIATED WITH LESS SEVERE OCULAR INVOLVEMENT IN PATIENTS WITH PRIMARY SJOGREN SYNDROME

C.G. Buzatu, S. Daia-Iliescu, I. Saulescu, D. Zaharia, C. Purice, T. Gudu, A. Borangiu, L. Groseanu, V. Bojinca, F. Berghia, A. Balanescu, D. Predeteanu, R. Ionescu, D. Opris-Belinski. *Internal Medicine and Rheumatology, "SFANTA MARIA" Clinical Hospital, Carol Davila University of Medicine and Pharmacy, Bucharest, Romania*

Background: In patients with primary Sjogren Syndrome (pSS), haematological involvement – autoimmune cytopenia, might be present at the time of the diagnosis or can develop in time after the characteristic glandular involvement. ^(1,2)

Objectives: The objective of the study is to evaluate the correlation between glandular involvement (ocular) and presence of cytopenia in patients diagnosed with pSS.

Methods: A retrospective analysis was performed on a cohort of patients diagnosed with primary Sjogren Syndrome under surveillance in one Rheumatology Centre between 2009 and 2016. The documented cases have been diagnosed according to the 2002 American-European Consensus group classification criteria, the 2012 ACR criteria or 2016 ACR/EULAR Classification Criteria for pSS. The EULAR Sjogren's Syndrome Disease Activity Index (ESSDAI) was calculated for all patients. Ocular assessment and follow-up were performed in collaboration with the same ophthalmologist. The data was analysed using Windows Excel/SPSS20.0.

Results: 30 female patients diagnosed with pSS were included in the study. The mean age at the time of diagnosis was 52.1 years±SD 9.1. The ESSDAI was calculated for all patients at baseline: 5 (17%) patients presented high disease activity (ESSDAI >14), 14 (46%) patients moderate disease activity (5≤ESSDAI≤13) and 11 (37%) patients low disease activity (ESSDAI <5). The domain weight for glandular involvement when calculating ESSDAI is fairly low (2),¹ so in the studied group there wasn't obtained a statistically significant correlation between ocular involvement and disease activity as evaluated by ESSDAI.

In the clinical case series, Spearman's rank correlation coefficient between haematological (autoimmune cytopenia), and biological markers (hypocomplementemia) and ocular involvement were calculated. A strong negative correlation was found between autoimmune cytopenia and glandular manifestations (ocular involvement-xerophthalmia) ($r=-0.60$; $p<0.05$). Another strong negative correlation was obtained between hypocomplementemia and severe ocular involvement (corneal ulceration) ($r=-0.59$, $p<0.05$), respectively.

Conclusions: Patients diagnosed with primary Sjogren Syndrome that presented at disease's onset cytopenia and hypocomplementemia had a less severe ocular involvement.

REFERENCES:

- [1] Seror R, Theander E, Brun JG, et al, Validation of EULAR primary Sjögren's syndrome disease activity (ESSDAI) and patient indexes (ESSPRI) *Annals of the Rheumatic Diseases* 2015;74:859–866.
- [2] Shiboski CH, Shiboski SC, Seror R, Criswell LA, Labetoulle M, Lietman TM, Rasmussen A, Scofield H, Vitali C, Bowman SJ, Mariette X, International Sjögren's Syndrome Criteria Working Group, *Ann Rheum Dis*. 2017 Jan;76(1):9–16.
- [3] Koh JH, Lee J, Chung SH, Kwok SK, Park SH, Relation of Autoimmune Cytopenia to Glandular and Systemic Manifestations in Primary Sjögren Syndrome: Analysis of 113 Korean Patients, *J Rheumatol*. 2015 Oct;42(10):1817–24.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2018-eular.7189

AB0558

ARE REGULATORY T CELL LEVELS DIFFERENT IN ACTIVE AND INACTIVE SLE PATIENTS?

H. Üsküdar Teke¹, D. Üsküdar Cansu², C. Bal³, C. Korkmaz². ¹Internal Medicine, Hematology; ²Internal Medicine, Rheumatology; ³Biostatistics, Eskişehir Osmangazi University, Eskişehir, Turkey

Background: The subgroups of T helper cells and regulatory T cells (T-reg) are involved in the pathogenesis of systemic lupus erythematosus (SLE). T-reg cells

suppress immune response to autoantigens and prevent autoimmune diseases (AID). Although there are studies suggesting that Treg cells are reduced during the active period of SLE, there are also studies claiming reversal.

Objectives: To determine whether there is a difference between T-reg and Th-17 cell levels in active and inactive SLE patients and to demonstrate the effects of these cells on disease course.

Methods: 21 SLE patients without active infection were included. Erythrocyte sedimentation rate, CRP, C3, C4, anti-ds-DNA levels, SLEDAI values were recorded in terms of organ and system involvement, hemogram results, disease activation. SLEDAI ≥6 were considered active disease. Concurrent peripheral blood Th17 and T-reg levels were studied. 15 healthy control subjects were included in the study.

Results: 19 of the patients were women, the mean age was 37.3±12.9 years and mean duration of the disease was 7.9±1.29 years. Hematologic involvement was present in 15 (71.4%) of the patients, renal involvement in 17 (81%) and joint involvement in 7 (42.9%). Both Treg cells and CD4 +IL17+cell levels were significantly higher in SLE group than HC group in terms of CD4 +CD25+, CD4 +FOXP3+T reg, CD4 +CD25+FOXP3+T reg ($p=0.011$, $p=0.001$, $p=0.001$ and $p=0.040$, respectively). There was no significant difference between active ($n=12$) and inactive ($n=9$) SLE patients in terms of Th17 and T-reg levels. However, in the inactive period, the levels of CD4 +FOXP3+T reg and CD4 +CD25+FOXP3+Treg cells tended to increase compared to the active period.

Conclusions: This study showed the tendency of increasing in Treg cells in the inactive period. This may be related to the modification of immunosuppressive drugs. It may be more appropriate to perform similar studies before and after treatment.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2018-eular.6151

AB0559

THE PREVALENCE OF NON-CRITERIA ANTI-PHOSPHOLIPID ANTIBODIES IN ANTI-PHOSPHOLIPID SYNDROME

C. Li¹, L. Zhu¹, Z. Wang¹, Y. Sun², R. Mu¹, Z. Li¹. ¹Peking University People's Hospital, Beijing; ²266 Hospital of Chinese People's Liberation Army, Chengde, China

Background: Antiphospholipid syndrome (APS) is an autoimmune disease characterised mainly by arterial and/or venous thrombosis, recurrent pregnancy morbidity, with the presence of a variety of heterogeneous circulating antiphospholipid antibodies. However, there are a group of APS patients with persistently negative antiphospholipid antibodies. It is necessary to validate new specific antibodies to better recognition of these APS patients.

Objectives: To explore the clinical significance of non-criteria anti-phospholipid antibodies in a large cohort of Chinese patients with anti-phospholipid syndrome (APS).

Methods: Serum samples were obtained from 214 APS patients, 122 disease control including systemic lupus erythematosus, sjogren syndrome, ankylosing spondylitis, rheumatoid arthritis, osteoarthritis and 50 healthy control. Antiphosphatidylserine(IgA/IgG/IgM) (aPS), anti-phosphatidylethanolamine (IgA/IgG/IgM) (aPE), anti-prothrombin-antibodies (IgA/IgG/IgM) (aPT), anti-annexin V-antibodies (IgA/IgG/IgM) (aAnxV), anti-phospholipid-antibody(IgG/IgM)(aPL), anti-oxLDL antibody(IgG/IgM)(aoxLDL) were tested by ELISA kits(HUMAN Diagnostics Products, Germany) and IgG/IgM APL were tested by ELISA(Louisville APL Diagnostic, USA). The Chi-square (χ^2) test was used to examine the difference of frequencies of antibodies in APS patients and patients with other diseases. Spearman correlation analysis was performed to investigate the relationship between aPS/PT and other clinical/laboratory parameters.

Results: The prevalence of aPS IgG, aPS IgM, aPT, aPE, aAnxV, aPL, aoxLDL and IgG APL, IgM APL were 44.8%, 22.4%, 46.3%, 9.8%, 17.8%, 44.4%, 22.4%, 45.3% and 22.9%. The specificity of the antibodies were 82.6%, 95.3%, 96.0%, 99.4%, 95.3%, 92.4%, 87.2%, 95.3% and 94.8%. The highest positive predictive values of these antibodies were aPE. aPE, aPS IgM and APL IgM were associated with thrombotic events and oxLDL, aPE, aPL, aPS IgG, aPS IgM, aPT, aoxLDL, APL IgG and APL IgM were correlated with anti-cardiolipin antibody (ACL). aPL, aPS IgG, aPS IgM, aPT, APL IgG and APL IgM were associated with β 2-GP1 antibody. APL IgG, APL IgM and PS IgG were the highest prevalence in both ACL and lupus anticoagulant (LAC) negative patients. aPT has the highest prevalence in ACL and β 2-GP1 negative patients and APL IgG and aPS IgG were the highest two antibodies in LAC and β 2-GP1 negative patients and aPT has the highest prevalence in seronegative APS patients.

Conclusions: Non-criteria aPLs have a good diagnostic value in APS and were associated with thrombotic events.

REFERENCES:

- [1] Hughes GR, Khamashta MA (2003) Seronegative antiphospholipid syndrome. *Ann Rheum Dis* 62:1127